

Arylation of chloroanthraquinones by surprisingly facile Suzuki–Miyaura cross-coupling reactions

Thies Thiemann^{a,f*}, Yasuko Tanaka^b, Jesus Iniesta^c, H. Tresa Varghese^d and C. Yohannan Pannicker^e

^aInterdisciplinary Graduate School of Engineering Sciences and ^bInstitute of Materials Chemistry and Engineering, Kyushu University, 6-1, Kasuga-koh-en, Kasuga-shi, Fukuoka 816-8580, Japan

^cDepartment of Physical Chemistry, University of Alicante, E-03080 Alicante, Spain

^dDepartment of Physics, Fatima Mata National College, Kollam-1, India

^eDepartment of Physics, TKM College of Arts and Science, Kollam-5, Kerala, India

^fPresent address: Department of Chemistry, Faculty of Science, United Arab Emirates University, PO Box 17551, Al Ain, United Arab Emirates.

Chloroanthraquinones were found to undergo facile Suzuki–cross coupling with substituted phenyl boronic acids using a commercial catalyst $\text{Pd}(\text{PPh}_3)_4$ and with $\text{Pd}(\text{PPh}_3)_4$ prepared *in situ* from $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ and PPh_3 .

Keywords: chloroarenes, anthraquinones, Suzuki–Miyaura cross-coupling, tetrakis(triphenylphosphino)palladium

The Suzuki–Miyaura cross coupling reaction of aryl- and alkylboronic acids with bromo- and iodoarenes is a well established reaction. As bromo- and iodoarenes are expensive in comparison to chloroarenes, the recent focus has been on the development of new catalysts and reaction protocols for the Suzuki reaction with chloroarenes. A number of catalysts had been proposed by Fu,¹ Buchwald² and others^{3–8} in the last 20 years, in which the triphenylphosphine ligand of the original tetrakis[triphenylphosphino]palladium (0) was replaced by other ligands such as tris(*tert*-butyl)phosphine. With the advent of ligandless catalysts for Suzuki–Miyaura reactions, a number of highly reactive nanopalladium catalysts⁹ have been developed for the coupling of chloroarenes. It has also been found that ligandless palladium catalysts can be used with aryl tetrafluoroborates in coupling reactions with aryl halides.¹⁰ However, in the synthesis of arylated anthraquinones, as part of our search for highly conjugated π -systems, we have found that the use of specially prepared palladium catalysts are not always necessary in the reaction of chloroarenes. Thus, chloroanthraquinones easily undergo Suzuki cross coupling reactions with arylboronic acids in the presence of the original, commercially available $\text{Pd}(\text{PPh}_3)_4$.

Arylated anthraquinones **1** (Fig. 1) have attracted attention because of their interesting physical organic properties,^{11,12} due to the interaction of the attached aryl groups with the π -system of the anthraquinone core as revealed by the UV and luminescence^{13,14} spectra, the redox behaviour of the molecules,^{12,15} and in the NMR shift values. Specifically, the interaction of the substituents with the C=O function of the anthraquinones has been subjected to investigation.¹¹ In practical applications, arylated anthraquinones have also been used as stabilisers of light-modulating fluids such as those containing liquid polybenzyltoluenes.¹⁶ Our interest in these molecules lies in the study of their electrochemical behaviour. Having prepared a number of arylated anthraquinones from bromo substituted anthraquinones by Suzuki–Miyaura coupling reactions,¹⁷ we pursued the question of whether it was possible to carry out the transformation with the respective chloro substituted anthraquinones.

Results and discussion

First, we had to synthesise chlorinated anthraquinones for which a number of synthetic procedures are known. Thus, 1,4-dichloroanthraquinone (**1a**) can be synthesised from





Figure 1

1-hydroxyanthraquinone by chlorination and subsequent treatment of the 1-chloro-4-hydroxyanthraquinone with PCl_5 .¹⁸ 1,4-Dichloroanthraquinone (**1a**) can also be prepared by acylation of 1,4-dichlorobenzene (**3**) with phthaloyl chloride (**2**) and subsequent heating of the 2-(2,5-dichlorobenzoyl) benzoic acid intermediate with H_2SO_4 (Scheme 1),¹⁹ or by treatment of 9,10-dihydroxy-2,3-dihydro-1,4-anthraquinone with PCl_5 .^{20,21}




In the present work, we chose to prepare chloroanthraquinones from 2,5-dichlorothiophene (**4a**). This was oxidised by *m*-CPBA to 2,5-dichlorothiophene-*S*-oxide and reacted *in situ* with naphthoquinones (Scheme 2). The use of thiophene *S*-oxides *in situ* as dienes in Diels Alder reactions has been found to be an efficient route to multifunctionalised arenes.^{22–24} This is especially true for thiophenes with electron donating substituents. Thiophenes with electron-withdrawing substituents^{23,25,26} such as chloro- or bromothiophenes **4** are much more difficult to oxidise. Nevertheless, at elevated temperatures it was possible to isolate chloroanthraquinones when chlorothiophenes were reacted with *m*-CPBA in the presence of naphthoquinones. Even in cases where the halothiophene *S*-oxides were oxidised further to halothiophene *S,S*-dioxides prior to undergoing a Diels Alder reaction, cycloaddition reactions were expected to proceed with the halothiophene *S,S*-dioxides equally well. Electron poor thiophene *S,S*-dioxides have been found to undergo cycloaddition reactions readily.^{27,28} Thus, under the present conditions, halothiophene *S,S*-dioxides may also make a contribution to the reaction.

The resultant chloroanthraquinones **1a–e** were subjected to Suzuki–Miyaura cross coupling reactions with a number of different arylboronic acids. Tetrakis(triphenylphosphine) palladium(0) [$\text{Pd}(\text{PPh}_3)_4$] was used as catalyst and the reactions were run in a biphasic system of 1,2-dimethoxyethane (DME) and an aqueous sodium carbonate solution. The arylated anthraquinones were isolated in good yield. In case of

* Correspondent. E-mail: thies@uaeu.ac.ae



Scheme 1



Scheme 2


1,4-dibromo-5,8-dichloroanthraquinone (1c), all of the halo-groups exchanged with equal ease, so that a 5,8-diaryl-1,4-dichloroanthraquinone intermediate could not be isolated. Also, 1-bromo-5,8-dichloro-4-hydroxyanthraquinone (1e), which is a side-product in the oxidative cycloaddition of 2,5-dibromothiophene (4b) to 1,4-dichlorobenzoquinone (7), was transformed easily to 1-hydroxy-4,5,8-tris(4-methoxyphenyl)anthraquinone (9k) by a Suzuki–Miyaura cross-coupling reaction with 4-methoxyphenylboronic acid (8a). The use of Pd(PPh₃)₄, prepared *in situ* from Pd(PPh₃)₂Cl₂ and PPh₃, worked equally well.

In conclusion, we have shown that chloroanthraquinones undergo Suzuki–Miyaura coupling with the commercially available tetrakis(triphenylphosphine)palladium(0)[Pd(PPh₃)₄] and with the combination of Pd(PPh₃)₂Cl₂ and triphenylphosphine.

Experimental

CAUTION: Working with meta chloroperoxybenzoic acid at elevated temperatures is hazardous. The reactions should be carried out in a well-ventilated hood. Protection against an explosion should be set up. (The authors have not experienced any difficulties with these reactions. These measures are precautions.)

IR spectra were measured with JASCO IR-700, JASCO FTIR-6300 and Nippon Denshi JIR-AQ20M machines. ¹H and ¹³C NMR spectra were recorded with a JEOL EX-270 spectrometer (¹H at 270 MHz and ¹³C at 67.8 MHz). The chemical shifts are relative to TMS (solvent CDCl₃, unless otherwise noted). Mass spectra were measured with a JMS-01-SG-2 spectrometer [electron impact mode (EI), 70 eV or fast atom bombardment (FAB)]. Column chromatography was carried out on Wakogel C-300.



Scheme 3

Table 1

Starting Material	Product	Starting Material	Product
1a: R = H; 1b: R = CH₃	9a: R = H, R¹ = CH₃ (83%) 9b: R = H, R¹ = C₆H₁₁ (85%) 9c: R = H, R¹ = C₇H₁₅ (81%) 9d: R = H, R¹ = C₁₀H₂₁ (80%) 9e: R = CH₃, R¹ = C₅H₁₁ (78%)	1e	9j (70%)
1a	9f: R¹ = CHO; Z = <i>p</i>-C₆H₄-CHO (45%) 9g: R¹ = CH₃; Z = Cl (53%) 9h: R¹ = H; Z = C₆H₅ (68%)	1f	9k: R = CH₃ (56%) 9l: R = C₂H₅ (59%)
1d	9i (72%)	9g	9m (71%)

2,3-Dimethyl-5,8-naphthoquinone (**5b**) was prepared by the cycloaddition of 2,3-dimethylbuta-1,3-diene to *p*-benzoquinone under EuCl₃ catalysis (96 h, ClCH₂CH₂Cl, rt).²⁹ This reaction has been reported to proceed under YbCl₃ catalysis and we have found that it also proceeds in the presence of EuCl₃. With subsequent base catalysed enolisation³⁰ of the 4a,5,8,8a-tetrahydro-6,7-dimethyl-1,4-naphthoquinone which was formed and oxidation of the 6,7-dimethyl-5,8-dihydronaphthalene-1,4-diol (Ag₂O, Na₂SO₄, benzene)³¹ to 6,7-dimethyl-5,8-dihydro-1,4-naphthoquinone, which in the last step was dehydrogenated (DDQ, benzene, reflux). [Here, we have used a biphasic system of 4N aq. NaOH and ether under ultrasonication. The enolisation has been reported to also go very well in the presence of triethylamine (Et₃N) or in the presence of acids such as HCl]. We used Ag₂O in benzene as the oxidation agent as described for the oxidation of other hydroquinones to quinones. The reaction gives quantitative yields of 6,7-dimethyl-5,8-dihydro-1,4-naphthoquinone, when carried out at rt. Specifically for the transformation of 6,7-dimethyl-

5,8-dihydronaphthalene-1,4-diol to 6,7-dimethyl-5,8-dihydro-1,4-naphthoquinone, the use of MnO₂ in acetone has been described.³¹ *p*-Methoxyphenylboronic acid (**8a**) (TCI), *p*-formylphenylboronic acid (**8e**) (TCI), phenylboronic acid (**8g**) (TCI) and *p*-tolylboronic acid (**8f**) (Aldrich) were acquired commercially. *p*-Ethoxy-phenylboronic acids, **8c** and the homologous *p*-alkoxyphenylboronic acids **8b-d** were prepared from the corresponding *p*-alkoxy-bromobenzenes (a. *n*-BuLi, B(OR)₃, THF; b. HCl).³²

1,4-Dichloroanthraquinone (1a):¹⁸ *general procedure A:* A stirred solution of 2,5-dichlorothiophene (**4a**) (640 mg, 4.16 mmol) and 1,4-naphthoquinone (517 mg, 3.47 mmol) in CHCl₃ (20 mL) at 75 °C was treated with *m*-CPBA (70 w%, 4.76 g) in small portions. After 48 h, the mixture was cooled and poured into an aq. sat. Na₂CO₃ solution. After the mixture had been stirred for 15 min. at rt, it was extracted with chloroform (3 × 25 mL). The organic phase was dried over anhydrous MgSO₄ and concentrated *in vacuo*. The residue was subjected to column chromatography on silica gel (hexane/ether/

CHCl₃) to give **1a** (425 mg, 37%) as yellow needles, m.p. 187°C; (Found: M⁺, 275.9748. C₁₄H₆O₂³⁵Cl₂ requires M⁺, 275.9745). δ_{H} (270 MHz, CDCl₃) 7.68 (2H, s), 7.77–7.81 (2H, m), 8.17–8.21 (2H, m); δ_{C} (67.8 MHz, CDCl₃) 126.9 (2C, CH), 132.1 (2C, C_{quat}), 133.6 (2C, C_{quat}), 134.0 (2C, C_{quat}), 134.2 (2C, CH), 137.2 (2C, CH), 181.6 (2C, C_{quat}, CO). MS (EI, 70 eV) *m/z* (%) = 276 (M⁺) (100).

1,4-Dichloro-6,7-dimethylantraquinone (1b); general procedure A: Pale yellow needles, m.p. 194°C; (Found: M⁺, 304.0060. C₁₆H₁₀O₂³⁵Cl₂ requires M⁺, 304.0058). δ_{H} (270 MHz, CDCl₃) 2.42 (6H, s, 2 CH₃), 7.64 (2H, s), 7.92 (2H, s); δ_{C} (67.8 MHz, CDCl₃) 20.3 (2C, CH₃), 127.7 (2C, CH), 129.6 (2C, C_{quat}), 131.6 (2C, C_{quat}), 133.8 (2C, C_{quat}), 140.0 (2C, CH), 144.3 (2C, C_{quat}), 181.7 (2C, C_{quat}, CO); MS (EI, 70 eV) *m/z* (%) = 304 (M⁺) (100).

1,4-Dibromo-5,8-dichloroanthraquinone (1c); general procedure A: Colourless solid, m.p. 210°C; (Found: M⁺, 433.7930. C₁₄H₄O₂³⁵Cl₂⁷⁹Br₂ requires M⁺, 433.7933). δ_{H} (270 MHz, CDCl₃) 7.60 (2H, s), 7.72 (2H, s); MS (EI, 70 eV) *m/z* (%) = 438 (3.3), 436 (9.2), 434 (9.6), 432 (3.9), 149 (34), 58 (100).

1,4-Bis(4-methoxyphenyl)anthraquinone (9a); general procedure B: A solution of **1a** (245 mg, 0.89 mmol), 4-methoxyphenylboronic acid (430 mg, 2.83 mmol), Pd(PPh₃)₄ (46 mg, 4.0 × 10⁻⁵ mol) [or Pd(PPh₃)₂Cl₂ (30 mg, 4.10⁻⁵ mol) and triphenylphosphine (30 mg, 0.11 mmol)] in a solvent mixture of DME (10 mL) and aq. Na₂CO₃ (2.32 g Na₂CO₃ in 15 mL H₂O, 6 mL) was kept at 70°C for 18 h in an inert atmosphere. The solution was then cooled and poured into water (25 mL) and extracted with chloroform (3 × 15 mL). The combined organic phase was dried over anhydrous MgSO₄ and concentrated *in vacuo*. Column chromatography of the residue on silica gel (hexane/CHCl₃/ether 3:1:1) gave **9a** (310 mg, 83%) as orange needles; m.p. 231°C; (Found: M⁺, 420.1367. C₂₈H₂₀O₄ requires M⁺, 420.1362). ν_{max} (KBr/cm⁻¹) 2950, 2867, 1671, 1604, 1279, 1242, 1204, 1023, 971, 857, 830; δ_{H} (270 MHz, CDCl₃) 3.89 (6H, s, 2 OCH₃), 7.00 (4H, d, ³*J* = 8.6 Hz), 7.26 (4H, d, ³*J* = 8.6 Hz), 7.53 (2H, s), 7.68–7.72 (2H, m), 8.06–8.09 (2H, m); δ_{C} (67.8 MHz, CDCl₃) 55.2 (2C, OCH₃), 113.7 (4C, CH), 126.7 (2C, CH), 129.3 (4C, CH), 133.7 (2C, CH), 132.9 (2C, C_{quat}), 134.1 (2C, C_{quat}), 134.5 (2C, C_{quat}), 136.6 (2C, CH), 143.6 (2C, C_{quat}), 158.9 (2C, C_{quat}), 184.3 (2C, C_{quat}, CO); MS (EI, 70 eV) *m/z* (%) 420 (M⁺) (100), 389 (32), 333 (18), 313 (13), 276 (17). UV-Vis (CH₃CN, nm) λ_{max} 253 (59610), 271 (sh, 23890), 313 (13280).

1,4-Bis(4-pentoxypheyl)anthraquinone (9b); general procedure B: Rose-coloured solid; m.p. 202°C; (Found: MH⁺, 533.2697. C₃₆H₃₇O₄ requires MH⁺, 533.2692 [FAB]). ν_{max} (KBr/cm⁻¹) 2953, 2869, 1604, 1515, 1385, 1375, 1320, 1281, 859, 828; δ_{H} (270 MHz, CDCl₃) 0.96 (6H, t, ³*J* = 7.0 Hz), 1.46 (8H, m), 1.84 (4H, m), 4.04 (4H, t, ³*J* = 6.5 Hz), 6.99 (4H, d, ³*J* = 8.6 Hz), 7.24 (4H, d, ³*J* = 8.6 Hz), 7.53 (2H, s), 7.67–7.71 (2H, m), 8.06–8.09 (2H, m); δ_{C} (67.8 MHz, CDCl₃) 14.1 (2C, CH₃), 22.5 (2C, CH₂), 28.3 (2C, CH₂), 29.1 (2C, CH₂), 67.9 (2C, OCH₂), 114.2 (4C, CH), 126.7 (2C, C_{quat}), 129.2 (4C, CH), 132.9 (2C, C_{quat}), 133.6 (2C, CH), 134.2 (2C, C_{quat}), 136.6 (2C, CH), 143.6 (2C, C_{quat}), 158.5 (2C, C_{quat}), 184.3 (2C, C_{quat}, CO); MS (FAB, 3-nitrobenzyl alcohol) *m/z* (%) 533 (MH⁺) (100), 391 (36). Calcd for C₃₆H₃₆O₄: C, 81.17; H, 6.81. Found: C, 80.79; H, 6.77%.

1,4-Bis(4-heptoxypheyl)anthraquinone (9c); general procedure B: Rose-coloured solid, m.p. 205°C; (Found: MH⁺, 589.3313. C₄₀H₄₅O₄ requires MH⁺, 589.3318 [FAB]). ν_{max} (KBr/cm⁻¹) 2950, 2867, 1673, 1604, 1515, 1318, 1280, 1244, 1206, 1023, 970, 858, 828; δ_{H} (270 MHz, CDCl₃) 0.92 (6H, t, ³*J* = 6.8 Hz, 2 CH₃), 1.34–1.56 (16H, m), 1.83 (4H, m), 4.03 (4H, t, ³*J* = 6.7 Hz, 2 OCH₂), 6.98 (4H, d, ³*J* = 8.6 Hz), 7.24 (4H, d, ³*J* = 8.6 Hz), 7.53 (2H, s), 7.67–7.70 (2H, m), 8.06–8.09 (2H, m); δ_{C} (67.8 MHz, CDCl₃) 14.1 (2C, CH₃), 22.6 (2C, CH₂), 26.1 (2C, CH₂), 29.1 (2C, CH₂), 29.4 (2C, CH₂), 31.8 (2C, CH₂), 67.9 (2C, OCH₂), 114.1 (4C, CH), 126.7 (2C, C_{quat}), 129.2 (4C, CH), 132.9 (2C, C_{quat}), 133.6 (2C, CH), 134.2 (2C, CH), 134.3 (2C, C_{quat}), 136.6 (2C, CH), 143.6 (2C, C_{quat}), 158.5 (2C, C_{quat}), 184.3 (2C, C_{quat}, CO); MS (FAB, 3-nitrobenzyl alcohol) *m/z* (%) 589 (MH⁺) (30).

1,4-Bis(4-decyloxyphenyl)anthraquinone (9d); general procedure B: Orange-yellow solid, m.p. 197°C; (Found: MH⁺, 673.4277. C₄₆H₅₇O₄ requires MH⁺, 673.4275 [FAB]). ν_{max} (KBr/cm⁻¹) 2950, 2867, 1673, 1604, 1515, 1280, 1243, 1204, 1023, 971, 857, 828; δ_{H} (270 MHz, CDCl₃) 0.89 (6H, t, ³*J* = 7.0 Hz, 2 CH₃), 1.29–1.55 (28H, m), 1.78–1.86 (4H, m), 4.03 (4H, t, ³*J* = 6.5 Hz, 2 OCH₂), 6.98 (4H, d, ³*J* = 8.6 Hz), 7.24 (4H, d, ³*J* = 8.6 Hz), 7.53 (2H, s), 7.67–7.71 (2H, m), 8.06–8.09 (2H, m); δ_{C} (67.8 MHz, CDCl₃) 14.1 (2C, CH₃), 22.7 (2C, CH₂), 26.1 (2C, CH₂), 29.3 (2C, CH₂), 29.4 (2C, CH₂), 29.5 (2C, CH₂), 29.6 (2C, CH₂), 29.7 (2C, CH₂), 31.9 (2C, CH₂), 67.9 (2C, OCH₂), 114.2 (4C, CH), 126.7 (2C, C_{quat}), 129.2 (4C, CH), 132.9 (2C, C_{quat}), 133.6 (2C, CH), 134.2 (2C, CH), 134.3 (2C, C_{quat}), 136.6 (2C, CH), 143.6 (2C, C_{quat}), 158.5 (2C, C_{quat}), 184.3 (2C, C_{quat}, CO); MS (FAB, 3-nitrobenzyl alcohol) *m/z* (%) = 673 (MH⁺) (8), 672 (M⁺) (10).

1,4-Bis(4-pentoxypheyl)-6,7-dimethylantraquinone (9e); general procedure B: Pale orange needles, m.p. 215°C; (Found: MH⁺, 561.3006. C₃₈H₄₁O₄ requires MH⁺, 561.3005 [FAB]). ν_{max} (KBr/cm⁻¹) 2950, 2867, 1673, 1604, 1515, 1458, 1389, 1373, 1320, 1279, 1242, 1204, 1023, 971, 857, 828; δ_{H} (270 MHz, CDCl₃) 0.96 (6H, t, ³*J* = 7.0 Hz, 2 CH₃), 1.81–1.86 (4H, m), 1.41–1.54 (8H, m), 2.35 (6H, s, 2 CH₃), 4.03 (4H, t, ³*J* = 6.5 Hz), 6.98 (4H, d, ³*J* = 8.6 Hz), 7.24 (4H, d, ³*J* = 8.6 Hz), 7.51 (2H, s), 7.83 (2H, s); δ_{C} (67.8 MHz, CDCl₃) 14.1 (2C), 20.2 (2C), 22.5 (2C), 28.3 (2C), 29.1 (2C), 67.9 (2C), 114.1 (4C), 127.6 (2C), 129.2 (4C), 132.1 (2C), 133.0 (2C), 134.4 (2C), 136.4 (2C), 143.4 (2C), 143.6 (2C), 158.4 (2C), 184.5 (2C); MS (FAB, 3-nitrobenzyl alcohol) *m/z* (%) = 561 (MH⁺) (28), 419 (7.4), 326 (10). Calcd for C₃₈H₄₀O₄: C, 81.40; H, 7.19. Found: C, 80.87; H, 7.14%.

1,4-Bis(4-formylphenyl)anthraquinone (9f); general procedure B: Yellow solid; m.p. 243°C; (Found: M⁺, 417.1130. C₂₈H₁₇O₄ requires M⁺, 417.1127). ν_{max} (KBr/cm⁻¹) 3068, 1687, 1677, 1602, 1380, 1319, 1259, 1211, 1168, 1091, 958, 827, 730; δ_{H} (270 MHz, CDCl₃) 7.51 (4H, d, ³*J* = 8.1 Hz), 7.60 (2H, s), 7.73–7.76 (2H, m), 8.02 (4H, d, ³*J* = 8.1 Hz), 8.06–8.09 (2H, m), 10.1 (2H, s, 2 CHO); δ_{C} (67.8 MHz, CDCl₃) 127.0 (2C, CH), 128.5 (4C, CH), 129.7 (4C, CH), 132.6 (2C, C_{quat}), 133.5 (2C, C_{quat}), 134.2 (2C, CH), 135.2 (2C, C_{quat}), 135.9 (2C, CH), 143.3 (2C, C_{quat}), 148.6 (2C, C_{quat}), 183.5 (2C, CO), 191.92 (2C, CHO); MS (FAB, 3-nitrobenzyl alcohol) *m/z* (%) 417 (MH⁺) (3.4).

1-Chloro-4-(4-methylphenyl)anthraquinone (9g); general procedure B: Yellow solid; m.p. 224°C; (Found: 333.0685. C₂₁H₁₄O₂³⁵Cl requires M⁺, 333.0682). ν_{max} (KBr/cm⁻¹) 3001, 1672; δ_{H} (270 MHz, CDCl₃) 2.45 (3H, s, CH₃), 7.15 (2H, d, ³*J* = 8.4 Hz), 7.27 (2H, d, ³*J* = 8.4 Hz), 7.46 (1H, d, ³*J* = 8.1 Hz), 7.69–7.80 (2H, m), 7.75 (1H, d, ³*J* = 8.1 Hz), 8.03–8.06 (1H, m), 8.22–8.26 (1H, m); δ_{C} (67.8 MHz, CDCl₃) 21.3 (CH₃), 126.7 (CH), 126.9 (CH), 127.7 (2C, CH), 127.9 (C_{quat}), 129.0 (2C, CH), 131.2 (C_{quat}), 133.7 (C_{quat}), 133.9 (CH), 134.0 (CH), 134.1 (C_{quat}), 134.3 (C_{quat}), 136.4 (CH), 137.1 (C_{quat}), 137.3 (CH), 138.5 (C_{quat}), 143.8 (C_{quat}), 182.6 (C_{quat}, CO), 183.3 (C_{quat}, CO); MS (FAB, 3-nitrobenzyl alcohol) *m/z* (%) 333 (MH⁺) (4.4). Calcd for C₂₁H₁₄O₂Cl: C, 75.79; H, 3.94. Found: C, 75.86; H, 3.99%.

1,4-Diphenylanthraquinone (9h);^{12,33} general procedure B: Yellow solid; (Found: MH⁺, 361.1232. C₂₆H₁₇O₂ requires MH⁺, 361.1229 [FAB]). ν_{max} (KBr/cm⁻¹) 3002, 1673, 758; δ_{H} (270 MHz, CDCl₃) 7.29–7.35 (4H, m), 7.43–7.48 (6H, m), 7.56 (2H, s), 7.66–7.71 (2H, m), 8.05–8.08 (2H, m); δ_{C} (67.8 MHz, CDCl₃) 126.8 (2C, CH), 127.2 (2C, CH), 127.9 (4C, CH), 128.2 (4C, CH), 132.7 (2C, C_{quat}), 133.7 (2C, CH), 134.0 (2C, C_{quat}), 136.4 (2C, CH), 142.3 (2C, C_{quat}), 144.1 (2C, C_{quat}), 184.0 (2C, C_{quat}, CO); MS (FAB, 3-nitrobenzyl alcohol) *m/z* (%) 361 (MH⁺) (5.6). UV-Vis (CH₃CN, nm) λ_{max} 253 (36370), 269 (sh, 19190), 288 (sh, 7320).

1,4,5,8-Tetrakis(4-methoxyphenyl)anthraquinone (9i); general procedure B: Using 4-methoxyphenylboronic acid (860 mg, 5.66 mmol) and Pd(PPh₃)₄ (60 mg, 5.2 × 10⁻⁵ mol) [or Pd(PPh₃)₂Cl₂ (40 mg, 5.2 × 10⁻⁵ mol) and triphenylphosphine (37 mg, 0.14 mmol)], in DME (15 mL) and aq. Na₂CO₃ (9 mL): pale orange solid; m.p. 251°C; (Found: MH⁺, 633.2286. C₄₂H₃₃O₆ requires MH⁺, 633.2277 [FAB]). ν_{max} (KBr/cm⁻¹) 1671, 1608, 1575, 1514, 1460, 1328, 1291, 1247, 1029, 831; δ_{H} (270 MHz, CDCl₃) 3.84 (12H, s, 4 OCH₃), 6.85 (8H, d, ³*J* = 8.4 Hz), 7.21 (8H, d, ³*J* = 8.4 Hz), 7.48 (4H, s); δ_{C} (67.8 MHz, CDCl₃) 55.2 (4C, OCH₃), 113.4 (8C, CH), 130.3 (8C, CH), 131.9 (4C, C_{quat}), 134.5 (4C, CH), 135.5 (4C, C_{quat}), 140.3 (4C, C_{quat}), 159.0 (4C, C_{quat}), 188.4 (2C, C_{quat}, CO); MS (FAB, 3-nitrobenzyl alcohol) *m/z* (%) = 633 (MH⁺) (1.0).

1,4,5,8-Tetrakis(4-ethoxyphenyl)anthraquinone (9j); general procedure: Analogous to the synthesis of **9i**: yellow orange solid; m.p. 307°C; (Found: MH⁺, 689.2910. C₄₆H₄₁O₆ requires MH⁺, 689.2903 [FAB]). ν_{max} (KBr/cm⁻¹) 3002, 1673, 1245, 1205, 831; δ_{H} (270 MHz, CDCl₃) 1.42 (12H, t, ³*J* = 7.0 Hz, 4 CH₃), 4.06 (8H, q, ³*J* = 7.0 Hz, 4 OCH₂), 6.82 (8H, d, ³*J* = 8.6 Hz), 7.19 (8H, d, ³*J* = 8.6 Hz), 7.47 (4H, s); δ_{C} (67.8 MHz, CDCl₃) 14.9 (4C, CH₃), 63.4 (4C, OCH₂), 113.9 (8C, CH), 130.3 (8C, CH), 131.7 (4C, C_{quat}), 134.4 (4C, CH), 135.6 (4C, C_{quat}), 140.3 (4C, C_{quat}), 158.4 (4C, C_{quat}), 188.5 (2C, C_{quat}, CO); MS (FAB, 3-nitrobenzyl alcohol) *m/z* (%) = 689 (MH⁺) (27).

1-Hydroxy-4,5,8-tris(4-methoxyphenyl)anthraquinone (9k); general procedure B: Analogous to the synthesis of **9i**: reddish solid; m.p. 238°C; (Found: MH⁺, 543.1805. C₃₅H₂₇O₆ requires MH⁺, 543.1808 [FAB]). ν_{max} (KBr/cm⁻¹) 3430 (bs, OH), 3054, 3002, 1673, 1515, 1461, 1295, 1243, 1205, 1179, 1031, 827, 754; δ_{H} (270 MHz, CDCl₃) 3.82 (3H, s, OCH₃), 3.83 (3H, s, OCH₃), 3.90 (3H, s, OCH₃), 6.84 (2H, d, ³*J* = 8.6 Hz), 6.87 (2H, d, ³*J* = 8.6 Hz), 7.00 (2H, d, ³*J* = 8.6 Hz), 7.14–7.29 (7H, m), 7.46 (1H, d, ³*J* = 7.8 Hz), 7.47 (1H, d, ³*J* = 8.4 Hz), 7.56 (1H, d, ³*J* = 7.8 Hz), 12.21 (s, 1H, OH); δ_{C} (67.8 MHz, CDCl₃) 55.2 (2C, OCH₃), 55.3 (OCH₃), 113.6 (6C,

CH), 117.1 (C_{quat}), 122.1 (CH), 129.4 (2C, CH), 129.9 (2C, CH), 130.2 (2C, CH), 131.0 (C_{quat}), 132.2 (C_{quat}), 132.5 (C_{quat}), 134.0 (C_{quat}), 134.1 (C_{quat}), 134.2 (C_{quat}), 136.0 (CH), 136.4 (CH), 136.8 (C_{quat}), 139.5 (CH), 142.0 (C_{quat}), 142.9 (C_{quat}), 158.8 (C_{quat}), 159.0 (2C, C_{quat}), 161.0 (C_{quat}), 188.0 (C_{quat}, CO), 189.5 (C_{quat}, CO); MS (FAB, 3-nitrobenzyl alcohol) *m/z* (%) = 543 (MH⁺) (1.4).

1-Hydroxy-4,5,8-tris(4-ethoxyphenyl)anthraquinone (9i); general procedure B: Analogous to the synthesis of **9i**: red-orange needles; m.p. 259°C; (Found: M⁺, 585.2270. C₃₈H₃₃O₆ requires M⁺, 585.2277). $\nu_{\max}$ (KBr/cm⁻¹) 3430 (bs, OH), 3054, 3002, 1671, 1513, 1460, 1295, 1243, 1205, 1178, 1030, 825, 752; δ_{H} (270 MHz, CDCl₃) 1.44 (6H, t, ³J = 7.0 Hz, 2 CH₃), 1.48 (3H, t, ³J = 7.0 Hz, CH₃), 4.05 (2H, q, ³J = 7.0 Hz, OCH₂), 4.06 (2H, q, ³J = 7.0 Hz, OCH₂), 4.13 (2H, q, ³J = 7.0 Hz, OCH₂), 6.83 (2H, d, ³J = 8.6 Hz), 6.85 (2H, d, ³J = 8.6 Hz), 6.99 (2H, d, ³J = 8.6 Hz), 7.14–7.29 (7H, m), 7.46 (1H, d, ³J = 7.8 Hz), 7.47 (1H, d, ³J = 8.4 Hz), 7.56 (1H, d, ³J = 7.8 Hz), 12.20 (s, 1H, OH); δ_{C} (67.8 MHz, CDCl₃) 14.8 (2C, CH₃), 14.9 (CH₃), 63.3 (OCH₂), 63.4 (2C, OCH₂), 114.1 (6C, CH), 117.1 (C_{quat}), 122.1 (CH), 129.4 (2C, CH), 129.9 (2C, CH), 130.2 (2C, CH), 131.0 (C_{quat}), 132.2 (C_{quat}), 132.5 (C_{quat}), 134.0 (C_{quat}), 134.1 (C_{quat}), 134.2 (C_{quat}), 136.0 (CH), 136.4 (CH), 136.8 (C_{quat}), 139.5 (CH), 142.0 (C_{quat}), 142.9 (C_{quat}), 158.2 (C_{quat}), 159.0 (2C, C_{quat}), 161.0 (C_{quat}), 188.0 (C_{quat}, CO), 189.5 (C_{quat}, CO); MS (FAB, 3-nitrobenzyl alcohol) *m/z* (%) = 585 (MH⁺) (14), 539 (3.5).

1-(4-Ethoxyphenyl)-4(4-methylphenyl)anthraquinone (9m); general procedure B: Yellow-orange solid, m.p. 247°C; (Found: MH⁺, 419.1651. C₂₉H₂₃O₃ requires MH⁺, 419.1647). δ_{H} (270 MHz, CDCl₃) 1.47 (3H, t, ³J = 7.0 Hz, CH₃), 2.46 (3H, s, CH₃), 4.12 (2H, q, ³J = 7.0 Hz, OCH₂), 6.99 (2H, d, ³J = 8.6 Hz), 7.21 (2H, d, ³J = 7.6 Hz), 7.27 (2H, d, ³J = 8.6 Hz), 7.28 (2H, d, ³J = 7.6 Hz), 7.51 (1H, d, ³J = 8.1 Hz), 7.55 (1H, d, ³J = 8.1 Hz), 7.67–7.70 (2H, m), 8.06–8.09 (2H, m); δ_{C} (67.8 MHz, CDCl₃) 14.9 (CH₃), 21.4 (CH₃), 63.4 (OCH₂), 114.1 (2C, CH), 127.8 (2C, CH), 128.9 (2C, CH), 129.2 (2C, CH), 132.8 (C_{quat}), 132.9 (C_{quat}), 133.6 (2C, CH), 134.1 (C_{quat}), 134.2 (C_{quat}), 134.3 (C_{quat}), 136.4 (CH), 136.6 (CH), 136.8 (C_{quat}), 139.4 (C_{quat}), 143.7 (C_{quat}), 143.8 (C_{quat}), 158.3 (C_{quat}), 184.2 (C_{quat}, CO), 184.3 (C_{quat}, CO); FAB, 3-nitrobenzyl alcohol) *m/z* (%) 419 (MH⁺) (100). Calcd for C₂₉H₂₃O₃: C, 83.23; H, 5.30. Found: C, 83.15; H, 5.27%.

The research was supported partially by the Global Centre of Excellence on New Carbon Resources, Kyushu University, Japan.

Received 29 September 2009; accepted 9 November 2009

Paper 09/0804 doi: 10.3184/030823409X12586584303880

Published online: 8 December 2009

References

- G.C. Fu, *Pure Appl. Chem.*, 2002, **74**, 33.
- J.P. Wolfe, R.A. Singer, B.H. Yang and S.L. Buchwald, *J. Am. Chem. Soc.*, 1999, **121**, 9550.
- C. Zhang and M.L. Trudell, *Tetrahedron Lett.*, 2000, **41**, 595.
- C. Zhang, J. Huang, M.L. Trudell and S.P. Nolan, *J. Org. Chem.*, 1999, **64**, 3804.
- M.G. Andreu, A. Zapf and M. Beller, *Chem. Commun.*, 2000, 2475.
- G.Y. Li, *Angew. Chem. Int. Ed. Engl.*, 2001, **40**, 1513.
- L. Botella and C. Najera, *Angew. Chem. Int. Ed. Engl.*, 2002, **41**, 179.
- J. Zhang, L. Zhao, M. Song, T.C.W. Mak and Y. Wu, *J. Organomet. Chem.*, 2006, **691**, 1301.
- B.M. Choudary, S. Madhi, N.S. Chowdari, M.L. Kantam and B. Sreedhar, *J. Am. Chem. Soc.*, 2002, **124**, 14127.
- G. Lu, R. Franzen, Q. Zhang and Y. Xu, *Tetrahedron Lett.*, 2005, **46**, 4253.
- J. Josien and N. Fuson, *Bull. Soc. Chim. Fr.*, 1952, 389.
- J.E. Gautrot, P. Hodge, D. Cupertino and M. Helliwell, *New J. Chem.*, 2006, **30**, 1801.
- N.A. Shcheglova, D.N. Shigorin and N.S. Dokunikhin, *Zh. Fiz. Khim.*, 1966, **40**, 1048; *Chem. Abstr.*, 1966, **65**, 9958b.
- N.A. Shcheglova, D.N. Shigorin and N.S. Dokunikhin, *Zh. Fiz. Khim.*, 1965, **39**, 3039; *Chem. Abstr.*, 1965, **64**, 12058f.
- H.D.H. Showalter, E.M. Berman, J.L. Johnson and W.E. Hunter, *Tetrahedron Lett.*, 1985, **26**, 157.
- D.A. Orser (General Electric Company, Schenectady, N.Y., USA) US Pat. 3764549 (Oct. 9, 1973); *Chem. Abstr.*, 1973, **79**, 141431j.
- T. Thiemann, Y. Tanaka and J. Iniesta, *Molecules*, 2009, **14**, 1013.
- V.V. Kozlov, *Zh. Obshch. Khim.*, 1947, **17**, 289; *Chem. Abstr.*, **42**, 2581.
- H. Schoenhagen and R. Schmitz (Bayer AG), DE 3513981 (Oct. 30, 1986); *Chem. Abstr.*, 1987, **106**, 20008.
- P.A. Krapcho and Z. Getahun, *Synth. Commun.*, 1985, **15**, 907.
- F. Ihlan, D.S. Tyson and M.A. Meador, *Org. Lett.*, 2006, **8**, 577.
- Y.Q. Li, T. Thiemann, T. Sawada, S. Mataka and M. Tashiro, *J. Org. Chem.*, 1997, **62**, 7926.
- for reviews, see: T. Thiemann and K. Gopal Dongol, *J. Chem. Res. (S)*, 2002, 303; (M), 2002, 701.
- T. Thiemann, D.J. Walton, A.O. Brett, J. Iniesta, F. Marken and Y.Q. Li, *ARKIVOC*, 2009, **ix**, 96.
- T. Thiemann, M.L. Sae Melo, A.S. Campos Neves, Y. Li, S. Mataka, M. Tashiro, U. Geißler and D. Walton, *J. Chem. Res. (S)*, 1998, 346.
- see also: K. Torssell, *Acta Chem. Scand., Ser. B*, 1976, **30**, 353.
- M.S. Raasch, *J. Org. Chem.*, 1980, **45**, 856.
- D.P.G. Hamon and P.R. Spurr, *J. Chem. Soc., Chem. Commun.*, 1982, 373.
- X. Fang, B.P. Warner and J.G. Watkin, *Synth. Commun.*, 2000, **30**, 2669.
- M.P. Fernandez, B. Gonzalez, M. Pardo and J.L. Soto, *Anal. Quim.*, 1994, **90**, 477 (for Et₃N) and see ref. 31.
- S. Kotha and E. Manivannan, *Ind. J. Chem., Sect. B*, 2002, **41B**, 808 (for HCl).
- cf., K. Fuchibe and T. Akiyama, *J. Am. Chem. Soc.*, 2006, **128**, 1434.
- N.I. Ganushchak, A.V. Dombrovskii and O.A. Vislobitskaya, *Zh. Obshch. Khim.*, 1963, **33**, 2532.