

CONVENIENT SYNTHESIS OF CYCLOHEXA[*a*]PYRROLO[2,1-*b*][3]BENZAZEPINE, A CEPHALOTAXUS ALKALOID ANALOGUE


Hitoshi Tanaka,^{*a} Mitsunobu Doi,^b Hiroshi Shimizu,^c and Hideo Etoh^d

^aFaculty of Pharmacy, Meijo University, Yagoto, Tempaku-ku, Nagoya 468-8503, Japan, ^bOsaka University of Pharmaceutical Sciences, 4-20-1, Nasahara, Takatsuki, Osaka, 569-1094, Japan, ^cGifu Pharmaceutical University, 6-1 Mitahora-higashi 5 chome, Gifu 502-8585, Japan, ^dFaculty of Agriculture, Shizuoka University, Shizuoka 422-8529, Japan

Abstract — Irradiation of *N*-(3-hydroxy-4-methoxyphenethyl)-3-(2-iodo-5-methoxyphenyl)propionamide (**11**) in methanol in the presence of sodium hydroxide furnished 3-hydroxy-2,12-dimethoxy-6,7,9,10-tetrahydrodibenz[*d,f*]azecin-8(5*H*)-one (**14**) which was successfully led to cephalotaxine analogue (**17**) bearing pyrrolo[2,1-*b*][3]benzazepine skeleton by reduction with borane followed by Birch reduction and subsequent acidic treatment. The structure of **17** was established by means of X-Ray crystallography.


We previously reported that bromophenolic amides underwent photochemical cyclization reaction to form easily dibenz[*d,f*]azecine skeleton,^{1,2} which was subjected to reduction with borane followed by Birch reduction and subsequent acidic treatment to give homoerythrinan derivative.² The dibenz[*d,f*]azecine base (**4**) might be the important intermediate in the biosynthesis of homoerythrinan alkaloids and cephalotaxan alkaloids³ (Scheme 1) whose ester derivatives, harringtonine (**2**) and homoharringtonine (**3**), have potent antileukemic activity.⁴ From view of spectrum of the biological activity and unique structural features, there are many reports on the synthesis of cephalotaxine (**1**)⁵ and its analogues, homocephalotaxine⁶ and aza-cephalotaxine derivative,⁷ whose pivotal skeleton is pyrrolo[2,1-*b*][3]benzazepine. In this paper, we wish to describe a novel route for the synthesis of a cephalotaxine analogue (**17**) having the pyrrolo[2,1-*b*][3]benzazepine skeleton from a dibenz[*d,f*]azecine base (**15**) using Birch reduction followed by acidic treatment.

First, the dibenz[*d,f*]azecine base (**15**) was prepared as shown in Scheme 2. The condensation reaction of β-(2-iodo-5-methoxyphenyl)propionic acid (**8**)⁸ and 3-benzyloxy-4-methoxyphenethylamine (**9**) in refluxing xylene gave **10** in 73% yield. Treatment of **10** with methanesulfonic acid and thioanisole in




trifluoroacetic acid according to the procedure of Kiso and coworkers⁹ afforded the iodophenolic amide (**11**) in 69% yield. Irradiation of **11** in methanol in the presence of sodium hydroxide with a 100 W high pressure mercury lamp and subsequent chromatographic separation provided three products **12**, **13** and **14** in 7, 1 and 22% yields, respectively. The slower moving product (**14**) was recrystallized from EtOH and showed a molecular ion peak at m/z 327 in the MS and the IR spectrum indicated the presence of hydroxyl and amide groups. The ^1H NMR spectrum revealed two singlet signals at δ 6.54 and 6.86 due to H-1 and H-4 together with ABX type signals (δ 6.80, 6.88 and 7.01). Thus, the structure of this product is clearly represented as formula (**14**) and it indicates that the photochemical coupling reaction occurred at *para* position to hydroxyl group of **11**. The second slower moving product (**13**) had the same molecular formula and functional groups (hydroxyl and amide groups) as those of **14** from the observation of the MS and IR spectrum of **13**. In the ^1H NMR spectrum, *ortho* coupling signals (δ 6.71 and 6.87) were observed along with ABX type signals (δ 6.55, 6.66 and 6.75). Therefore, this compound was assigned to **13**. The fast moving product (**12**) had the same hydroxyl and amide groups as those of starting material (**11**) by the MS and IR spectrum of **12**. The ^1H NMR spectrum showed a similar signal pattern to that of **11** except for the aromatic proton regions, in which four proton signals (δ 6.73-6.78 (3H) and 7.20 (1H)) were observed together with the typical ABX pattern signals (δ 6.56, 6.69 and 6.76). The MS of this compound indicated the molecular ion peak at m/z 329, which corresponds to the lack of an iodine atom of **11**. Therefore, this compound was assigned to **12**. Reduction of **14** with borane gave the dibenz[*d,f*]azecine (**15**) in good yield.

The dibenz[*d,f*]azecine (**15**) was treated with sodium in liquid ammonia and methanol to afford reduction product (**16**) in 34% yield. In the ^1H NMR spectrum of **16**, an olefinic proton appeared at δ 4.69. Finally,



Scheme 2

20% hydrochloric acid treatment of **16** in dioxane furnished cyclization product (**17**) in 26% yield. The IR spectrum of **17** showed carbonyl absorption band at 1700 cm^{-1} and in the ^1H NMR spectrum, the olefinic proton and methoxyl group signals disappeared and the methine proton signal due to H-5 appeared newly as doublet of doublet at δ 3.44. The structural determination of compound (**17**) was finally carried out by X-Ray analysis (see EXPERIMENTAL). The molecular structure of compound (**17**) is illustrated in Figure 1. Thus, we have achieved a convenient synthesis of cephalotaxine analogue (**17**) *via* Birch reduction of the dibenz[*d,f*]azecine (**15**) followed by acidic treatment.

Figure 1. The X-Ray structure of **17**

EXPERIMENTAL

General. The instruments used for this study were as follows: a JASCO IR-810 spectrophotometer (for IR spectra); a JEOL JMS-D 300 spectrometer (for MS and HRMS spectra); a JEOL JNM-A 400 spectrometer (for NMR spectra using tetramethylsilane as an internal standard). Column chromatography was carried out with Merck silica gel 60 (70-230 mesh).

N-(3-Benzyloxy-4-methoxyphenethyl)-3-(2-iodo-5-methoxyphenyl)propionamide (**10**)

A mixture of **8**⁸ (8.8 g, 28.77 mmol) and **9** (7.4 g, 28.79 mmol) in xylene (200 mL) was refluxed for 3 h. The mixture was evaporated to dryness and the residue was dissolved in CHCl₃. The CHCl₃ solution was washed in turn with 5% HCl, 5% NaOH and H₂O, dried over Na₂SO₄, and then evaporated. The crude solid was recrystallized from EtOH to give colorless prisms (**10**) (11.4 g, 73%), mp 150 °C. IR (KBr) $\nu_{\max}$ cm⁻¹: 3300 (NH), 1640 (CO), 1590. ¹H NMR (CDCl₃): δ 2.34 (2H, *t*, *J*=7.4 Hz, CH₂CH₂CO), 2.65 (2H, *t*, *J*=6.7 Hz, CH₂CH₂NH), 2.97 (2H, *t*, *J*=7.4 Hz, CH₂CH₂CO), 3.41 (2H, *q*, *J*=6.7 Hz, CH₂CH₂NH), 3.75, 3.87 (6H, 2 *x* *s*, 2 *x* OCH₃), 5.13 (2H, *s*, OCH₂Ph), 5.33 (1H, *br s*, NH), 6.51 (1H, *dd*, *J*=3.0, 8.7 Hz, H-4), 6.66 (1H, *dd*, *J*=2.0, 8.1 Hz, H-6'), 6.70 (1H, *d*, *J*=2.0 Hz, H-2'), 6.82 (1H, *d*, *J*=8.1 Hz, H-5'), 6.83 (1H, *d*, *J*=3.0 Hz, H-6), 7.28-7.53 (5H, *m*, 5 *x* arom. H), 7.64 (1H, *d*, *J*=8.7 Hz, H-3). MS *m/z*: 545 [M]⁺, 418, 328, 306, 268, 247, 240. *Anal.* Calcd for C₂₆H₂₈NO₄I: C, 57.26; H, 5.17; N, 2.57. Found: C, 57.16; H, 5.12; N, 2.44.

N-(3-Hydroxy-4-methoxyphenethyl)-3-(2-iodo-5-methoxyphenyl)propionamide (**11**)

A mixture of **10** (8 g, 14.68 mmol), PhSCH₃ (4.3 mL, 36.63 mmol) and CH₃SO₃H (2 mL, 30.82 mmol) in TFA (14 mL) was stirred under a nitrogen atmosphere at 0 °C for 2 h. The reaction mixture was poured into ice water and extracted with CHCl₃. The CHCl₃ extract was washed with H₂O, dried over Na₂SO₄, and evaporated. The crude solid was recrystallized from benzene to give colorless needles (**11**) (4.6 g, 69%), mp 122 °C. IR (CHCl₃) $\nu_{\max}$ cm⁻¹: 3550 (OH), 3450 (NH), 1670 (CO), 1595. ¹H NMR (CDCl₃): δ 2.38 (2H, *t*, *J*=7.4 Hz, CH₂CH₂CO), 2.66 (2H, *t*, *J*=6.7 Hz, CH₂CH₂NH), 2.99 (2H, *t*, *J*=7.4 Hz, CH₂CH₂CO), 3.46 (2H, *q*, *J*=6.7 Hz, CH₂CH₂NH), 3.76, 3.87 (6H, 2 *x* *s*, 2 *x* OCH₃), 5.41 (1H, *br s*, NH or OH), 5.65 (1H, *br s*, OH or NH), 6.51 (1H, *dd*, *J*=3.0, 8.7 Hz, H-4), 6.59 (1H, *dd*, *J*=2.0, 8.1 Hz, H-6'), 6.72 (1H, *d*, *J*=2.0 Hz, H-2'), 6.77 (1H, *d*, *J*=8.1 Hz, H-5'), 6.83 (1H, *d*, *J*=3.0 Hz, H-6), 7.64 (1H, *d*, *J*=8.7 Hz, H-3). MS *m/z*: 455 [M]⁺, 328, 306, 289, 247. *Anal.* Calcd for C₁₉H₂₂NO₄I: C, 50.12; H, 4.87; N, 3.08. Found: C, 50.02; H, 4.85; N, 3.06.

Irradiation of **11** (Formation of **12**, **13** and **14**)

A mixture of **11** (800 mg, 1.76 mmol) and NaOH (650 mg, 16.25 mmol) in MeOH (250 mL) was irradiated with a 100 W high pressure mercury lamp at rt for 2 h under a nitrogen atmosphere. The same procedure was run eleven times and then the reaction mixture was combined. After the whole mixture was evaporated, the residue was acidified with 5% HCl and extracted with CHCl₃. The CHCl₃ extract was

washed with H₂O, dried over Na₂SO₄, and evaporated to give a brown oil. The oil was chromatographed on a silica gel column and eluted with a mixture of CHCl₃ and acetone (5:1). The starting material (**11**)(414 mg) was first eluted. Then the second (**12**)(452.7 mg, 7% based on consumed starting material), the third (**13**)(63.9 mg, 1%), and the fourth substance (**14**)(1.3 g, 22%) were in turn eluted.

12: A crude solid was recrystallized from EtOH to give colorless prisms, mp 114 °C. TLC (silica gel/CHCl₃-acetone (5:1), R_f=0.38). IR (CHCl₃) $\nu_{\max}$ cm⁻¹: 3550 (OH), 3450 (NH), 1660 (CO), 1600. ¹H NMR (CDCl₃): δ 2.41 (2H, *t*, *J*=7.4 Hz, CH₂CH₂CO), 2.64 (2H, *t*, *J*=6.7 Hz, CH₂CH₂NH), 2.93 (2H, *t*, *J*=7.4 Hz, CH₂CH₂CO), 3.44 (2H, *q*, *J*=6.7 Hz, CH₂CH₂NH), 3.79, 3.87 (6H, 2 *x s*, 2 *x* OCH₃), 5.31 (1H, *br s*, NH or OH), 5.62 (1H, *s*, OH or NH), 6.56 (1H, *dd*, *J*=2.0, 8.1 Hz, H-6'), 6.69 (1H, *d*, *J*=2.0 Hz, H-2'), 6.76 (1H, *d*, *J*=8.1 Hz, H-5'), 6.73-6.78 (3H, *m*, 3 *x* arom. H), 7.20 (1H, *m*, arom. H). MS *m/z*: 329 [M]⁺, 293, 281, 255, 243, 231, 219, 205. HRMS calcd for C₁₉H₂₃NO₄: 329.1626. Found: 329.1616.

13: A crude solid was recrystallized from benzene to give colorless prisms, mp 224 °C. TLC (silica gel/CHCl₃-acetone (5:1), R_f=0.30). IR (CHCl₃) $\nu_{\max}$ cm⁻¹: 3550 (OH), 3450 (NH), 1665 (CO), 1605. ¹H NMR (benzene-*d*₆)¹⁰: δ 3.18, 3.36 (6H, 2 *x s*, 2 *x* OCH₃), 3.78 (1H, *br s*, NH or OH), 5.60 (1H, *s*, OH or NH), 6.55 (1H, *d*, *J*=8.4 Hz, H-14), 6.66 (1H, *dd*, *J*=3.0, 8.4 Hz, H-13), 6.71 (1H, *d*, *J*=8.4 Hz, H-4 or H-3), 6.75 (1H, *d*, *J*=3.0 Hz, H-11), 6.87 (1H, *d*, *J*=8.4 Hz, H-3 or H-4). MS *m/z*: 327 [M]⁺, 282, 268, 265, 255, 253, 239, 223. HRMS calcd for C₁₉H₂₁NO₄: 327.1469. Found: 327.1460.

14: A crude solid was recrystallized from EtOH to give colorless prisms, mp 228 °C. TLC (silica gel/CHCl₃-acetone (5:1), R_f=0.14). IR (CHCl₃) $\nu_{\max}$ cm⁻¹: 3550 (OH), 3450 (NH), 1670 (CO), 1610. ¹H NMR (CDCl₃): δ 3.81, 3.85 (6H, 2 *x s*, 2 *x* OCH₃), 4.51 (1H, *br s*, NH or OH), 5.58 (1H, *br s*, OH or NH), 6.54 (1H, *s*, H-1 or H-4), 6.80 (1H, *dd*, *J*=3.0, 8.4 Hz, H-13), 6.86 (1H, *s*, H-4 or H-1), 6.88 (1H, *d*, *J*=3.0 Hz, H-11), 7.01 (1H, *d*, *J*=8.4 Hz, H-14). MS *m/z*: 327 [M]⁺, 268, 255, 239, 237, 227, 225, 223, 211, 209. *Anal.* Calcd for C₁₉H₂₁NO₄·1/2H₂O: C, 67.84; H, 6.59; N, 4.16. Found: C, 67.69; H, 6.25; N, 4.20.

3-Hydroxy-2,12-dimethoxy-5,6,7,8,9,10-hexahydrodibenz[*d,f*]azecine (**15**)

BH₃-THF complex (1 M solution)(8 mL) was added dropwise to a stirred solution of **14** (233 mg, 0.71 mmol) in anhydrous THF (15 mL) at 0 °C under a nitrogen atmosphere. The mixture was continued to stir at rt overnight and then the excess BH₃ was decomposed by addition of EtOH. The reaction mixture was evaporated to give a brown oil that was heated with a mixture of MeOH (1 mL) and conc. HCl (1 mL) at 60 °C for 30 min. After the acidic solution was evaporated to dryness, the residue was made alkaline with 5% NH₄OH and extracted with CHCl₃. The CHCl₃ extract was washed with H₂O, dried over Na₂SO₄, and evaporated to give a crude solid. The solid was recrystallized from EtOH to afford colorless prisms (**15**)(192 mg, 86%), mp 227 °C. IR (CHCl₃) $\nu_{\max}$ cm⁻¹: 3550 (OH), 1605. ¹H NMR (CDCl₃): δ 3.78, 3.84 (6H, 2 *x s*, 2 *x* OCH₃), 6.52 (1H, *s*, H-1 or H-4), 6.71 (1H, *s*, H-4 or H-1), 6.77 (1H, *dd*, *J*=3.0, 8.4 Hz, H-13), 6.81 (1H, *d*, *J*=3.0 Hz, H-11), 6.99 (1H, *d*, *J*=8.4 Hz, H-14). MS *m/z*: 313 [M]⁺, 298, 281, 255.

HRMS calcd for $C_{19}H_{23}NO_3$: 313.1677. Found: 313.1670.

3-Hydroxy-2,12-dimethoxy-5,6,7,8,9,10,11,14-octahydrodibenz[*d,f*]azecine (16)

Sodium (3 g, 130 matom) was added to a stirred mixture of **15** (182 mg, 0.58 mmol), anhydrous MeOH (8 mL), anhydrous THF (9 mL), and liquid ammonia (40 mL) at $-70\text{ }^{\circ}\text{C}$ over a period of 2h. After cautious addition of MeOH, H_2O and NH_4Cl , the ammonia was allowed to evaporate off and the reaction mixture was extracted with $CHCl_3$. The $CHCl_3$ extract was washed with H_2O , dried over Na_2SO_4 , and evaporated to afford a crude solid. The solid was recrystallized from MeOH to give colorless needles (**16**) (63 mg, 34%), mp $229\text{ }^{\circ}\text{C}$. IR ($CHCl_3$) $\nu_{max}\text{ cm}^{-1}$: 3550 (OH), 1510. 1H NMR ($CDCl_3$): δ 3.60, 3.83 (6H, 2 x s, 2 x OCH_3), 4.69 (1H, *br s*, H-13), 6.47 (1H, *s*, H-1 or H-4), 6.70 (1H, *s*, H-4 or H-1). MS m/z : 315 $[M]^+$, 300, 284, 258, 243, 239, 231, 227, 225, 211. HRMS calcd for $C_{19}H_{25}NO_3$: 315.1833. Found: 315.1838.

17-Hydroxy-16-methoxy-8,9,11,12-tetrahydro-7H-cyclohexa[*a*]pyrrolo[2,1-*b*][3]benzazepin-2-one (17)

A stirred mixture of **16** (45 mg, 0.14 mmol) in dioxane (10 mL) and 20% HCl (0.5 mL) was heated at $60\text{ }^{\circ}\text{C}$ for 20 min. After cooling, the reaction mixture was poured into ice water. The whole mixture was made alkaline with 5% NH_4OH and extract with AcOEt. The AcOEt extract was washed with H_2O , dried over Na_2SO_4 , and evaporated to give a crude solid. The solid was recrystallized from MeOH to afford colorless prisms (**17**) (11.1 mg, 26%), mp $90\text{ }^{\circ}\text{C}$. IR ($CHCl_3$) $\nu_{max}\text{ cm}^{-1}$: 3550 (OH), 1700 (CO), 1600. 1H NMR ($CDCl_3$): δ 1.39 (1H, *m*, H-8 β), 1.43 (1H, *m*, H-7 β), 1.51-1.62 (2H, *m*, H-7 α and H-8 α), 1.91 (1H, *m*, H-4 β), 2.14 (1H, *m*, H-4 α), 2.24 (1H, *d*, $J=13.7\text{ Hz}$, H-1 β), 2.38 (1H, *m*, H-9 α), 2.46 (1H, *ddd*, $J=5.9$, 12.9, 14.4 Hz, H-3 α), 2.53 (1H, *ddd*, $J=2.7$, 5.9, 14.4 Hz, H-3 β), 2.70 (1H, *d*, $J=13.7\text{ Hz}$, H-1 α), 2.72 (1H, *m*, H-9 β), 2.86 (1H, *ddd*, $J=5.1$, 10.0, 15.1 Hz, H-12 β), 3.10 (1H, *ddd*, $J=4.4$, 10.0, 14.7 Hz, H-11 α), 3.25 (1H, *ddd*, $J=4.4$, 9.8, 15.1 Hz, H-12 α), 3.38 (1H, *ddd*, $J=5.1$, 9.8, 14.7 Hz, H-11 β), 3.44 (1H, *dd*, $J=3.2$, 12.5 Hz, H-5), 3.88 (3H, *s*, OCH_3), 5.46 (1H, *br s*, OH), 6.57 (1H, *s*, H-15), 6.74 (1H, *s*, H-18). These signals were assigned by the NOE experiment. MS m/z : 301 $[M]^+$, 286, 283, 281, 258, 244, 231, 216. HRMS calcd for $C_{18}H_{23}NO_3$: 301.1677. Found: 301.1690.

X-Ray Diffraction Study of 17.

The 2737 reflections were measured on Rigaku RU200/AFC5R up to $\theta_{max}=62.1^{\circ}$ by $2\theta-\omega$ scan method. Crystal data --- formula: $C_{18}H_{23}NO_3 \cdot CH_3OH$, $Mr=330.4$, $F(000)=648$, $\mu(\text{Cu K}\alpha)=0.630\text{ mm}^{-1}$, crystal system: monoclinic, space group: $P2_1/c$, $a=8.486(3)\text{ \AA}$, $b=20.361(9)\text{ \AA}$, $c=10.735(3)\text{ \AA}$, $\beta=111.46(2)^{\circ}$, $V=1726.3(11)\text{ \AA}^3$, $Z=4$. The structure was solved by *Crystan G* and was refined by *SHELXL-93*. The non-hydrogen atoms were refined with anisotropic temperature factors and the hydrogen atoms were calculated at geometrically ideal positions. In the difference Fourier map, two peaks were found and were separated *ca.* 1.4 $\text{\AA}$ to each other. These peaks were located at the distance of 2.726 $\text{\AA}$ from the hydroxyl

oxygen (position 17) of compound (**17**) and were considered as a solvent molecule (methanol). In the final cycle of the refinement, the *R* and *wR* values were converged to 0.0676 and 0.1715 for 2386 reflections with over $2\sigma(I)$, respectively. The other parameters are as follows. No. of parameters=226, goodness of fit=1.138, $(\Delta/\sigma)_{\max} < 0.001$, $(\Delta\rho)_{\max}=0.654 \text{ e}\text{\AA}^{-3}$, $(\Delta\rho)_{\min}=-0.411 \text{ e}\text{\AA}^{-3}$.

ACKNOWLEDGMENT

We wish to thank Professor Itsuo Mori at Osaka University of Pharmaceutical Sciences for the valuable help.

REFERENCES AND NOTES

1. H. Tanaka, Y. Takamura, K. Ito, K. Ohira, and M. Shibata, *Chem. Pharm. Bull.*, 1984, **32**, 2063.
2. H. Tanaka, Y. Takamura, M. Shibata, and K. Ito, *Chem. Pharm. Bull.*, 1986, **34**, 24.
3. Y. Tsuda and T. Sano, *The Alkaloids*, Vol. 48, ed. by G. A. Cordell, Academic press, 1996, p. 249.
4. a) L. Huang and Z. Xue, *The Alkaloids*, Vol. 23, ed. by A. Brossi, Academic press, 1984, p. 157; b) J. -Y. Zhou, D. -L. Chen, Z. -S. Shen, and H. P. Koeffler, *Cancer Res.*, 1990, **50**, 2031.
5. Total synthesis of cephalotaxine: a) S. M. Weinreb and J. Auerbach, *J. Am. Chem. Soc.*, 1975, **97**, 2503; b) M. F. Semmelhack, B. P. Chong, R. D. Stauffer, T. D. Rogerson, A. Chong, and L. D. Jones, *J. Am. Chem. Soc.*, 1975, **97**, 2507; c) S. Yasuda, T. Yamada, and M. Hanaoka, *Tetrahedron Lett.*, 1986, **27**, 2023; d) M. E. Kuehne, W. G. Bornmann, W. H. Parsons, T. D. Spitzer, J. F. Blount, and J. Zubieta, *J. Org. Chem.*, 1988, **53**, 3439; e) T. P. Burkholder and P. L. Fuchs, *J. Am. Chem. Soc.*, 1990, **112**, 9601; f) M. Ikeda, M. Okano, K. Kosaka, M. Kido, and H. Ishibashi, *Chem. Pharm. Bull.*, 1993, **41**, 276; g) N. Isono and M. Mori, *J. Org. Chem.*, 1995, **60**, 115; h) X. Lin, R. W. Kavash, and P. S. Mariano, *J. Org. Chem.*, 1996, **61**, 7335; i) L. F. Tietze and H. Schirok, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 1124; j) T. Nagasaka, H. Sato, and S.-I. Saeki, *Tetrahedron: Asymmetry*, 1997, **8**, 191.
6. M. Okano, N. Nishimura, K. Maruyama, K. Kosaka, H. Ishibashi, and M. Ikeda, *Chem. Pharm. Bull.*, 1991, **39**, 3163.
7. M. Nyerges, I. Bitter, I. Kadas, G. Toth, and L. Toke, *Tetrahedron*, 1995, **51**, 11489.
8. D. Papa, E. Schwenk, and E. Klingsberg, *J. Am. Chem. Soc.*, 1950, **72**, 2623.
9. Y. Kiso, H. Isawa, K. Kitagawa, and T. Akita, *Chem. Pharm. Bull.*, 1978, **26**, 2562.
10. Although aromatic proton signals of **13** in CDCl_3 were overlapped, all of those signals in benzene- d_6 were clearly separated.