

Preparation, Characterization, and X-Ray Structure Studies of Tertiary Phosphine Coordinated Molybdenocene

Takashi ITO,* Tadayuki TOKUNAGA, Makoto MINATO, and Takashi NAKAMURA

Department of Materials Chemistry, Faculty of Engineering,

Yokohama National University, 156 Tokiwadai, Hodogaya-ku, Yokohama 240



$\text{Cp}_2\text{MoH}(\text{OTs})$ ($\text{Cp} = \eta\text{-C}_5\text{H}_5$, $\text{Ts} = p\text{-CH}_3\text{C}_6\text{H}_4\text{SO}_2$), which was prepared from Cp_2MoH_2 by its treatment with TsOH in EtOH , reacted with tertiary phosphines and phosphite to give cationic $[\text{Cp}_2\text{MoH}(\text{PR}_3)]^+\text{TsO}^-$ where $\text{R} = \text{Ph}$, Et , OEt , Cy , and Bu^n . Deprotonation of the latter with NaOH in EtOH afforded $\text{Cp}_2\text{Mo}(\text{PR}_3)$; X-ray structure ($\text{R} = \text{Bu}^n$) and some reactions including those with alkyl halides and with H_2 to revert to Cp_2MoH_2 of which were studied.

Recently we reported the diastereoselective reduction of organic carbonyl compounds with the system composed of Cp_2MoH_2 (**1**) ($\text{Cp} = \eta\text{-C}_5\text{H}_5$) and acid (HA) such as RCO_2H , TsOH ($\text{Ts} = p\text{-CH}_3\text{C}_6\text{H}_4\text{SO}_2$), and HCl (Eq. 1).¹⁾



In an effort to convert this reaction into catalytic one, we succeeded in obtaining and characterizing, including X-ray crystal structure analysis, a series of neutral tertiary phosphine derivatives of molybdenocene, $\text{Cp}_2\text{Mo}(\text{PR}_3)$ (**2**). The formation of **2** has so far been suggested by Azevedo *et al.* in the reaction of $[\text{Cp}_2\text{MoH}(\text{PR}_3)]^+\text{PF}_6^-$ with $\text{NaH}^{2)}$ and by Geoffroy *et al.* in the photochemical reaction of **1** with PR_3 in isoocane.³⁾ Development of easy access to the complex of the type **2** and establishment of its structural characterization are important in view of the fact that it behaves as a highly reactive molybdenocene precursor.^{2,3)} In fact, complex **2** ($\text{R} = \text{Bu}^n$) reacted with H_2 under fairly moderate conditions to give **1** (*vide infra*).

Hydridotosylato complex $\text{Cp}_2\text{MoH}(\text{OTs})$ (**3**), which was prepared in almost quantitative yield by the reaction of **1** with equimolar amount of $\text{TsOH} \cdot \text{H}_2\text{O}$ in ethanol at 80°C ,^{4,5)} was allowed to react with tertiary phosphines and phosphite to give cationic phosphine adducts of monohydride (**4**) (Eqs. 2 and 3 and Table 1).



The complex **4** with $\text{PR}_3 = \text{PPh}_3$, which has been prepared by the reaction of $[\text{Cp}_2\text{MoH}_3]^+\text{TsO}^-$ with PPh_3 ,⁴⁾ was also formed in 82% yield by the direct reaction of **1** with TsOH in the presence of PPh_3 and acetone (Eq. 4). In the reaction of $[\text{Cp}_2\text{MoH}_3]^+\text{TsO}^-$ with PEt_3 , the latter worked as a Lewis base to trap dissociated TsOH yielding $[\text{P}(\text{H})\text{Et}_3]^+\text{TsO}^-$ together with **1**.⁴⁾ In contrast, the present route (Eqs. 2 and 3) can afford **4** with PEt_3 ligand successfully although its purification was somewhat difficult.

PCy_3 ligand in **4d** was found to be replaced easily with PPh_3 , which suggests the existence of an equi-


Table 1. Preparation (Eq. 3) and Spectroscopic Properties of Cationic Complexes **4**^{a)}

PR ₃ /mmol		3 /mmol	Solvents/cm ⁻³	Time/h	Yield of 4 /%	Yield of 4 /%	IR /cm ⁻¹ ^{c)}		¹ H NMR/ppm ^{d)}	
							v(Mo-H)	δ(Cp)	δ(Mo-H)	
PPh ₃	(a)	2.68	1.34	EtOH 10	20	58 (94)	1820	4.96 (1.8)	-8.06 (34)	
PEt ₃	(b)	1.26	0.63	THF 10	10	e) (92)	1855	5.11 (1.5)	-8.51 (35)	
P(OEt) ₃	(c)	1.22	0.81	THF 10	4	50 (85)	1865	5.23 (1.8)	-8.69 (39)	
PCy ₃ ^{f)}	(d)	1.76	1.17	THF 10	22	17 (34)	1865	5.15 (1.2)	-8.38 (35)	
PBu ⁿ ₃	(e)	0.74	0.49	THF 10	15	73 (96)	1845	5.13 (2.4)	-8.51 (35)	

a) All complexes except for PR₃ = PEt₃ gave satisfactory analytical results. b) Crude yields in parentheses.


c) KBr disc. d) 270 MHz in CD₃OD, J(P-H) in Hz are in parentheses. e) Not crystallizable.

f) Cy = *cyclo*-C₆H₁₂.



librium as shown in Eq. 3. The cationic complex analogous to **4** with a halide counter anion has been prepared by Dias *et al.* starting from Cp₂MoHX (X = Cl, Br, and I)⁶⁾ for PR₃ = PPh₃, PMe₂Ph, PEt₂Ph,⁷⁾ and the X-ray structure of [Cp₂MoH(PPh₃)]⁺I⁻ was determined to be a distorted tetrahedron.²⁾


Treatment of the cationic complex **4** with an equivalent NaOH in EtOH afforded phosphine-adduct of molybdenocene **2** (Eq. 5).⁸⁾ When PR₃ = PCy₃, **2d** was not formed but some uncharacterizable products were yielded together with **1**. In the case of PR₃ = PEt₃, the reaction was carried out *in situ* since isolation of corresponding **4b** in a pure state was difficult.



The phosphine derivatives of molybdenocene of the type **2** have been prepared for PR₃ = PPh₃ and PEt₃ independently by Azevedo *et al.*²⁾ and Geoffroy *et al.*³⁾ However, the present route finds an advantage over the previous routes in that it works under milder reaction conditions and thence it is convenient for the preparation of a series of complexes with a variety of tertiary phosphines and phosphite ligands. Complex **2c** reverted to the cationic hydrido complex **4c** on treatment with TsOH (Scheme 1).

Since it was difficult to obtain satisfactory analytical results for complex **2** due to its highly sensitive nature to the air, characterization by means of X-ray crystal structure analysis was attempted. A reddish brown prismatic crystal of **2e** (PR₃ = PBuⁿ₃) suitable for X-ray analysis was obtained by recrystallization from hexane. A full-matrix least-squares refinement procedure was used with anisotropic thermal parameters for the non-hydrogen atoms of the complex. The hydrogen atoms were placed at distance of C-H = 1.00 Å and included in least-square calculations without refinement of their parameters. ORTEP drawing of the molecule is shown in Fig. 1.⁹⁾

The molecule consists of wedge-formed two Cp rings coordinated to molybdenum in a mutually eclipsed conformation and a tributylphosphine ligand. Mo-P distance of 2.494(3) Å is a little shorter than that reported for [Cp₂MoH(PPh₃)]⁺I⁻ [2.501(4) Å].²⁾ Mean bond distances between molybdenum and Cp carbons (2.287


Fig. 1. Structure of $\text{Cp}_2\text{Mo}(\text{PBu}^n_3)$ (**2e**).

and 2.273 Å) are not unusual for this type of complexes.²⁾

Complex **2** reacted with alkyl halides ($\text{R}'\text{X}$) to give corresponding cationic alkyl complexes $[\text{Cp}_2\text{MoR}'(\text{PR}_3)]^+\text{X}^-$ (**5**) (Scheme 1). Recently Azevedo *et al.* have isolated **5** ($\text{R}' = \text{Me}$, $\text{R} = \text{Ph}$, $\text{X} = \text{PF}_6^-$) by an *in situ* reaction between **2a** and MeI and determined its X-ray structure.²⁾ They mentioned that alkyl halide other than MeI did not work similarly and failed to obtain analogous alkyl complexes. When complex **2c** with $\text{PR}_3 = \text{P}(\text{OEt})_3$ was allowed to react with EtI, PhCH_2Br , and $\text{CH}_2=\text{CH}-\text{CH}_2\text{I}$ as well as MeI, the orange colored corresponding alkyl derivatives (yields 72, 20, and 63%, respectively) as well as methyl complex (yield 63%) were obtained (Scheme 1). The product in the reaction of allyl iodide was found to possess η^1 -allyl ligand on the basis of spectral evidence.¹⁰⁾ The reaction between **2** ($\text{R} = \text{OEt}$) and Pr^iI afforded violet $[\text{Cp}_2\text{MoI}\{\text{P}(\text{OEt})_3\}]^+\text{I}^-$ (yield 74%) instead of an *i*-propyl derivative, probably via a rather unstable intermediary $[\text{Cp}_2\text{MoPr}^i(\text{PR}_3)]^+\text{I}^-$. The analogous cationic bromo complex, $[\text{Cp}_2\text{MoBr}\{\text{P}(\text{OEt})_3\}]^+\text{Br}^-$, was obtained as a dark colored solid by the reaction of $\text{Cp}_2\text{Mo}\{\text{P}(\text{OEt})_3\}$ (**2c**) with bromine in THF at room temperature (Scheme 1).

As reported for the PPh_3 derivative of **2**,²⁾ the other type **2** complexes with $\text{PR}_3 = \text{P}(\text{OEt})_3$ and PBu^n_3 reacted similarly with diethyl disulfide at 50 °C to give $\text{Cp}_2\text{Mo}(\text{SEt})_2$ possibly via an oxidative addition involving S-S bond cleavage. Furthermore, $\text{Cp}_2\text{Mo}(\text{PBu}^n_3)$ (**2e**) was found to activate dihydrogen under fairly mild conditions. Thus, the reaction of **2e** with 5 atm of H_2 in benzene at 50 °C for 4 h yielded dihydride **1** in 38% yield (Scheme). The rest of the complexes **2** other than **2e** failed to react with H_2 . Probably, feasibility of a dissociation of the phosphine ligand in **2e** due to its bulkiness may play an important role in this reaction.

Finally, complex **2c** was found to catalyze selective trimerization of phenyl isocyanate to give triphenyl isocyanurate (Scheme 1). In a typical experiment (not optimized), phenyl isocyanate (1.9 cm^3) and **2c** (0.43 mmol) in hexane (15 cm^3) were stirred *in vacuo* at room temperature for 18.5 h. During the period, a colorless precipitate accumulated gradually in the flask, which was filtered and recrystallized from acetone. 44% of phenyl isocyanate was converted into isocyanurate (589% for **2c**), which was identified by IR and mass spectrometry.



Scheme 1.

Although isocyanate is known to be trimerized by Lewis base such as tertiary amine,¹¹⁾ a control experiment using $P(OEt)_3$ as catalyst in place of **2c** did not give the cyclic trimer. Cyclo-addition reaction to give heterometallacycle in the reaction of $Cp_2Mo=O$ with $PhNCO$ and the coupling of $PhNCO$ with coordinated carbonyl ligand in $Cp_2W(CO)$ to give metallacycloimides have been reported.¹²⁻¹⁴⁾

We are grateful to Prof. K. Osakada of Tokyo Institute of Technology and Dr. S. Kurishima for X-ray structure analysis and to Dr. M. Tanaka of Tokyo Institute of Technology for elemental analysis. This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, and Culture of Japan (No. 02650621). T. I. thanks to Iwatani Scholarship and Shinsei-Shigen Kyokai for financial support.

References

- 1) T. Ito, M. Koga, S. Kurishima, M. Natori, N. Sekizuka, and K. Yoshioka, *J. Chem. Soc., Chem. Commun.*, **1990**, 988.
- 2) C. G. Azevedo, M. J. Calhorda, M. A. A. F. de C. T. Carrondo, A. R. Dias, V. Felix, and C. C. Romão, *J. Organomet. Chem.*, **391**, 345 (1990).
- 3) G. L. Geoffroy, M. G. Bradley, and R. Pierantozzi, "Transition Metal Hydrides," ed by R. Bau, American Chemical Society, Washington D. C. (1978), Chap. 14; G. L. Geoffroy and M. G. Bradley, *Inorg. Chem.*, **17**, 2410 (1978).
- 4) T. Igarashi and T. Ito, *Chem. Lett.*, **1985**, 1699.
- 5) **3**: IR (KBr) 1875 cm^{-1} [$\nu(\text{Mo-H})$]; ^1H NMR (CD_3OD) δ 5.3 (s, Cp), δ -9.4 ppm (s, Mo-H); E.A. Found: C, 51.77; H, 4.70; S, 7.91%. Calcd for $\text{C}_{17}\text{H}_{18}\text{MoO}_3\text{S}$: C, 51.26; H, 4.55; S, 8.05%.
- 6) A. R. Dias and C. C. Romão, *J. Organomet. Chem.*, **233**, 223 (1982).
- 7) C. G. de Azevedo, A. R. Dias, A. M. Martins, and C. C. Romão, *J. Organomet. Chem.*, **368**, 57 (1989).
- 8) $\text{Cp}_2\text{Mo}\{P(OEt)_3\}$ (**2c**), yield 76%; ^1H NMR (C_6D_6) δ = 4.19 [5H, d, Cp, $J(\text{H-P}) = 4.88\text{ Hz}$]; $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6) δ = 206.9 ppm down field from external PPh_3 . $\text{Cp}_2\text{Mo}(\text{P}^n\text{Bu}_3)$ (**2e**), yield 60%; ^1H NMR (C_6D_6) δ = 3.99 [5H, d, Cp, $J(\text{H-P}) = 4.29\text{ Hz}$]; $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6) δ = 42.0 ppm down field from external PPh_3 .
- 9) Crystal data: $\text{C}_{22}\text{H}_{37}\text{MoP}$, $M = 428.454$, orthorhombic, space group $Pbca$, $a = 18.193(5)$, $b = 25.414(3)$, $c = 9.541(2)\text{ \AA}$, $V = 4411.3\text{ \AA}^3$, $Z = 8$, $D_c = 1.29\text{ g cm}^{-3}$, $\lambda(\text{Mo K}\alpha) = 0.71069\text{ \AA}$, $R = 0.080$, $R_w = 0.073$ for 1697 reflections [$F_o > 3\sigma(F_o)$].
- 10) $[\text{Cp}_2\text{Mo}(\eta^1\text{-C}_3\text{H}_5)\{P(OEt)_3\}]^+\text{I}^-$, ^1H NMR (CD_3OD) δ = 5.21 [5H, d, Cp, $J(\text{H-P}) = 2.0\text{ Hz}$], 1.85 [2H, t, Mo- CH_2 -, $J(\text{H-P}) = 7.44\text{ Hz}$]; ^{13}C NMR (CD_3OD) δ = 94.2 (s, Cp), 8.3 [d, Mo- CH_2 -, $J(\text{C-P}) = 14.3\text{ Hz}$]. Although the η^1 -allyl derivative was not able to be isolated analytically pure due to contamination of $[\text{Cp}_2\text{MoI}\{P(OEt)_3\}]^+\text{I}^-$ in the product, satisfactory analytical results were obtained for the rest of the alkyl complexes reported here.
- 11) A. W. Hoffmann, *Chem. Ber.*, **3**, 761 (1860); **4**, 246 (1861).
- 12) P. Braunstein and D. Nobel, *Chem. Rev.*, **89**, 1927 (1989).
- 13) P. Jernakoff, G. L. Geoffroy, A. L. Rheingold, and S. J. Geib, *J. Chem. Soc., Chem. Commun.*, **1987**, 1610.
- 14) P. Jernakoff and N. J. Cooper, *J. Am. Chem. Soc.*, **109**, 2173 (1987).

(Received July 23, 1991)