

### Biological Activity

The estrogenic and antiestrogenic activities of these compounds were assessed on the basis of stimulation of the growth of the uterus of immature female rats. The test compounds were given by gavage either alone or in combination with estradiol (0.002 mg/kg per day) administered subcutaneously. On the 4th day the

uteri were excised, blotted dry, and weighed. The antifertility activity was determined by administering the compound by gavage to mature female rats for 6 days beginning the morning after a proven insemination. The rats were autopsied 9 days after the last medication and their uteri were removed and examined for implantation sites and gross abnormalities.

## Synthesis and Antiarrhythmic Activity of Naphthylalkylamines


GIANFRANCO PALA, ARTURO DONETTI, CARLA TURBA, AND SILVANO CASADIO

Research Laboratories of Istituto De Angeli, 20135 Milan, Italy

Received March 10, 1970

A further series of naphthylalkylamines was prepared and assayed for antiarrhythmic activity. Many of the compounds were found to be active *in vitro*, but only for five of them was activity confirmed *in vivo*. Comparative regression line analysis revealed that among the naphthylalkylamines so far investigated for antiarrhythmic activity, 1,5-dimorpholino-3-( $\alpha$ -naphthyl)pentane is still the most interesting one.

Our finding<sup>1</sup> that some  $\alpha$ -naphthylalkylamines, especially 1,5-dimorpholino-3-( $\alpha$ -naphthyl)pentane, possess marked antiarrhythmic activity led us to extend this investigation to 83 chemically related compounds. The new naphthylalkylamines had the general structures I-IV, in which R was an alkyl or aminoalkyl group; R<sub>1</sub> was a primary amino or aminomethyl group; NAA was a tertiary amino group;  $n = 2-4$ .



Naphthylalkylamines with R<sub>1</sub> = NH<sub>2</sub> were prepared from the corresponding amides by the Hofmann reaction. Reduction of the related nitriles with excess LAH in Et<sub>2</sub>O afforded naphthylalkylamines with R<sub>1</sub> = CH<sub>2</sub>NH<sub>2</sub>, reaction time and excess LAH depending on the steric hindrance of the nitriles.

**Pharmacology.**—All of the substances listed in Table II were submitted to the *in vitro* antiarrhythmic test, using quinidine and 1,5-dimorpholino-3-( $\alpha$ -naphthyl)pentane as reference standards. Many of them considerably reduced the maximal rate of stimulation of electrically driven isolated guinea pig auricles but did not inhibit the amplitude of contractions. These results are included in Table II in terms of relative potency, which was calculated from ED<sub>50</sub> values as

previously described<sup>2</sup> and expressed in relation to the antiarrhythmic activity of quinidine, which has been assigned the potency of 1.0.

Due to the promising results *in vitro*, all of the above compounds were tested subcutaneously in rats for the action on arrhythmias induced by CaCl<sub>2</sub>. The procedure was essentially the same as previously described,<sup>2</sup> except that 120 mg/kg of CaCl<sub>2</sub> was infused. Reference standards and expression of results were as *in vitro*. Of all the tested substances, only **51**, **64**, **82**,





Figure 1.—CaCl<sub>2</sub>-induced arrhythmias in rats. Regression lines of: **51** ( $\Delta$ — $\Delta$ ); **64** ( $\circ$ — $\circ$ ); **82** ( $\square$ — $\square$ ); **84** ( $\diamond$ — $\diamond$ ); **120** (x—x); 1,5-dimorpholino-3-( $\alpha$ -naphthyl)pentane (+—+); and quinidine ( $\nabla$ — $\nabla$ ).

(1) S. Casadio, G. Pala, T. Bruzzese, C. Turba, and E. Marazzi-Uberti, *J. Med. Chem.*, **13**, 418 (1970).

(2) C. Bianchi, G. P. Sanna, and C. Turba, *Arzneim. Forsch.*, **18**, 845 (1968).

TABLE I  
INTERMEDIATE NITRILES

Compd	R	R <sub>1</sub>		Structure	Yield, % <sup>a</sup>	Bp (mm) or mp, °C	Formula <sup>b</sup>
1	<i>n</i> -C <sub>3</sub> H <sub>7</sub>	CN	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	53.9	145–150 (0.2)	C <sub>19</sub> H <sub>24</sub> N <sub>2</sub>
2	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	CN	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	62.8	177–180 (0.2)	C <sub>20</sub> H <sub>26</sub> N <sub>2</sub>
3	<i>i</i> -C <sub>4</sub> H <sub>9</sub>	CN	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	82.3	147–150 (0.4)	C <sub>20</sub> H <sub>26</sub> N <sub>2</sub>
4	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	CH <sub>3</sub> (C <sub>2</sub> H <sub>5</sub> )N(CH <sub>2</sub> ) <sub>2</sub>	II	78	167–169 (0.2)	C <sub>21</sub> H <sub>28</sub> N <sub>2</sub>
5	CH <sub>3</sub> (C <sub>2</sub> H <sub>5</sub> )N(CH <sub>2</sub> ) <sub>2</sub>	CN	CH <sub>3</sub> (C <sub>2</sub> H <sub>5</sub> )N(CH <sub>2</sub> ) <sub>2</sub>	II	77.6	190–195 (0.6)	C <sub>22</sub> H <sub>30</sub> N <sub>2</sub>
6	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	74	170–175 (0.5)	C <sub>22</sub> H <sub>30</sub> N <sub>2</sub>
7	(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	CN	(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	84.6	174–178 (0.1)	C <sub>24</sub> H <sub>32</sub> N <sub>2</sub>
8	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CN	( <i>i</i> -C <sub>3</sub> H <sub>7</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	68.6	167–170 (0.3)	C <sub>23</sub> H <sub>32</sub> N <sub>2</sub>
9	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	( <i>i</i> -C <sub>3</sub> H <sub>7</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	51.6	168–170 (0.1)	C <sub>24</sub> H <sub>34</sub> N <sub>2</sub>
10	( <i>i</i> -C <sub>3</sub> H <sub>7</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	CN	( <i>i</i> -C <sub>3</sub> H <sub>7</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	70	200–205 (0.6)	C <sub>28</sub> H <sub>40</sub> N <sub>2</sub>
11	CH <sub>3</sub>	CN	<i>c</i>	II	35.3	180–182 (0.5)	C <sub>19</sub> H <sub>22</sub> N <sub>2</sub>
12	C <sub>2</sub> H <sub>5</sub>	CN	<i>c</i>	II	77.5	185–188 (0.7)	C <sub>20</sub> H <sub>24</sub> N <sub>2</sub>
13	<i>n</i> -C <sub>3</sub> H <sub>7</sub>	CN	<i>c</i>	II	74.8	190–191 (0.75)	C <sub>21</sub> H <sub>26</sub> N <sub>2</sub>
14	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CN	<i>c</i>	II	77.1	85–87	C <sub>21</sub> H <sub>26</sub> N <sub>2</sub>
15	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>c</i>	II	69.5	195–200 (0.15)	C <sub>22</sub> H <sub>28</sub> N <sub>2</sub>
16	<i>i</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>c</i>	II	69.3	180–183 (0.2)	C <sub>22</sub> H <sub>28</sub> N <sub>2</sub>
17	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>c</i>	II	89	187–190 (0.5)	C <sub>22</sub> H <sub>28</sub> N <sub>2</sub>
18	<i>c</i>	CN	<i>c</i>	II	78.8	215–220 (0.15)	C <sub>24</sub> H <sub>31</sub> N <sub>2</sub>
19	<i>n</i> -C <sub>3</sub> H <sub>7</sub>	CN	<i>d</i>	II	66.2	179–182 (0.05)	C <sub>22</sub> H <sub>28</sub> N <sub>2</sub>
20	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>d</i>	II	79.2	190–195 (0.2)	C <sub>23</sub> H <sub>30</sub> N <sub>2</sub>
21	<i>i</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>d</i>	II	63.5	205–209 (0.8)	C <sub>23</sub> H <sub>30</sub> N <sub>2</sub>
22	<i>n</i> -C <sub>3</sub> H <sub>7</sub>	CN	<i>e</i>	II	43.7	109–111	C <sub>21</sub> H <sub>26</sub> N <sub>2</sub> O
23	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>e</i>	II	39	200–205 (0.2)	C <sub>22</sub> H <sub>28</sub> N <sub>2</sub> O
24	<i>i</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>e</i>	II	52.4	190–195 (0.2)	C <sub>22</sub> H <sub>28</sub> N <sub>2</sub> O
25	CH <sub>3</sub>	CN	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	48.7	155–157 (0.2)	C <sub>18</sub> H <sub>22</sub> N <sub>2</sub>
26	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	79	164–167 (0.4)	C <sub>21</sub> H <sub>28</sub> N <sub>2</sub>
27	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>f</i>	II	54.3	96–99	C <sub>23</sub> H <sub>30</sub> N <sub>2</sub>
28	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>g</i>	II	44.4	123–124	C <sub>24</sub> H <sub>32</sub> N <sub>2</sub>
29	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>h</i>	II	58	90–92	C <sub>27</sub> H <sub>39</sub> N <sub>2</sub> O
30	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>4</sub>	II	29	165–170 (0.25)	C <sub>22</sub> H <sub>30</sub> N <sub>2</sub>
31	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CN	<i>i</i>	II	80	182–185 (0.25)	C <sub>23</sub> H <sub>30</sub> N <sub>2</sub>
32	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>i</i>	II	75.7	187–190 (0.1)	C <sub>24</sub> H <sub>32</sub> N <sub>2</sub>
33	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CN	<i>j</i>	II	71.7	207–210 (0.5)	C <sub>24</sub> H <sub>32</sub> N <sub>2</sub>
34	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>j</i>	II	61.7	196–200 (0.2)	C <sub>25</sub> H <sub>34</sub> N <sub>2</sub>
35	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CN	<i>k</i>	II	55	200–202 (0.25)	C <sub>23</sub> H <sub>30</sub> N <sub>2</sub> O
36	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>k</i>	II	70	205–210 (0.1)	C <sub>24</sub> H <sub>32</sub> N <sub>2</sub> O
37	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	III	75.3	150–155 (0.09)	C <sub>10</sub> H <sub>26</sub> N <sub>2</sub>
38	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CN	<i>e</i>	III	66	192–195 (0.15)	C <sub>21</sub> H <sub>26</sub> N <sub>2</sub> O
39	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>e</i>	III	74.5	193–196 (0.1)	C <sub>22</sub> H <sub>28</sub> N <sub>2</sub> O
40	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	(CH <sub>2</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	IV	41.5	62–63	C <sub>11</sub> H <sub>28</sub> N <sub>2</sub>
41	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CN	<i>d</i>	IV	41	96–98	C <sub>23</sub> H <sub>30</sub> N <sub>2</sub>
42	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CN	<i>d</i>	IV	66	102–103	C <sub>24</sub> H <sub>32</sub> N <sub>2</sub>

<sup>a</sup> Purified product. <sup>b</sup> All compounds were analyzed for C, H, N and the analytical results were within  $\pm 0.4\%$  of the theoretical values. <sup>c</sup> 2-(1-Pyrrolidinyl)ethyl. <sup>d</sup> 2-Piperidinoethyl. <sup>e</sup> 2-Morpholinoethyl. <sup>f</sup> 3-(1-Pyrrolidinyl)propyl. <sup>g</sup> 3-Piperidinopropyl. <sup>h</sup> 3-Morpholinopropyl. <sup>i</sup> 4-(1-Pyrrolidinyl)butyl. <sup>j</sup> 4-Piperidinobutyl. <sup>k</sup> 4-Morpholinobutyl.

84, and 120 had confirmed activity *in vivo*; the relative potencies (quinidine = 1.0) were 1.6, 0.7, 1.5, 3.4, and 0.8, respectively; the potency of 1,5-dimorpholino-3-( $\alpha$ -naphthyl)pentane was 1.2. However, an examination of the regression lines (Figure 1) revealed that the new compounds had a range of active doses narrower than both the reference standards.

On the basis of previous<sup>1</sup> and present results the conclusion may be drawn that, among the naphthylalkylamines so far investigated for antiarrhythmic activity, 1,5-dimorpholino-3-( $\alpha$ -naphthyl)pentane is still to be considered as the most interesting one.

### Experimental Section<sup>3</sup>

**Intermediates.**—Many of the nitriles were prepared as pre-

viously described.<sup>4–8</sup> The new nitriles (Table I) were obtained similarly. Except for the following compound, all the amides were prepared as previously reported.<sup>8–10</sup>

**$\alpha$ -(2-Dimethylaminoethyl)-1-naphthylacetamide.**—A solution of  $\alpha$ -(2-dimethylaminoethyl)-1-naphthylacetonitrile (10 g, 0.042 mol) and KOH (7 g, 0.125 mol) in 95% EtOH (40 ml) was refluxed for 3 hr with stirring, cooled, and poured into H<sub>2</sub>O. The separated pasty product was extracted (Et<sub>2</sub>O), washed (H<sub>2</sub>O), and dried (Na<sub>2</sub>SO<sub>4</sub>). Evaporation of the solvent yielded a

(4) S. Casadio, G. Pala, and T. Bruzzese, *Farmaco Ed. Sci.*, **17**, 871 (1962).

(5) S. Casadio, G. Pala, E. Crescenzi, T. Bruzzese, E. Marazzi-Uberti, and G. Coppi, *J. Med. Chem.*, **8**, 589 (1965).

(6) G. Pala, S. Casadio, T. Bruzzese, E. Crescenzi, and E. Marazzi-Uberti, *ibid.*, **8**, 698 (1965).

(7) G. Pala, S. Casadio, T. Bruzzese, and G. Coppi, *ibid.*, **9**, 786 (1966).

(8) S. Casadio, T. Bruzzese, G. Pala, G. Coppi, and C. Turba, *ibid.*, **9**, 707 (1966).

(9) S. Casadio, G. Pala, T. Bruzzese, E. Crescenzi, E. Marazzi-Uberti, and G. Coppi, *ibid.*, **8**, 594 (1965).

(10) M. Julia and M. Baillargé, *Bull. Soc. Chim. Fr.*, 928 (1957).

(3) Boiling points are uncorrected. Melting points are corrected and were taken on a Büchi capillary melting point apparatus.

TABLE II  
PHYSICAL PROPERTIES AND ANTIARRHYTHMIC ACTIVITY OF NAPHTHYLALKYLAMINES



Compd	R	R <sub>1</sub>		Structure	Method	Reaction conditions			Yield, % <sup>a</sup>	Bp (mm), °C <sup>b</sup>	Formula <sup>c</sup>	Rel potency
						Time, hr	mol ratio	LAH: nitrile				
43	CH <sub>3</sub>	NH <sub>2</sub>		I	A				66	118-120 (0.1)	C <sub>12</sub> H <sub>13</sub> N	0.6
44	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	NH <sub>2</sub>		I	A				62	130-133 (0.2)	C <sub>17</sub> H <sub>20</sub> N <sub>2</sub>	0.3
45	CH <sub>3</sub>	CH <sub>2</sub> NH <sub>2</sub>		I	B	20	2		81	112-113 (0.1)	C <sub>13</sub> H <sub>15</sub> N	<i>c</i>
46	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>		I	B	20	2		83	130-132 (0.2)	C <sub>15</sub> H <sub>19</sub> N	0.9
47	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>		I	B	10	4		82	130-132 (0.2)	C <sub>16</sub> H <sub>21</sub> N	<i>c</i>
48	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	CH <sub>2</sub> NH <sub>2</sub>		I	B	10	2		63	147-150 (0.3)	C <sub>16</sub> H <sub>22</sub> N <sub>2</sub>	0.4
49	CH <sub>3</sub> (C <sub>2</sub> H <sub>5</sub> )N(CH <sub>2</sub> ) <sub>2</sub>	CH <sub>2</sub> NH <sub>2</sub>		I	B	10	2		67	141-142 (0.3)	C <sub>17</sub> H <sub>22</sub> N <sub>2</sub>	<i>c</i>
50	(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	CH <sub>2</sub> NH <sub>2</sub>		I	B	10	2		60	150-153 (0.2)	C <sub>18</sub> H <sub>26</sub> N <sub>2</sub>	<i>c</i>
51	<i>d</i>	CH <sub>2</sub> NH <sub>2</sub>		I	B	20	2		76	183-184 (0.2)	C <sub>17</sub> H <sub>20</sub> N <sub>2</sub>	1.2
52	<i>c</i>	CH <sub>2</sub> NH <sub>2</sub>		I	B	40	5		53	180 (0.1)	C <sub>18</sub> H <sub>24</sub> N <sub>2</sub> O	<i>c</i>
53	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>3</sub>	CH <sub>2</sub> NH <sub>2</sub>		I	B	10	2		57	170-172 (0.2)	C <sub>17</sub> H <sub>21</sub> N	0.3
54	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	A				83	137-139 (0.2)	C <sub>15</sub> H <sub>20</sub> N <sub>2</sub>	<i>c</i>
55	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	NH <sub>2</sub>	<i>d</i>	II	A				26	180-183 (0.5)	C <sub>21</sub> H <sub>30</sub> N <sub>2</sub>	<i>c</i>
56	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>3</sub>	II	A				45	140-141 (0.1)	C <sub>16</sub> H <sub>23</sub> N <sub>2</sub>	<i>c</i>
57	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	NH <sub>2</sub>	<i>f</i>	II	A				23	190-194 (0.4)	C <sub>22</sub> H <sub>32</sub> N <sub>2</sub>	<i>c</i>
58	CH <sub>3</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	B	15	4		78	145 (0.2)	C <sub>17</sub> H <sub>24</sub> N <sub>2</sub>	0.4
59	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	B	15	2		62	152-155 (0.2)	C <sub>18</sub> H <sub>26</sub> N <sub>2</sub>	0.5
60	<i>n</i> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	B	10	2		67	157-158 (0.1)	C <sub>17</sub> H <sub>25</sub> N <sub>2</sub>	0
61	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	B	20	4		77	148-149 (0.1)	C <sub>18</sub> H <sub>27</sub> N <sub>2</sub>	0.7
62	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	B	4	2		73	167-169 (0.1)	C <sub>21</sub> H <sub>30</sub> N <sub>2</sub>	<i>c</i>
63	<i>i</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	B	20	2		85	157-158 (0.5)	C <sub>20</sub> H <sub>30</sub> N <sub>2</sub>	<i>c</i>
64	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	B	15	5		80	168-170 (0.4)	C <sub>20</sub> H <sub>30</sub> N <sub>2</sub>	0.4
65	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	B	60	4		68	160-162 (0.1)	C <sub>20</sub> H <sub>31</sub> N <sub>3</sub>	<i>c</i>
66	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	CH <sub>3</sub> (C <sub>2</sub> H <sub>5</sub> )N(CH <sub>2</sub> ) <sub>2</sub>	II	B	20	4		82	174-176 (0.8)	C <sub>20</sub> H <sub>30</sub> N <sub>2</sub>	<i>c</i>
67	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	CH <sub>3</sub> (C <sub>2</sub> H <sub>5</sub> )N(CH <sub>2</sub> ) <sub>2</sub>	II	B	20	4		81	176-178 (0.7)	C <sub>21</sub> H <sub>32</sub> N <sub>2</sub>	<i>c</i>
68	CH <sub>3</sub> (C <sub>2</sub> H <sub>5</sub> )N(CH <sub>2</sub> ) <sub>2</sub>	CH <sub>2</sub> NH <sub>2</sub>	CH <sub>3</sub> (C <sub>2</sub> H <sub>5</sub> )N(CH <sub>2</sub> ) <sub>2</sub>	II	B	10	4		58	163-164 (0.1)	C <sub>22</sub> H <sub>35</sub> N <sub>3</sub>	<i>c</i>
69	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	B	10	4		67	163-165 (0.5)	C <sub>21</sub> H <sub>32</sub> N <sub>2</sub>	<i>c</i>
70	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	B	15	6		82	168-170 (0.2)	C <sub>22</sub> H <sub>35</sub> N <sub>2</sub>	<i>c</i>
71	(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	CH <sub>2</sub> NH <sub>2</sub>	(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	B	10	3		82	180-182 (0.2)	C <sub>24</sub> H <sub>37</sub> N <sub>2</sub>	<i>c</i>
72	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	CH <sub>3</sub> (C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> )N(CH <sub>2</sub> ) <sub>2</sub>	II	B	15	6		72	184-185 (0.1)	C <sub>21</sub> H <sub>30</sub> N <sub>2</sub>	<i>c</i>
73	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	( <i>i</i> -C <sub>3</sub> H <sub>7</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	B	40	6		68	180-182 (0.1)	C <sub>20</sub> H <sub>30</sub> N <sub>2</sub>	<i>c</i>
74	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	( <i>i</i> -C <sub>3</sub> H <sub>7</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	B	30	9		60	176-178 (0.4)	C <sub>24</sub> H <sub>37</sub> N <sub>2</sub>	<i>c</i>
75	( <i>i</i> -C <sub>3</sub> H <sub>7</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	CH <sub>2</sub> NH <sub>2</sub>	( <i>i</i> -C <sub>3</sub> H <sub>7</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	II	B	10	3		80	184-185 (0.5)	C <sub>25</sub> H <sub>41</sub> N <sub>3</sub>	<i>c</i>
76	CH <sub>3</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>g</i>	II	B	10	2		76	168-171 (0.1)	C <sub>19</sub> H <sub>26</sub> N <sub>2</sub>	<i>c</i>
77	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>g</i>	II	B	10	2		67	179-181 (0.2)	C <sub>20</sub> H <sub>28</sub> N <sub>2</sub>	0.7
78	<i>n</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>g</i>	II	B	4	2		71	190-192 (0.2)	C <sub>21</sub> H <sub>30</sub> N <sub>2</sub>	<i>c</i>
79	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>g</i>	II	B	20	5		81	188-190 (0.3)	C <sub>21</sub> H <sub>30</sub> N <sub>2</sub>	0.3
80	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>g</i>	II	B	4	3		85	188-190 (0.4)	C <sub>22</sub> H <sub>32</sub> N <sub>2</sub>	<i>c</i>
81	<i>i</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>g</i>	II	B	10	3		78	182-184 (0.4)	C <sub>22</sub> H <sub>32</sub> N <sub>2</sub>	1.5
82	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>g</i>	II	B	40	4		75	194-196 (0.3)	C <sub>22</sub> H <sub>32</sub> N <sub>2</sub>	0.7
83	<i>g</i>	CH <sub>2</sub> NH <sub>2</sub>	<i>g</i>	II	B	10	3		77	216-218 (0.3)	C <sub>23</sub> H <sub>36</sub> N <sub>3</sub>	<i>c</i>
84	CH <sub>3</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>d</i>	II	B	15	3		75	170-173 (0.1)	C <sub>20</sub> H <sub>25</sub> N <sub>2</sub>	0.4
85	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>d</i>	II	B	10	2		81	172-174 (0.2)	C <sub>21</sub> H <sub>30</sub> N <sub>2</sub>	1.8
86	<i>n</i> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>d</i>	II	B	20	2		59	170-173 (0.1)	C <sub>22</sub> H <sub>32</sub> N <sub>2</sub>	0.6
87	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>d</i>	II	B	40	3		79	172-175 (0.1)	C <sub>22</sub> H <sub>32</sub> N <sub>2</sub>	1.3
88	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>d</i>	II	B	4	2		89	192-194 (0.5)	C <sub>23</sub> H <sub>34</sub> N <sub>2</sub>	<i>c</i>
89	<i>i</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>d</i>	II	B	10	4		86	190-192 (0.4)	C <sub>23</sub> H <sub>34</sub> N <sub>2</sub>	1.3
90	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>d</i>	II	B	30	2		68	196-199 (0.3)	C <sub>23</sub> H <sub>34</sub> N <sub>2</sub>	1.3
91	<i>d</i>	CH <sub>2</sub> NH <sub>2</sub>	<i>d</i>	II	B	15	2		79	220-223 (0.1)	C <sub>24</sub> H <sub>37</sub> N <sub>2</sub>	<i>c</i>
92	CH <sub>3</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>e</i>	II	B	15	3		76	175-176 (0.1)	C <sub>19</sub> H <sub>25</sub> N <sub>2</sub> O	<i>c</i>
93	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>e</i>	II	B	15	3		74	195-196 (0.2)	C <sub>20</sub> H <sub>28</sub> N <sub>2</sub> O	0.3
94	<i>n</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>e</i>	II	B	20	2		58	190-191 (0.2)	C <sub>21</sub> H <sub>30</sub> N <sub>2</sub> O	<i>c</i>
95	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>e</i>	II	B	105	4		71	193-195 (0.2)	C <sub>21</sub> H <sub>30</sub> N <sub>2</sub> O	0.6
96	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>e</i>	II	B	10	4		68	200-201 (0.4)	C <sub>22</sub> H <sub>32</sub> N <sub>2</sub> O	<i>c</i>
97	<i>i</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>e</i>	II	B	15	3		72	198-200 (0.5)	C <sub>22</sub> H <sub>32</sub> N <sub>2</sub> O	2.0
98	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>e</i>	II	B	20	6		77	203-204 (0.3)	C <sub>22</sub> H <sub>32</sub> N <sub>2</sub> O	0.9
99	<i>c</i>	CH <sub>2</sub> NH <sub>2</sub>	<i>e</i>	II	B	180	4		43	230-233 (0.1)	C <sub>24</sub> H <sub>36</sub> N <sub>3</sub> O <sub>2</sub>	<i>c</i>
100	CH <sub>3</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>3</sub>	II	B	10	4		81	150-151 (0.1)	C <sub>18</sub> H <sub>26</sub> N <sub>2</sub>	<i>c</i>
101	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>3</sub>	II	B	20	2		66	166-168 (0.2)	C <sub>20</sub> H <sub>30</sub> N <sub>2</sub>	0.4
102	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>3</sub>	II	B	15	4		78	167-169 (0.2)	C <sub>21</sub> H <sub>32</sub> N <sub>2</sub>	0.4
103	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>3</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>3</sub>	II	B	10	4		60	186-188 (0.3)	C <sub>22</sub> H <sub>30</sub> N <sub>2</sub>	<i>c</i>
104	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>h</i>	II	B	30	5		83	180-182 (0.2)	C <sub>22</sub> H <sub>32</sub> N <sub>2</sub>	<i>c</i>
105	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>h</i>	II	B	60	4		84	195-196 (0.4)	C <sub>23</sub> H <sub>34</sub> N <sub>2</sub>	<i>c</i>

TABLE II (Continued)

Compd	R	R <sub>1</sub>		Reaction —conditions—				Bp (mm), °C	Formula <sup>b</sup>	Rel po- tency	
				Struc- ture	Method	Time, hr	Yield, mol ratio %				
106	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>f</i>	II	B	40	3	77	186–188 (0.2)	C <sub>23</sub> H <sub>34</sub> N <sub>2</sub>	<i>c</i>
107	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>f</i>	II	B	40	9	73	193–195 (0.2)	C <sub>24</sub> H <sub>36</sub> N <sub>2</sub>	<i>c</i>
108	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>i</i>	II	B	20	4	71	198–200 (0.1)	C <sub>22</sub> H <sub>32</sub> N <sub>2</sub> O	<i>c</i>
109	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>i</i>	II	B	30	5	87	190–192 (0.2)	C <sub>25</sub> H <sub>34</sub> N <sub>2</sub> O	<i>c</i>
110	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>4</sub>	II	B	10	2	71	156–158 (0.1)	C <sub>21</sub> H <sub>32</sub> N <sub>2</sub>	1.0
111	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>4</sub>	II	B	15	4	86	170–172 (0.1)	C <sub>22</sub> H <sub>34</sub> N <sub>2</sub>	0.7
112	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>j</i>	II	B	40	3	71	195–196 (0.1)	C <sub>23</sub> H <sub>34</sub> N <sub>2</sub>	<i>c</i>
113	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>j</i>	II	B	40	3	75	184–186 (0.1)	C <sub>24</sub> H <sub>36</sub> N <sub>2</sub>	<i>c</i>
114	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>k</i>	II	B	40	3	80	183–191 (0.2)	C <sub>24</sub> H <sub>36</sub> N <sub>2</sub>	<i>c</i>
115	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>k</i>	II	B	40	3	75	195–197 (0.2)	C <sub>25</sub> H <sub>38</sub> N <sub>2</sub>	<i>c</i>
116	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>l</i>	II	B	40	5	78	220–223 (0.4)	C <sub>23</sub> H <sub>34</sub> N <sub>2</sub> O	<i>c</i>
117	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>l</i>	II	B	15	3	92	210–212 (0.3)	C <sub>24</sub> H <sub>36</sub> N <sub>2</sub> O	<i>c</i>
118	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	III	B	15	4	56	154–156 (0.1)	C <sub>19</sub> H <sub>28</sub> N <sub>2</sub>	1.8
119	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	III	B	4	5	70	162–164 (0.2)	C <sub>20</sub> H <sub>30</sub> N <sub>2</sub>	2.3
120	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>e</i>	III	B	4	5	64	205–206 (0.2)	C <sub>21</sub> H <sub>30</sub> N <sub>2</sub> O	1.6
121	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>e</i>	III	B	4	5	77	207–208 (0.1)	C <sub>22</sub> H <sub>32</sub> N <sub>2</sub> O	2.0
122	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	IV	B	3	2	69	154–156 (0.1)	C <sub>23</sub> H <sub>30</sub> N <sub>2</sub>	0.7
123	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	IV	B	15	2	83	162–164 (0.1)	C <sub>21</sub> H <sub>32</sub> N <sub>2</sub>	<i>c</i>
124	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>d</i>	IV	B	15	2	87	184–183 (0.1)	C <sub>23</sub> H <sub>34</sub> N <sub>2</sub>	0.9
125	<i>sec</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>2</sub> NH <sub>2</sub>	<i>d</i>	IV	B	15	4	76	202–204 (0.5)	C <sub>24</sub> H <sub>36</sub> N <sub>2</sub>	<i>c</i>
1,5-Dimorpholino-3-( $\alpha$ -naphthyl)pentane											1.8
Quinidine											1.0

<sup>a</sup> Distilled product. <sup>b</sup> All compounds were analyzed for C, H, N and the analytical results were within  $\pm 0.4\%$  of the theoretical values. <sup>c</sup> Inactive or cardiotoxic compound. <sup>d</sup> 2-Piperidinoethyl. <sup>e</sup> 2-Morpholinoethyl. <sup>f</sup> 3-Piperidinopropyl. <sup>g</sup> 2-(1-Pyrrolidinyl)ethyl. <sup>h</sup> 3-(1-Pyrrolidinyl)propyl. <sup>i</sup> 3-Morpholinopropyl. <sup>j</sup> 4-(1-Pyrrolidinyl)butyl. <sup>k</sup> 4-Piperidinobutyl. <sup>l</sup> 4-Morpholinobutyl.

residue which, on trituration with 1:1 Et<sub>2</sub>O–petroleum ether (bp 40–70°) gave a colorless solid (6.9 g, 64%), mp 79–81°. Anal. (C<sub>16</sub>H<sub>20</sub>N<sub>2</sub>O) C, H, N.

**Naphthylalkylamines** with R<sub>1</sub> = NH<sub>2</sub> or CH<sub>2</sub>NH<sub>2</sub> are listed in Table II, and their preparation is illustrated by the following methods.

**Method A. 1-Dimethylamino-3-amino-3-( $\alpha$ -naphthyl)-4-methylpentane (54).**— $\alpha$ -Isopropyl- $\alpha$ -(2-dimethylaminoethyl)-1-naphthylacetamide (17.5 g, 0.059 mol) was added with stirring to a solution of Na (2.7 g, 1.17 g-atom) in anhydrous MeOH (100 ml), and then Br<sub>2</sub> (9.38 g, 0.059 mol) was rapidly dropped into the solution. After 6 hr stirring at room temperature, the mixture was allowed to stand overnight, and the solvent was removed under reduced pressure. The residue was dissolved in Et<sub>2</sub>O, washed (H<sub>2</sub>O), and dried (Na<sub>2</sub>SO<sub>4</sub>) and the solution was evaporated to dryness. The new residue was dissolved in 95% EtOH (130 ml), 50% KOH (130 ml) was added to it, and the mixture was refluxed for 6 hr, poured into cold H<sub>2</sub>O, and extracted (Et<sub>2</sub>O). The extract was washed (H<sub>2</sub>O) and dried (Na<sub>2</sub>SO<sub>4</sub>),

the solvent was evaporated, and the residue was distilled to give a viscous and colorless oil, bp 137–139° (0.2 mm).

**Method B. N-[3-Aminomethyl-3-( $\alpha$ -naphthyl)heptyl]piperidine (88).**—A solution of  $\alpha$ -*n*-butyl- $\alpha$ -(2-piperidinoethyl)-1-naphthylacetone (50 g, 0.15 mol) in dry Et<sub>2</sub>O (100 ml) was dropped at room temperature for 2 hr into a stirred suspension of LAH (11.35 g, 0.3 mol) in dry Et<sub>2</sub>O (900 ml). The mixture was refluxed for 4 hr with stirring, cooled, and cautiously decomposed with H<sub>2</sub>O (100 ml). The organic layer was separated, washed (H<sub>2</sub>O), and dried (Na<sub>2</sub>SO<sub>4</sub>). The solvent was evaporated and the residue was distilled to give a viscous and colorless oil, bp 192–194° (0.5 mm).

**Acknowledgments.**—The authors wish to thank Dr. R. Perego for performing the microanalyses, Mr. G. Bietti and Mr. E. Bellora for assistance in preparing the compounds, and Mr. G. Bertuzzi and Mr. P. Duranti for carrying out the antiarrhythmic tests.