

A Journal of the Gesellschaft Deutscher Chemiker

# Angewandte Chemie

GDCh

International Edition

[www.angewandte.org](http://www.angewandte.org)

## Accepted Article

**Title:** Chemoproteomic Profiling of Phosphoaspartate Modifications in Prokaryotes

**Authors:** Jae Won Chang, Gihoon Lee, Jeffrey E Montgomery, and Raymond Moellering

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

**To be cited as:** *Angew. Chem. Int. Ed.* 10.1002/anie.201809059  
*Angew. Chem.* 10.1002/ange.201809059

**Link to VoR:** <http://dx.doi.org/10.1002/anie.201809059>  
<http://dx.doi.org/10.1002/ange.201809059>

## COMMUNICATION


## Chemoproteomic Profiling of Phosphoaspartate Modifications in Prokaryotes

Jae Won Chang, Jeffrey E. Montgomery, Gihoon Lee, and Raymond E. Moellering\*

**Abstract:** Phosphorylation at aspartic acid residues represents an abundant and critical posttranslational modification (PTM) in prokaryotes. In contrast to most characterized PTMs, such as phosphorylation at serine or threonine, the phosphoaspartate moiety is intrinsically labile, and therefore incompatible with common proteomic profiling methods. Here we report a nucleophilic, desthiobiotin-containing hydroxylamine (DBHA) chemical probe that covalently labels modified aspartic acid residues in native proteomes. DBHA treatment coupled with LC-MS/MS analysis enabled detection of known phosphoaspartate modifications, as well as novel aspartic acid sites in the *E. coli* proteome. Coupled with isotopic labelling, DBHA-dependent proteomic profiling also permitted global quantification of changes in endogenous protein modification status, as demonstrated here with the detection of increased *E. coli* OmpR phosphorylation, but not abundance, in response to changes in osmolarity. The DBHA probe and proteomic workflow reported herein should be broadly useful for the discovery and quantitative interrogation of phosphoaspartate modifications in diverse organisms and biological contexts.

Posttranslational modifications (PTMs) have evolved as a selective and reversible means to alter protein function, and thus propagate signals within and between cells. These chemical tags are exceptionally diverse in structure and function, and are known to target a majority of proteins in many organisms, including mammals [1–2]. While the prevalence and perceived importance of PTMs is conserved across all forms of life, the relative abundance of specific PTMs is extremely variable. Humans, for example, encode more than 500 protein kinases that catalytically phosphorylate the alcohol functionality in serine, threonine and tyrosine residues [3–4]. Protein phosphorylation is recognized as a major communication mechanism in mammalian cells, a fact that is bolstered by high association with disease when phosphorylation networks are deregulated [5]. An important technical consideration when reflecting upon the vast number of phosphorylation marks that are known in mammalian systems is the fact that systematic, quantitative analysis of these PTMs is made possible through several technologies, including detection

with phospho-specific antibodies, and global identification with LC-MS/MS proteomic approaches [6–7]. Combined, these methods have identified thousands of phosphorylation sites in diverse cell types and organisms. The comprehensive set of tools available to study protein phosphorylation, which is invoked here as a prototypical PTM, hinges on a single aspect of the modification itself: chemical stability to biochemical and proteomic workflows.



**Figure 1.** DBHA probe labelling, detection and quantification of modified aspartic acids in native proteins. (a) Schematic of dynamic phosphorylation at aspartic acids, as well as labelling with a prototypical nucleophilic probe *in situ*, permitting detection and quantification. (b) Chemical structure of the desthiobiotin-linked hydroxylamine (DBHA) probe. (c) Western blot visualization of desthiobiotin-labelled proteins demonstrates dose-dependent and hydroxylamine competitive labelling of *E. coli* proteins in DBHA-treated lysates.

[a] Dr. J.W. Chang, J.E. Montgomery, G. Lee, Prof. Dr. R. E. Moellering  
Department of Chemistry  
Institute for Genomics and System Biology  
The University of Chicago  
5735 S. Ellis Ave, Chicago, IL 60637 (USA)  
E-mail: rmoellering@uchicago.edu

Supporting information for this article is given via a link at the end of the document.

Prokaryotes also employ phosphorylation as a major form of biochemical communication, but in contrast to eukaryotes the number of serine/threonine phosphorylation sites appears to be orders of magnitude lower [8]. Instead, these lower organisms rely more on two-component signalling pathways that involve phosphorylation of histidine “receiver” residues, which ultimately phosphorylate aspartic acids to form an acylphosphate modification [9–11]. The resulting phosphoaspartate residue (pD) is

## COMMUNICATION

generally thought to cause changes in protein structure, which in the case of many characterized response regulator (RR) transcription factors can promote or inhibit DNA binding and subsequent downstream signalling. While thought to represent the major form of signal transduction, the widespread discovery and dynamic quantification of these modifications has not been possible due to the intrinsic lability of the pD group itself. As a result, pD sites have almost exclusively been studied one-by-one with the use of  $^{32}\text{P}$ -ATP as a means to radiolabel the active protein for visualization by gel electrophoresis<sup>[12]</sup>. Even in these settings, however, protein purification is often required and the endogenous environment cannot be probed. Therefore, new methods with which to study these chemically labile yet functionally important modifications are necessary. Here we report a chemoproteomic approach to rapidly trap the electrophilic acylphosphate group unique to pD sites within native bacterial proteomes (Figure 1a). We demonstrate that a desthiobiotin-containing hydroxylamine (DBHA) probe selectively enriches and site-specifically identifies both known and novel electrophilic aspartic acid sites throughout the *E. coli* proteome. Furthermore, we demonstrate the potential to measure signalling dynamics in the osmolarity sensing OmpR/OmpF two-component signalling pathway by combining isotopic proteome labelling and pD-site enrichment from native proteomes.

We previously reported that intrinsically labile metabolites, such as the glycolytic metabolite 1,3-bisphosphoglycerate, could be efficiently trapped and converted into a stable hydroxamic acid derivative upon treatment with hydroxylamine *in situ*<sup>[13]</sup>. Based on the success of this approach for acylphosphate-containing metabolites, and precedence for mapping ADP-ribosylation<sup>[14]</sup>, we reasoned that a hydroxylamine-based chemical probe may be useful for capturing pD sites in the proteome. Such a probe would encompass two main components: 1) an O-alkyl hydroxylamine warhead capable of forming a stable intermediate with pD sites in native proteins under mild conditions; 2) a retrieval tag that permits direct enrichment and elution of modified proteins or peptides for LC-MS/MS or other quantitative detection methods (Figure 1a; Figure S1). We first attempted to use a minimal probe, O-propargyl hydroxylamine, which would label modified aspartates with a small alkyne tag that could be used for detection or enrichment following [3+2] Huisgen “click” chemistry with secondary retrieval/visualization tags<sup>[15]</sup>. This approach is similar to a recent report that employed an alkyne-containing hydrazine probe in mammalian cells<sup>[16]</sup>. Our preliminary experiments using O-propargyl hydroxylamine suffered from poor retrieval of tagged species as measured by either in-gel fluorescence or LC-MS/MS, which we attribute to decreased efficiency that comes with additional steps involved in click chemistry sample processing. Therefore, we sought to develop a probe that could be directly retrieved and processed for a variety of proteomic detection methods. We synthesized a desthiobiotin-containing O-alkyl hydroxylamine (DBHA) probe that could bypass the need for subsequent chemical modification prior to detection or enrichment (Figure 1b). Desthiobiotin was specifically used in place of biotin to permit efficient release of enriched peptides for LC-MS/MS detection, which can be a major bottleneck in chemical proteomic workflows<sup>[17]</sup>.

We first tested whether incubation of DBHA across assay conditions would covalently label native proteins in K12 *E. coli*.


We found that DBHA labelling was optimal under the denaturing conditions of 6 M urea, which likely improves probe access to pD modification sites. Proteome incubation with DBHA and Western blot detection of desthiobiotin labelled proteins revealed dose-dependent labelling of the *E. coli* proteome across a wide range of concentrations (Figure 1c). These labelling events were inhibited by pretreatment of proteome with soluble hydroxylamine or acidic conditions (pH = 2), confirming that the proteins and sites being visualized were dependent upon intact reactivity (Figure 1c; Figure S2). Determining the site of labelling is particularly important for probes such as DBHA because they could, in principle, react with numerous modifications on proteins. For example, hydrazine-probes have previously been shown to modify glycosylated proteins in their open-chain form<sup>[18]</sup>, as well as capture pyruvoyl and glyoxylyl modifications in mammalian cells<sup>[16]</sup>. Despite the potential for interaction with other electrophilic moieties that are present on proteins, our probe and workflow should differentiate these events through the detection of the DBHA-modified aspartic acids directly on tryptic peptides with high resolution LC-MS/MS and site-of-labelling analysis (*vide infra*; Figure S1). To identify the proteins and specific sites being modified by DBHA we developed an efficient workflow for modified protein enrichment and processing by LC-MS/MS analysis (Figure S3). DBHA treatments were performed at 10 mM and 100 mM concentrations in order to maximally capture both low and high abundance sites within the *E. coli* proteome. Modified proteins were directly enriched with streptavidin-agarose immunoprecipitation, washing to remove non-labelled proteins, and processed for on-bead tryptic digestion. Importantly, bulk tryptic peptides from enriched proteins could be collected in the digest eluent, followed by elution of DBHA-modified peptides. The combination of these two pools enables high-confidence detection of target proteins (i.e. from many detected tryptic peptides per protein) as well as site-specific resolution of DBHA-modification sites. DBHA-labelled sites were filtered based on stringent mass tolerance, decoy database false-discovery rates, and MS<sup>2</sup> statistics. Beyond these initial filters, we required that a putative modification site be observed in four or more biological replicates to be included in this dataset. Finally, in order to identify the specific DBHA-modified residue in each detected tryptic peptide we analyzed the “hits” that passed our initial specificity filters above using the LuciPHor modification site localization algorithm<sup>[19]</sup>. Application of this analysis pipeline to LC-MS/MS data from the 10 mM ( $n$  = eight replicates) and 100 mM ( $n$  = six replicates) DBHA datasets permitted detection of 138 high confidence pAsp sites from 98 proteins (site global false-localization rate in LuciPHor at < 1%; Table S1).

The identification of >100 modified aspartic acids in the *E. coli* proteome was surprising since there are ~20-30 predicted RR's present in this and other typical prokaryotes<sup>[20]</sup>. Analysis of the local sequence context of DBHA-modified aspartic acids revealed a modest motif preference for small, hydrophilic residues around the DBHA-modified aspartic acid, as well as preference for proline at the +4 position (Figure 2a). This overall profile was distinct from the motif surrounding known response regulator pD sites in *E. coli*. However, a conserved preference for proline at the +4 position was also present, confirming partial similarities between the two motifs (Figure 2a). Bioinformatic analysis of modified proteins with DAVID and KEGG pathway analysis revealed a dominant signature of proteins involved in metabolism, protein folding and

## COMMUNICATION

transcriptional regulation (Figure 2b; Figure S4). These sites were found in proteins spanning the entire range of relative abundance in *E. coli*, from metabolic proteins like Pkg (estimated at ~100,000 copies/cell) at one extreme, to transcription factors like BasR (estimated at ~20 copies/cell) on the other (Table S1; Figure S5)<sup>[20]</sup>. Among the lower abundance proteins identified were response regulators and related two-component signalling proteins with known pD sites. For example, the response regulators OmpR and BasR were identified harboring DBHA-modifications at D55 and D51, respectively (Figure 2c-d). These sites are known phosphorylation sites that regulate OmpR and

BasR DNA-binding activity<sup>[21-22]</sup>. Additional DBHA-modification sites included known phosphohistidine-containing proteins, like Asp38 in Crr. Intriguingly, the DBHA modification site in Crr was found to be highly proximal to a known phosphohistidine site, perhaps indicating that dynamic relay between pH and pD sites occurs within this and other proteins (Figure S6)<sup>[23]</sup>. Taken together, this dataset validated DBHA-labelling and enrichment as a method to detect both known and potentially novel phosphoaspartate sites in native proteins from whole prokaryotic proteome.



**Figure 2.** DBHA-dependent proteomic profiling identifies known and novel aspartic acid modification sites in *E. coli*. (a) Enriched sequence motifs derived from DBHA-modified aspartic acid sites (top) compared to the motif generated by known response regulator pD sites in *E. coli* (bottom). (b) Gene-ontology biological process categories (GOTERM\_BP; bottom) and KEGG pathways (top) enriched among DBHA-modified proteins. Representative, statistically significant categories are shown, with a complete list in **Suppl. Fig. 4**. (c-d) MS/MS spectra of DBHA-modified tryptic peptides at two known pD sites in the response regulator proteins OmpR (c) and BasR (d). Observed b- and y-ions are labelled on each spectrum and highlighted on the tryptic peptide. The known pD sites at D55 and D51 in OmpR and BasR, respectively, are highlighted and starred (\*) in red.

DBHA-enrichment of proteins via labelling of pD sites presents an opportunity to directly interrogate changes in phosphorylated proteins in parallel to, and distinct from, monitoring global changes in protein abundance. To determine whether DBHA-profiling could identify physiologically-relevant changes in two-component signalling, we interrogated the well-characterized EnvZ/OmpR pathway in *E. coli*. This pathway facilitates sensing of extracellular osmolarity by the histidine kinase and sensor protein EnvZ, which phosphorylates the response regulator OmpR at D55 in response to increased


osmolarity. Phosphorylated OmpR exhibits markedly increased DNA-binding activity and differential regulation of transcription of the membrane porin proteins OmpF and OmpC (Figure S7)<sup>[24-25]</sup>. To track this activation directly in cells, *E. coli* growing in standard LB media were switched to either 20% sucrose-containing NB (high osmolarity) or NB alone (mock), and grown for 12 hr. A minor fraction of each bulk proteome was trypsinized, isotopically labelled with unique tandem-mass tag (TMT) channels<sup>[26]</sup>, and pooled to compare protein abundance changes resulting from high osmolarity (Figure 3a). In parallel, the bulk of



## COMMUNICATION

the native proteome from each condition was labelled with DBHA probe, enriched on streptavidin-agarose resin and typrisinized on-bead. The resulting tryptic peptides were labelled with unique TMT channels and pooled for LC-MS/MS quantitation of phosphorylation changes in response to high osmolarity (Figure 3a). Abundance changes for more than 1,200 proteins were quantified between conditions (Figure 3b; Table S2). OmpR protein levels were increased 1.4-fold in response to high osmolarity, which matches literature precedent showing that OmpR levels remain relatively constant or slightly increase in response to increased osmolarity (Figure 3b)<sup>[27]</sup>. This abundance profile also detected significantly decreased levels of OmpF protein, consistent with repression by active OmpR. Significant changes in the levels of other proteins involved in osmoregulation were also apparent in the global profile, including the

transcriptional regulator BetI (downregulated in high osmolarity) and the target of its repression, BetB (upregulated in high osmolarity; Fig. 3b)<sup>[28-29]</sup>. Quantification of DBHA-enriched proteins, in contrast, identified OmpR as the most affected protein, with nearly 4-fold higher levels of DBHA-enriched OmpR present in the cells grown in 20% sucrose (Figure 3c; Table S2). Other known response regulators that harbor pD sites, including CpxR and ArcA, were detected and quantified in the enriched profile, but were not affected by osmolarity at either the protein abundance (Figure 3b) or pD level (Figure 3c). Combined, these data validate that DBHA enrichment coupled to quantitative proteomics enables global profiling of response regulator phosphorylation, as opposed to abundance, in response to physiological signals in *E. coli*.



**Figure 3.** DBHA-dependent LC-MS/MS enables global profiling of pD modification dynamics. (a) Proteomic workflow to quantify global changes in protein levels (top) and pD or other DBHA-reactive modifications (bottom) by LC-MS/MS. Lysates from high and low osmolarity samples of *E. coli* are split for protein level quantification (top) or treated with DBHA probe (bottom). This workflow thus generates two isotopically labelled tryptic peptide pools that provide quantitative information on bulk protein level changes (top) and DBHA-sensitive modification levels (bottom). (b) Global quantification of protein level changes in response to high osmolarity in *E. coli*. Proteins are ranked from most down-regulated (left) to up-regulated (right) under high osmolarity conditions. Proteins discussed in the text are labelled in red with their quantitative ratio listed. (c) Quantification of DBHA-enriched proteins in response to high osmolarity in *E. coli*, thus reporting on specific changes in modification status, in contrast to changes in abundance alone. The osmoregulatory response regulatory OmpR was the most affected protein in the dataset. All data points in (b) and (c) represent the mean ratios from  $n = 4$  (two technical of two biological) replicate runs.

In summary, we have developed a novel chemical proteomic workflow that offers several advantages over current approaches to interrogate phosphoaspartate sites, and likely other electrophilic modifications, in native proteomes. First, peptide-based enrichment and LC-MS/MS profiling provides unequivocal identification of modified residues. This not only enables quantitation of pD signalling dynamics at specific sites, but also allows for exploratory mapping in unannotated proteomes and organisms. In line with this attribute, the dataset herein contained many high-confidence, reproducible DBHA-modified residues that are not previously annotated as phosphoaspartate sites. These sites may represent heretofore unrecognized phosphoaspartate

sites that are introduced through enzymatic or non-enzymatic means, as has been observed for other intrinsically reactive metabolic intermediates like acetylphosphate<sup>[30-32]</sup>, which is used *in vitro* to chemically phosphorylate Asp sites in response regulators<sup>[33]</sup>. Beyond phosphoaspartate, other modifications could contribute to the profile observed here, including methylesterification, ADP-ribosylation, and potentially other less-characterized PTMs that occur at aspartic acids in prokaryotes. Despite this potential for overlap, we view the labelling of additional electrophilic modifications as a feature not a flaw of DBHA- profiling because the use quantitative LC-MS/MS provides site-specific resolution necessary for follow up analysis at a site-

## COMMUNICATION

by-site level. The dataset produced herein represents a starting point for these future efforts. Finally, we have demonstrated the ability to detect physiologically-relevant changes in phosphoaspartic acid levels, distinct from protein level differences, in the EnvZ/OmpR osmoregulatory pathway. This work is the first to enable global, quantitative measurement of phosphoaspartate modification dynamics in native proteome, which should be broadly applicable to other organisms and biologic perturbations, providing an unprecedented view of this elusive, yet critical, signalling mechanism.

## Acknowledgements

We thank M. Rust and G. Pattanayak for discussions surrounding the manuscript, and R. Park for conversations about proteomic analyses. We are grateful for financial support of this work from the following: Kwanjeong Educational Fellowship (to G.L.); NIH 2T32GM008720-16 (to J.E.M.); NIH R00CA175399, DP2GM128199-01, and 2R01CA093577-11 (to R.E.M.); the Damon Runyon Cancer Research Foundation (DFS08-14 to R.E.M.), and The University of Chicago.

## Conflict of interest

The authors declare no conflict of interest.

**Keywords:** Chemoproteomics • Two-component signalling • Response regulator • Phosphoaspartate modification

- [1] C. Walsh, Posttranslational modification of proteins: expanding nature's inventory (Roberts and Co. Publishers, Englewood, Colo), **2006**, pp. xxi, 490 p.
- [2] J.V. Olsen, M. Mann, *Molecular & cellular proteomics* **2013**, 12, 3444-3452.
- [3] P. Cohen, *Trends Biochem Sci* **2000**, 25, 596-601.
- [4] G. Manning, D.B. Whyte, R. Martinez, T. Hunter, S. Sudarsanam, *Science* **2002**, 298, 1912-1934.
- [5] J. Brognard, T. Hunter, *Curr Opin Genet Dev* **2011**, 21, 4-11.
- [6] M. Mann, O.N. Jensen, *Nature biotechnology* **2003**, 21, 255-261.
- [7] J.M. Kee, R.C. Oslund, D.H. Perlman, T.W. Muir, *Nature Chemical Biology* **2013**, 9, 416-421.
- [8] A.M. Stock, V.L. Robinson, P.N. Goudreau, *Annual review of biochemistry* **2000**, 69, 183-215.
- [9] J.B. Stock, A.J. Ninfa, A.M. Stock, *Microbiol Rev* **1989**, 53, 450-490.
- [10] C.M. Potel, M.H. Lin, A.J.R. Heck, S. Lemeer, *Nature Methods* **2018**, 15, 187-190.
- [11] P.V. Attwood, P.G. Besant, M.J. Piggott, *Amino acids* **2011**, 40, 1035-1051.
- [12] M.M. Igo, A.J. Ninfa, T.J. Silhavy, *Genes & development* **1989**, 3, 598-605.
- [13] J.W. Chang, G. Lee, J.S. Coukos, R.E. Moellering, *Analytical chemistry* **2016**, 88, 6658-6661.
- [14] Y. Zhang, J. Wang, M. Ding, Y. Yu, *Nature methods* **2013**, 10, 981-984.
- [15] A.E. Speers, B.F. Cravatt, *Chemistry & biology* **2004**, 11, 535-546.
- [16] M.L. Matthews, L. He, B.D. Horning, E.J. Olson, B.E. Correia, J.R.3<sup>rd</sup>. Yates, P.E. Dawson, B.F. Cravatt, *Nat Chem* **2017**, 9, 234-243.
- [17] J.D. Hirsch, L. Eslamizar, B.J. Filanoski, N. Malekzadeh, R.P. Haugland, J.M. Beechem, R.P. Haugland, *Analytical biochemistry* **2002**, 308, 343-357.
- [18] H. Zhang, X.J. Li, D.B. Martin, R. Aebersold, *Nature biotechnology* **2003**, 21, 660-666.
- [19] D. Fermin, S.J. Walmsley, A.C. Gingras, H.W. Choi, A.I. Nesvizhskii, *Molecular & Cellular Proteomics* **2013**, 12, 3409-3419.
- [20] A. Schmidt, K. Kochanowski, S. Vedelaar, E. Ahn, B. Volkmer, L. Callipo, K. Knoops, M. Bauer, R. Aebersold, M. Heinemann, *Nature biotechnology* **2016**, 34, 104-110.
- [21] K. Yamamoto, K. Hirao, T. Oshima, H. Aiba, R. Utsumi, A. Ishihama, *The Journal of biological chemistry* **2005**, 280, 1448-1456.
- [22] J. Delgado, S. Forst, S. Harlocker, M. Inouye, *Molecular microbiology* **1993**, 10, 1037-1047.
- [23] K.A. Presper, C.Y. Wong, L. Liu, N.D. Meadow, S. Roseman, *Proceedings of the National Academy of Sciences of the United States of America* **1989**, 86, 4052-4055.
- [24] Y.L. Jo, F. Nara, S. Ichihara, T. Mizuno, S. Mizushima, *The Journal of biological chemistry* **1986**, 261, 15252-15256.
- [25] H. Aiba, F. Nakasai, S. Mizushima, T. Mizuno, *The Journal of biological chemistry* **1989**, 264, 14090-14094.
- [26] L. Dayon, A. Hainard, V. Licker, N. Turck, K. Kuhn, D.F. Hochstrasser, P.R. Burkhard, J.C. Sanchez, *Analytical chemistry* **2008**, 80, 2921-2931.
- [27] S.J. Cai, M. Inouye, *The Journal of biological chemistry* **2002**, 277, 24155-24161.
- [28] T.P. Rkenes, T. Lamark, A.R. Strom, *J Bacteriol* **1996**, 178, 1663-1670.
- [29] O.B. Styrvoid, P. Falkenberg, B. Landfald, M.W. Eshoo, T. Bjornsen, A.R. Strom, *J Bacteriol* **1986**, 165, 856-863.
- [30] R.E. Moellering, B.F. Cravatt, *Science* **2013**, 341, 549-553.
- [31] B.T. Weinert, V. Iesmantavicius, S.A. Wagner, C. Scholz, B. Gummesson, P. Bell, T. Nystrom, C. Choudhary, *Molecular cell* **2013**, 51, 265-272.
- [32] R. Harmel, D. Fiedler, *Nature chemical biology* **2018**, 14, 244-252.
- [33] L. Rajeev, E.G. Luning, P.S. Dehal, M.N. Price, A.P. Arkin, A. Mukhopadhyay, *Genome Biol* **2011**, 12, R99.

## COMMUNICATION


## Entry for the Table of Contents

Layout 1:

## COMMUNICATION

**Phosphoaspartate**

**modifications:** A desthiobiotin-containing hydroxylamine (DBHA) chemical probe covalently labels modified aspartic acid residues in native proteomes. Coupled with isotopic labelling, DBHA-dependent proteomic profiling enables global quantification of changes in phosphoaspartate modification levels and protein levels.



Dr. Jae Won Chang, Jeffrey E. Montgomery, Gihoon Lee, Prof. Dr. Raymond E. Moellering\*

Page No. – Page No.

**Chemoproteomic Profiling of Phosphoaspartate Modifications in Prokaryotes**