## A platinum-catalyzed annulation reaction leading to medium-sized rings<sup>†</sup>

Dirk Hildebrandt, Wiebke Hüggenberg, Matthias Kanthak, Tobias Plöger, Iris M. Müller and Gerald Dyker\*

Received (in Cambridge, UK) 20th February 2006, Accepted 4th April 2006 First published as an Advance Article on the web 25th April 2006 DOI: 10.1039/b602498j

A platinum-catalyzed domino process with intermediate benzopyrylium cations reaches its optimum utility in the formation of 7- and 8-membered rings. With iron(III) chloride, a tetracyclic product is isolated, derived from an oxidative transformation of a metal–carbene intermediate.

Whereas synthetic methods for the construction of five- and sixmembered rings are numerous, the development of novel and efficient cyclization reactions leading to medium-sized carbocycles and heterocycles is still challenging because of the disadvantageous influence of both entropic and enthalpic factors.<sup>1</sup> In the course of our studies on gold- and platinum-catalyzed annulation reactions with ortho-alkynyl-substituted benzaldehydes,<sup>2</sup> we intended to test the generality of the transformation depicted in Scheme 1. The formation of five- and six-membered naphthalena-carbocycles of type 2, catalyzed by gold salts, has recently been described.<sup>3</sup> In the case of our total synthesis of heliophenanthrone,<sup>2c</sup> which resembles a substituted derivative of **2** (with n = 0), we found PtCl<sub>2</sub> to be the superior catalyst compared to gold chloride.<sup>4</sup> Therefore, we have concentrated on PtCl<sub>2</sub> as the catalyst for the reactions in Scheme 1, testing its scope and limitations with respect to ring size and the presence of heteroatoms.

For the synthesis of model compounds **1a–f**, we started from 2-bromobenzaldehyde (**3**), as illustrated in Scheme 2. The bisalkyne side chain<sup>5</sup> was either introduced directly in a one-step procedure by a Sonogashira coupling reaction<sup>6</sup> (for **1c–f**), or *via* the corresponding cyclic acetal<sup>7</sup> and intermediate alcohols **4a** and **4b**. In the latter case, the ether functionalities were prepared with propargyl bromide, using NaH as the base,<sup>8</sup> and the aldehyde was finally deprotected with dilute hydrochloric acid to give **1a** and **1b**.<sup>9</sup> Surprisingly, another established method of deprotection—FeCl<sub>3</sub>



Scheme 1 PtCl<sub>2</sub>-catalysed annulations. a: PtCl<sub>2</sub> (3 mol%), toluene, reflux.

Fakultät für Chemie, Ruhr-Universität Bochum, Universitätsstrasse 150, D-44780 Bochum, Germany. E-mail: Gerald.Dyker@rub.de; Fax: +49 (0)234 32-14353; Tel: +49 (0)234 32-24551

<sup>†</sup> Electronic supplementary information (ESI) available: Summary crystal data for **6b**, and full synthesis and characterisation data for all compounds prepared. See DOI: 10.1039/b602498j



Scheme 2 Synthesis of the precursors 1a–f. a: ethylene glycol, TMSCl, DCM, reflux; b:  $Pt(PPh_3)_2Cl_2$  (2 mol%), CuI (2 mol%), TEA, reflux; c: NaH, propargyl bromide, THF, 20 °C; d: 5% HCl, THF, 20 °C.

in dichloromethane at room temperature<sup>10</sup>—led to the formation of annulation product **6b** in 10% isolated yield, identified by NMR and X-ray crystal structure analysis (Fig. 1).<sup>11</sup> In the control experiment, substrate **1b** reacted under the same conditions to give the [3 + 2] cycloadduct **6b** in 13% yield; a related product being recently observed by Gupta *et al.* in a similar gold-catalyzed process with a participating allenyl unit.<sup>12</sup>

We generally performed the platinum-catalyzed domino process on a 0.5 mmol scale: substrates **1a–f** were heated in 20 ml toluene and a catalytic amount of PtCl<sub>2</sub> (3 mol%) in a screw-capped flask at 140 °C. The results are summarized in Table 1. Substrates **1a–c** were consumed in 3 h, and the corresponding annulation products **2a–c** were obtained in isolated yields of 67, 89 and 95%, respectively (Table 1, entries 1–3). Formation of an eightmembered ring occurred smoothly to give product **2d** in a very satisfactory 80% yield (Table 1, entry 4), although a somewhat enhanced reaction time was necessary. Consequently, for the generation of a nine-membered carbocycle in product **2e**, the time for the consumption of **1e** rose to 24 h (Table 1, entry 5). In



Fig. 1 Crystal structure of 6b.

Entry	Substrate	Reaction time/h	Product	Ring size	Yield (%)
1	1a	3	2a	6	67 <sup>a</sup>
2	1b	3	2b	7	89
3	1c	3	2c	7	95
4	1d	12	2d	8	80
5	1e	24	2e	9	13
6	1f	24	2f	15	$< 5^{b}$
a o o	1 1	b G1: 1 (1 )	11		

<sup>t</sup> 0.2 mmol scale. <sup>b</sup> Slightly impure, not fully characterized.



Scheme 3 Plausible mechanism for annulation reaction.

addition, the yield of annulation product 2e dropped to only 13%, indicating a limitation of this process. Since the ring strain of carbocycles generally reaches its maximum in nine-membered rings, we hoped for higher yields in the cases of macrocycles. However, an attempt to produce a 15-membered ring from substrate **1f** resulted in a very low yield (Table 1, entry 6).

For comparison, we also tried AuCl<sub>3</sub> (5 mol%, 80 °C in acetonitrile) as a catalyst for the annulation reaction. With **1a** and **1e** as substrates, we achieved yields of less than 10%, confirming that  $PtCl_2$  is superior.

A plausible mechanism for the platinum(II)-catalyzed annulation reaction is depicted in Scheme 3. The double annulation process most probably proceeds through the benzopyrylium cation **I**, which results from the nucleophilic attack of the carbonyl oxygen at the alkyne, activated by the Lewis-acidic platinum salt. A subsequent intramolecular Huisgen-type [3 + 2] cycloaddition of the second alkyne is assumed to generate intermediate **II**.<sup>13</sup> Rearrangement to **III** and the formal [4 + 2] cycloaddition product **IV** leads to the aromatized final **2b**, liberating the active catalyst, in analogy to the calculations of Straub.<sup>13</sup> In the case of FeCl<sub>3</sub> as the Lewis acid, we assume that intermediate **II** is oxidatively transformed to **6b**.<sup>14</sup>

The platinum-catalyzed intramolecular domino annulation reaction of *ortho*-alkynyl benzaldehydes described herein represents a versatile approach to naphthalenes with annulated carbocycles or heterocycles of various ring sizes. The easily accessible starting materials, simple reaction procedure and mild conditions make this method attractive for the production of common- and medium-sized rings.

We gratefully acknowledge financial support from Fonds der Chemischen Industrie and the Deutsche Forschungsgemeinschaft (DFG project DY 12/7-2).

## Notes and references

- (a) G. Illuminati and L. Mandoline, Acc. Chem. Res., 1981, 14, 95–102;
  (b) G. A. Molander, Acc. Chem. Res., 1998, 31, 603–609.
- G. Dyker, D. Hildebrandt, J. Liu and K. Merz, Angew. Chem., 2003, 115, 4536–4538, Angew. Chem., Int. Ed., 2003, 42, 4399–4002; (b) D. Kadzimirsz, D. Hildebrandt, K. Merz and G. Dyker, Chem. Commun., 2006, 661–662; (c) G. Dyker and D. Hildebrandt, J. Org. Chem., 2005, 70, 6093–6096.
- 3 (a) N. Asao, K. Sato, Menggenbateer and Y. Yamamoto, J. Org. Chem., 2005, **70**, 3682–3685; (b) K. Sato, N. Asao and Y. Yamamoto, J. Org. Chem., 2005, **70**, 8977–8981; (c) N. Kim, Y. Kim, W. Park, D. Sung, A. K. Gupta and C. H. Oh, Org. Lett., 2005, **7**, 5289–5291.
- 4 For reviews on gold catalysis, see: (a) G. Dyker, Angew. Chem., 2000, 112, 4407-4409, Angew. Chem., Int. Ed., 2000, 39, 4237-4239; (b) A. S. K. Hashmi, Gold Bull., 2004, 37, 51-65; (c) A. Hoffmann-Röder and N. Krause, Org. Biomol. Chem., 2005, 3, 387-391. Current examples of Pt(II) catalysis: (d) H. Kusawa, Y. Miyashita, J. Takaya and N. Iwasawa, Org. Lett., 2006, 8, 289-292; (e) C. Y. Lo, M. P. Kumar, H.-K. Chang, S.-F. Lush and R.-S. Liu, J. Org. Chem., 2005, 70, 10482-10487; (f) E. Soriano and J. Marco-Contelles, J. Org. Chem., 2005, 70, 9345-9353; (g) A. Fürstner and P. W. Davies, Am. Chem. Soc., 2005, 127, 15024–15025; (h) I. Nakamura, J. Y Mizushima and Y. Yamamoto, J. Am. Chem. Soc., 2005, 127, 15022-15023; (i) C. Nevado and A. M. Echavarren, Chem.-Eur. J., 2005, 11, 3155-3164; (j) M. P. Muñoz, J. Adrio, J. C. Carretero and A. M. Echavarren, *Organometallics*, 2005, **24**, 1293–1300; (*k*) G. B. Bajracharya, I. Nakamura and Y. Yamamoto, *J. Org. Chem.*, 2005, 70, 892-897; (1) E. Soriano and J. Marco-Contelles, Chem.-Eur. J., 2005, 11, 521-533.
- 5 The bis-alkynes are commercially available, except for 1,15-hexadecadiyne, which was prepared according to the literature: B. Hellbach, R. Gleiter and F. Rominger, *Synthesis*, 2003, 2535–2541.
- 6 (a) K. Sonogashira, Y. Tohda and N. Hagihara, *Tetrahedron Lett.*, 1975, **50**, 4467–4470; (b) K. R. Roesch and R. C. Larock, *J. Org. Chem.*, 2002, **67**, 86–94.
- 7 T. H. Chan, M. A. Brook and T. Chaly, Synthesis, 1983, 203-205.
- 8 B. C. Bales, J. H. Horner, X. Huang, M. Newcomb, D. Crich and M. M. Greenberg, J. Am. Chem. Soc., 2001, **123**, 3623–3629.
- 9 P. A. Grieco, Y. Yokoyama, G. O. Withers, F. J. Okuniewicz and C.-L. J. Wang, J. Org. Chem., 1978, 43, 4178–4182.
- 10 S. Sen, S. Roach, J. Boggs, G. Ewing and J. Magrath, J. Org. Chem., 1997, 62, 6684–6886.
- 11 Crystal data for **6b**:  $C_{14}H_{12}O_3$ , M = 228.24, monoclinic, space group  $P_{2_1/n}$ , a = 10.592(4), b = 6.912(6), c = 15.076(8) Å,  $\beta = 94.25(3)^\circ$ , V = 1100.6(11) Å<sup>3</sup>, Z = 4, T = 100(2) K,  $\mu = 0.097$  mm<sup>-1</sup>, 8305/1991/1603 reflections collected/unique/observed,  $R_{int} = 0.0435$ ,  $R_1 = 0.0521$ ,  $wR_2$  (all data) = 0.1172, 154 parameters. All non-hydrogen atoms calculated anisotropic; positions of the H atoms calculated for idealised positions. The structure was solved and refined using SHELXTL-97: G. M. Sheldrick, SHELXTL-97. Program for the solution and refinement of crystal structures, Universität Göttingen, Germany, 1997. CCDC 299431. For crystallographic data in CIF or other electronic format see DOI: 10.1039/b602498j.
- 12 A. K. Gupta, C. Y. Rhim, C. H. Oh, R. S. Mane and S.-H. Han, Green Chem., 2006, 8, 25–28.
- 13 For a related Huisgen-type [3 + 2] cycloaddition, which was calculated by a DFT study, see: B. F. Straub, *Chem. Commun.*, 2004, 1726–1728.
- 14 A. G. M. Barrett, J. Mortier, M. Sabat and M. A. Sturgess, Organometallics, 1988, 7, 2553–2561.