

0040-4039(95)02331-3

The Role of Alkoxyl Radicals in Gif (GoAgg^v) Chemistry.¹

Darren W. Snelgrove,² Philip A. MacFaul,³ Keith U. Ingold^{*} and Danial D. M. Wayner^{*}

Steacie Institute for Molecular Sciences, National Research Council of Canada Ottawa, Ontario, Canada, K1A 0R6

Abstract: The oxidation of cyclooctane using *tert*-alkyl hydroperoxides under $GoAgg^V$ conditions has been examined using selected alkyl hydroperoxides as mechanistic probes. The results indicate that the products are formed by alkoxyl radical-induced reactions, there being no need to invoke any chemistry involving high valent iron-oxo species.

Enzyme mimetic chemistry has long been a very popular research area. The cytochrome P-450's⁴ and methane monooxygenases (MMO's)⁵ have attracted particular attention because they can functionalize alkanes. High valent iron-oxo species are generally believed to be the active intermediates that cleave the alkanes' unactivated C-H bonds. Barton et al⁶ have conjectured that the numerous "Gif" chemical systems actually mimic the behavior and mode of action of P-450 and / or MMO enzymes. Most of the Gif systems use an iron salt as a catalyst, hydrogen peroxide or *tert*-butyl hydroperoxide (TBHP) as the primary oxidant,⁷ and a pyridine / acetic acid mixture as the solvent. The substrate is commonly a cyclic alkane and the reactivity of the catalyst is frequently enhanced by the addition of picolinic acid.

In Gif systems which use an Fe^{III} salt as the catalyst together with TBHP, Barton has proposed⁸ that a high valent iron-oxo species (formally $Fe^V=O$) is formed which can insert into an unactivated C-H bond to give an alkyl-Fe^V complex (e.g. c-C₆H₁₁-Fe^V-OH). This complex (together with dioxygen, if available) is the supposed precursor of the observed organic reaction products (mainly ketones from cycloalkanes, alcohols from *tert*-CH bonds and hydroperoxides in the presence of O₂). Of course, all these products could also be accounted for in terms of "normal" free radical chemistry initiated by the formation of the *tert*-butoxyl radical, see reactions 1-10. Indeed, Minisci et al⁹ have shown that *tert*-butoxyl radicals are present under certain Gif conditions by their trapping with electron-rich alkenes.¹⁰ Herein, we report results obtained with cyclooctane in a specific Gif system, GoAgg^V.^{8b} We have employed the "usual" TBHP and three other *tert*-alkyl hydroperoxides each of which serves as a mechanistic probe. These are "probes" because if the hydroperoxide, ROOH, yields the corresponding alkoxyl radical, RO*, this radical will show a characteristic reaction product "signature".



Oxidation of cyclooctane (0.142M) with TBHP (0.284M) using $GoAgg^V$ conditions (0.0142M Fe^{III} (NO₃)₃ in pyridine, acetic acid (10:1, v/v) and 0.0426 M picolinic acid at room temperature) under air gave cyclooctanone, cyclooctanol, cyclooctyl *tert*-butyl peroxide and 2-cyclooctylpyridine.¹¹ The

results of duplicate experiments are given in Table 1, with product yields expressed in equivalents relative to the Fe^{III} catalyst. Duplicate experiments were also carried out under an argon atmosphere after prior deoxygenation of the reactants (see also Table 1). Of course, these conditions are not strictly "anaerobic" because dioxygen is formed continuously via reaction 4. Naturally, under these "anaerobic" conditions the yield of ketone is reduced (by ca. 40%). However, the yields of mixed peroxide (reaction 9) and 2-cyclooctylpyridine are increased. These aerobic results (apart from the mixed peroxide) are consistent with earlier work on cyclooctane and cyclohexane in the GoAgg^V system (see Table 1).^{8b} "Anaerobic" results have also been reported for the oxidation of cyclooctane by a GoAgg^{IV} system, i.e. without the addition of picolinic acid, which indicated that the mixed peroxide was formed but the yields of ketone and alcohol were not given.^{8b}

Table 1. The Reaction of Cycloalkanes with tert-Butyl Hydroperoxide under GoAgg^V Conditions.

Cycloalkane	Conditions	<i>c</i> -C8H14O	с-С8H15ОН	c-C8H15OOBut	2-c-C8H15-C5H4N
Cyclooctane	Aerobic	3.78	0.02	0.09	Trace
Cyclooctane	Aerobic	3.53	0.05	0.13	а
Cyclooctaneb	Aerobic	3.72	0.16	с	c
Cyclohexaned	Aerobic	5.10 ^e	0.56 ^f	c,g	с
Cyclooctane	Anaerobic	2.12	0.04	0.31	0.013
Cyclooctane	Anaerobic	2.17	0.07	0.42	0.013

^a Not detected. ^b Yields of cyclooctane-derived products (in mmol) from a competitive oxidation of cyclooctane and cyclohexane, duration of experiment unspecified (ref. 8b). ^c Not reported. ^d Yields (in mmol) for the oxidation of cyclohexane, duration of experiment unspecified (ref. 8b). ^e c-C₆H₁₀O. ^f c-C₆H₁₀OH. ^g c-C₆H₁₀OBu^t.

We have previously employed 2-methyl-1-phenyl-2-propyl hydroperoxide (MPPH) as a mechanistic probe to explore the real chemistry lying behind another purported example of alkane oxidation via a high valent iron-oxo species.¹² The alkoxyl radical derived from this hydroperoxide undergoes extremely rapid β -scission.

$$PhCH_2CMe_2O \bullet \qquad \longrightarrow \qquad PhCH_2 \bullet + Me_2CO \qquad (11)$$

Replacement of the "usual" TBHP in a $GoAgg^V$ system by MPPH gave mainly products derived from the benzyl radical formed in reaction 11, viz. benzaldehyde, benzyl alcohol, bibenzyl and the mixed peroxide formed by the cross-coupling of PhCH₂[•] and PhCH₂CMe₂OO[•] radicals (see Table 2). A very small amount of cyclooctanone was produced under aerobic conditions but not under "anaerobic" conditions. We suggest that under aerobic conditions further oxidation of the main product, benzaldehyde, yields the highly reactive benzoylperoxyl radical¹³ and that it is this radical which abstracts a hydrogen atom from the cyclooctane.

$$PhCHO \xrightarrow{\text{ROO}} PhC(O)OO \xrightarrow{\text{c-C}_8H_{16}} PhC(O)OOH + c-C_8H_{15} \xrightarrow{\text{O}_2} c-C_8H_{14}O \quad (12)$$

Table 2. The Reaction of Cyclooctane with MPPH under GoAgg^V Conditions.

Conditions	<i>c</i> -C ₈ H ₁₄ O	PhCHO	PhCH ₂ OH	(PhCH ₂) ₂	PhCH ₂ OO- CMe ₂ CH ₂ Ph
Aerobic	0.19	10.4	0.04	0.07	0.06
Aerobic	0.27	10.5	0.02	0.05	0.06
Anaerobic	а	1.7	0.46	0.56	2.96
Anaerobic	а	1.7	0.66	0.72	3.56

a Not detected.

The alkoxyl radical derived from *o*-methyl cumene hydroperoxide (*o*-MCH) is about 100 times longer lived than the alkoxyl radical derived from MPPH.¹⁴ Nevertheless, when *o*-MCH is employed in place of TBHP in the $GoAgg^V$ system only minor amounts of cyclooctanone were produced (see Table 3). The main

products are those derived from unimolecular decay reactions of the o-methylcumyloxyl radicals, viz. β -scission (reaction 13) and intramolecular hydrogen abstraction (reaction 14).

(13)
$$o-CH_3C_6H_4COMe + Me^* \stackrel{\beta}{\longleftarrow} o-CH_3C_6H_4CMe_2O^* \stackrel{\text{intra}}{\longrightarrow} o-CH_2C_6H_4CMe_2OH$$
 (14)

Table 3. The Reaction of Cyclooctane with o-Methyl Cumene Hydroperoxide under GoAgg^V Conditions.

Conditions	<i>c</i> -C ₈ H ₁₄ O	o-Me-	o-OCH-	o-HOCH2-	o-MeC6H4CMe2OO-	
		C ₆ H ₄ COMe	C ₆ H ₄ CMe ₂ OH	C ₆ H ₄ CMe ₂ OH	CH ₂ C ₆ H ₄ CMe ₂ OH	
Aerobic	0.09	0.02	1.98	0.86	0.75	
Aerobic	0.11	0.10	0.81	0.76	0.97	
Anaerobic	0.10	0.13	0.64	0.95	1.07	
Anaerobic	0.09	0.12	0.58	0.90	1.11	

With cumene hydroperoxide (CH) the GoAgg^V reaction yields cyclooctane derived products reminiscent of the TBHP reaction, however acetophenone (formed by the β -scission of "free" cumyloxyl radicals, reaction 15), is also a major product, particularly under "anaerobic" conditions (see Table 4).

$$C_6H_5CMe_2O^{\bullet} \longrightarrow C_6H_5COMe + Me^{\bullet}$$
 (15)

Table 4. The Reaction of Cyclooctane with Cumene Hydroperoxide under GoAgg^V Conditions.

Conditions	<i>c</i> -C ₈ H ₁₄ O	<i>с</i> -С ₈ Н ₁₅ ОН	PhC(O)Me	c-C ₈ H ₁₅ OOCMe ₂ Ph	2-c-C8H15-C5H4Na
Aerobic	3.36	0.20	5.24	1.00	b
Aerobic	3.42	0.19	6.78	1.01	b
Anaerobic	1.61	0.11	5.89	0.59	0.048
Anaerobic	1.40	0.12	7.27	0.79	0.016
CU-C-U-N	as not received from	Collent b Not data	ated		

^a CH₃C₅H₄N was not resolved from C₅H₅N. ^b Not detected.

The tertiary alkyl hydroperoxides we have employed were chosen because they yield *tert*-alkoxyl radicals having vastly different rate constants for unimolecular β -scission (plus intramolecular H-atom abstraction for the *o*-MCH-derived radical). The lifetimes¹⁵ of the alkoxyl radicals are: PhCH₂CMe₂O• ~ 5 ns;¹⁴ *o*-MeC₆H₄CMe₂O• ~ 400 ns in CH₃CN;¹⁴ C₆H₅CMe₂O• ~ 1.6 µs in CH₃CN,¹⁶ ~ 0.5 µs in CH₃COOH;¹⁶ and Me₃CO•¹⁷ ~ 12 µs in CH₃CN, ~ 3.6 µs in CH₃COOH. The pseudo-first-order rate constant for H-atom abstraction from cyclooctane by an alkoxyl radical is¹² k₅[C₈H₁₆] = 2.7 x 10⁶ M⁻¹ s⁻¹ x 0.142 M = 3.8 x 10⁵ s⁻¹, corresponding to a lifetime of 2.6 µs. Thus, with the two short-lived alkoxyls, PhCH₂CMe₂O• and *o*-MeC₆H₄CMe₂O•, the principal products are those derived from their unimolecular reactions, whereas the relatively long-lived Me₃CO• radical yields products derived mainly via H-atom abstraction from the cycloalkane. With CH under "anaerobic" conditions acetophenone constitutes ca. 74% of the products which is roughly the yield expected on the basis of the lifetimes given above, i.e., on the basis of the competition between reactions 15 and 5. In a similar vein, cyclooctane has been reported to be 2.8 times as reactive as cyclohexane with TBHP under GoAgg^V conditions at room temperature.^{8b} This is the expected reactivity ratio for alkoxyl radical chemistry since $k_5 = 2.7 \times 10^6 M^{-1} s^{-1} at 25 °C^{12}$ and $k(RO^{\bullet} + c-C_6H_{12}) = 1.2 \times 10^6 M^{-1} s^{-1} at 30 °C.¹⁶$

The conclusion is inescapable. $GoAgg^V$ chemistry is most succinctly explained in terms of classical free radical chemistry with initiation via alkoxyl radicals formed in reaction 1. Our results are congruent with those of Minisci et al^{9,19} and Newcomb et al²⁰ and our conclusions are the same: $GoAgg^V$ chemistry (and possibly a lot more of Gif chemistry) involves *free-radicals* and an Fe^{III} / Fe^{IV} couple, not a "*radical-free*" Fe^{III} / Fe^V couple.

Acknowledgment. We thank Dr. I.W.C.E. Arends for preliminary work on MPPH and the Association for International Cancer Research and the National Foundation for Cancer Research for partial support of this work.

REFERENCES AND NOTES

- 1. Issued as NRCC Nº. 39079.
- 2. Fourth year honours student, Department of Chemistry, Carleton University, Ottawa, Ontario, Canada.
- 3. Research Associate 1995-1996.
- 4. Ortiz de Montellano, P. R. Ed. Cytochrome P-450: Structure, Mechanism and Biochemistry; Plenum: New York, 1986.
- 5. Lippard, S. J. Acc. Chem. Res. 1994, 27, 229-236.
- 6. Barton, D. H. R.; Doller, D. Acc. Chem. Res. 1992, 25, 504-512.
- 7. Dioxygen from the hydroperoxide, e.g., reaction 4, and from the air plays a role as a secondary oxidant, see, e.g., Knight, C.; Perkins, M. J. J. Chem. Soc., Chem. Commun. 1991, 925-927.
- (a) Barton, D. H. R.; Chavasiri, W. Tetrahedron 1994, 50, 19-30. (b) Barton, D. H. R.; Bévière, S D.; Chavasiri, W. Tetrahedron 1994, 50, 31-46.
- 9. Minisci, F.; Fontana, F.; Araneo, S.; Recupero, F. Tetrahedron Lett. 1994, 35, 3759-3762.
- Tert-butoxyl radicals are acknowledged to be formed with Fe^{II} catalysts under Gif conditions, see:
 (a) Bardin, C.; Barton, D. H. R.; Hu, B.; Rojas-Wahl, R.; Taylor, D. K. Tetrahedron Lett, 1994, 35, 5805-5808.
 (b) Barton, D. H. R.; Chabot, B. M.; Delanghe, N. C.; Hu, B.; Le Gloahec, V. N.; Rojas Wahl, R. U. Tetrahedron Lett 1995, 36, 7007-7010.
- 11. Reagents were mixed at 0 °C, the temperature was raised to 25 °C and after 3 h at 25 °C part of the reaction mixture was analyzed directly by LC/MS mainly for the mixed peroxides (authentic standards for the TBHP, MPPH and CH reactions). Ph₃P was added to the remainding part of the mixture to convert *all* hydroperoxides to the corresponding alcohols, and then was followed by GC and GC/MS analysis.
- 12. Arends, I. W. C. E.; Ingold, K. U.; Wayner, D. D. M. J. Am. Chem. Soc. 1995, 117, 4710-4711.
- PhC(O)OO• is ca. 860 times as reactive in H-atom abstraction as HOO•. See: Zaikov, G. E.; Howard, J. A.; Ingold, K. U. Can J. Chem. 1969, 47, 3017-3029.
- 14. Laser flash photolysis (266 nm) of *o*-MCH in deoxygenated CH₃CN at 25 °C showed that the *o*-methyl cumyloxyl radical decayed with a rate constant of 2.5 x 10⁶ s⁻¹. We have previously estimated that $k_{11} \sim 2.2 \times 10^8 \text{ s}^{-1}$ at 40 °C in cyclopentane (see footnote 19 in ref. 12).
- 15. The lifetime is the reciprocal of the first-order rate constant for radical decay. Rate constants for β-scission increase with solvent polarity so lifetimes decrease.¹⁶ No alkoxyl radical lifetimes have been measured in pyridine / acetic acid mixtures. Lifetimes at 30 °C for *tert*-butoxyl¹⁷ and cumyloxyl¹⁶ are available in a number of pure solvents.
- 16. Avila, D. V.; Brown, C. E.; Ingold, K. U.; Lusztyk, J. J. Am. Chem. Soc. 1993, 115, 466-470.
- 17. For *tert*-butoxyl radicals the ratios of the rate constants for H-atom abstraction from cyclohexane to β -scission $(k_{\rm H} / k_{\beta})$ are available over a temperature range in several solvents.¹⁸ Cumyloxyl radicals abstract hydrogen from cyclohexane with $k_{\rm H} = 1.2 \times 10^6 \,{\rm M}^{-1} \,{\rm s}^{-1}$ at 30 °C in all solvents.¹⁶ With the assumption that $k_{\rm H}$ values are the same for these two alkoxyl radicals, k_{β} (and, hence, lifetimes) can be calculated for *tert*-butoxyl at 30 °C.
- 18. Walling, C.; Wagner, P. J. J. Am. Chem. Soc. 1964, 86, 3368-3375.
- (a) Minisci, F.; Fontana, F.; Araneo, S.; Recupero, F. J. Chem. Soc., Chem. Commun. 1994, 1823-1824.
 (b) Minisci, F.; Fontana, F. Tetrahedron Lett. 1994, 35, 1427-1430.
 (c) Minisci, F.; Fontana, F.; Araneo, S.; Recupero, F.; Banfi, S.; Quici, S. J. Am. Chem. Soc. 1995, 117, 226-232.
- 20. Newcomb, M.; Simakov, P. A.; Park, S.-U. accompying Letter in this issue.

(Received in USA 9 November 1995; revised 1 December 1995; accepted 3 December 1995)