Iron–carbene functionalities supported by a macrocyclic ligand: iron–carbon double bond stabilized by tetramethyldibenzotetraazaannulene

Alain Klose,^a Euro Solari,^a Carlo Floriani,^{*a} Nazzareno Re,^b Angiola Chiesi-Villa^c and Corrado Rizzoli^c

^a Institut de Chimie Minérale et Analytique, BCH, Université de Lausanne, CH-1015 Lausanne, Switzerland

^b Dipartimento di Chimica, Università di Perugia, I-06100 Perugia, Italy

^c Dipartimento di Chimica, Università di Parma, I-43100 Parma, Italy

We report an unprecedented entry to the chemistry of the iron-carbene functionality supported by an easily available macrocycle.

The metal–carbene unit is a key functionality both in organometallic and organic chemistry.¹ It is usually bonded to conventional ancillary ligands, with a few remarkable exceptions in metal–porphyrin chemistry,^{2–6} and particularly in the case of iron² and ruthenium.^{2b,3,5} Despite of considerable effort in this field, owing to the relevance of metal–porphyrin carbene derivatives in chemistry and biology,^{2,3} some facts are particularly surprising: (*i*) there is an absence of any modeling approach in macrocycle chemistry, *i.e.* the use of readily available macrocycles; (*ii*) the potential of the metal–carbene bonded to a macrocyclic structure is still to be explored; (*iii*) very few (three, at present) metal–porphyrin carbenes have been structurally characterized.^{2a,6}

Here, we report a straightforward synthesis of iron–carbenes stabilized by the easily accessible tetramethyldibenzotetraazaannulene ligand $(H_2 \text{tmtaa})^7$ along with their structural characterization in solution and in the solid state. The availability of tetramethyldibenzotetraazaannulene encourages the use of [Fe(tmtaa)] for developing iron–carbene chemistry based on this and other macrocycles. The synthesis was performed by reacting 2^{8a} with the corresponding diazoalkane (Scheme 1).

The reaction of **2** with Ph₂CN₂ in THF at -30 °C, and then at room temperature, gave high yields of **3** as a green crystalline solid.† Complex **2** showed a very high thermal stability and resistance to hydrolysis. The reaction of **3** with dioxygen gave, almost quantitatively, benzophenone and the well known μ -oxo dimer **4**.^{7a} The stability and the nature of the final compound derived from the reaction of 2 with diazoalkanes is strongly dependent on the substituents at the carbene carbon. In the case of PhCHN₂, the carbene derivative forms only at low temperature. It decomposes at room temperature leading to the starting material and trans-stilbene (81%) and cis-stilbene (19%), while in the presence of dioxygen it forms PhCHO and complex 4. Owing to its diamagnetism, 3 has been readily characterized by ¹H and ¹³C NMR spectroscopies and structural details have been revealed by X-ray analysis[‡] (see Fig. 1). The coordination environment of iron is a distorted tetragonal pyramid, unlike iron-prophyrin^{2a} or osmium-porphyrin^{6b} examples, where the metal is always six-coordinate. The metal is displaced by 0.335(1) Å from the N₄ average plane. The Fe-C(23) vector is perpendicular to the N₄ core, the dihedral angle with the normal to the N_4 plane being $1.0(1)^\circ$. The carbene plane C(23)C(31)C(41) is almost parallel to the $N(2)\cdots N(4)$ vector, the torsion angles C(31)-C(23)-Fe-N(4) and C(41)-C(23)-Fe-N(2) being -15.5(3) and $-15.2(3)^{\circ}$, respectively. The Fe-C(23) bond distance [1.794(3) Å] is particularly short compared to the only available iron-porphyrin [Fe=CCl₂ 1.83(3) Å]^{2a} or organometallic derivatives of iron, where it ranges from 1.978(3) to 1.85(3) Å.⁹ The question whether the oxidation state of iron in 3 is +2 or +4 can be reasonably answered, in the absence of Mössbauer measurements, by considering the Fe distance from the N₄ plane. A correlation in tmtaa complexes has been established between the out of plane distance of the metal and its dⁿ configuration.^{7b} The value found in the present case [0.335(1) Å] is expected for low-spin d⁶ five-coordinate iron(II), e.g. 0.29 Å in [Fe(tmtaa)(CO)].^{8c} Much longer distances would be found for a d⁴ configuration. The extended





Fig. 1 ORTEP drawing of complex **3** (50% probability ellipsoids. Selected interatomic distances (Å) and angles (°): Fe–N(1) 1.921(3), Fe–N(2) 1.926(2), Fe–N(3) 1.911(3), Fe–N(4) 1.920(2), Fe–C(23) 1.794(3), C(23)–C(31) 1.487(5), C(23)–C(41) 1.472(4); N(3)–Fe–N(4) 94.6(1), N(2)–Fe–N(4) 163.3(1), N(2)–Fe–N(3) 82.7(1), N(1)–Fe–N(4) 82.3(1), N(1)–Fe–N(3) 156.4(1), N(1)–Fe–N(2) 93.6(1), Fe–C(23)–C(41) 122.6(2), Fe–C(23)–C(31) 121.7(2), C(31)–C(43)–C(43) 115.7(3).

Chem. Commun., 1997 2297

Hückel calculations further support the +2 oxidation state for iron in **3** and the use of carbene rather than alkylidene terminology for the present complexes. The tmtaa, which shows the usual saddle shape conformation,⁷ has structural parameters quite close to those reported for the other iron complexes.⁸

We thank the 'Fonds National Suisse de la Recherche Scientifique' (Grant No. 20-46'590.96), Ciba Speciality Chemicals (Basel, Switzerland), and 'Fondation Herbette' (University of Lausanne, Switzerland) for financial support.

Footnotes and References

* E-mail: carlo.floriani@icma.unil.ch

† *Preparation* of **3**: To a cooled (−30 °C) THF solution (140 ml) of [Fe(tmtaa)]·thf^{8a} (1.79 g, 3.80 mmol) was added dropwise a THF solution (40 ml) of diphenyldiazomethane (0.78 g, 4.0 mmol). When warmed to room temperature the solution became red. Nitrogen gas was slowly given off. After stirring for 2 d, the green solution was concentrated *in vacuo* (20 ml) and then hexane (15 ml) added. A green crystalline solid was obtained (1.70 g, 75%). Crystals suitable for X-ray analysis were grown in toluene (Found: C, 74.53; H, 5.77; N, 9.91. C₃₅H₃₂FeN₄ requires C, 74.47; H, 5.71; N, 9.92%). ¹H NMR (200 MHz, C₆D₆, 25 °C): δ 6.88–6.76 (m, 8 H), 6.35–6.33 (m, 10 H), 5.13 (s, 2 H), 2.12 (s, 12 H). ¹³C NMR (100.6 MHz, C₆D₆, 25 °C): δ 313.2 (C carbene), 166.6, 157.9, 149.3, 127.1, 124.8, 120.9, 120.5, 119.7, 107.3, 67.8, 25.8, 23.2.

‡ *Crystal data* for **3**: C₃₅H₃₂FeN₄·C₇H₈, M = 656.7, triclinic, space group $P\overline{1}$, a = 11.670(3), b = 13.593(4), c = 11.234(3) Å, $\alpha = 106.72(2)$, $\beta = 96.21(2)$, $\gamma = 77.62(2)^\circ$, U = 1665.0(8) Å³, Z = 2, $D_c = 1.310$ g cm⁻³, F(000) = 692, Mo-Kα radiation ($\lambda = 0.7109$ Å), μ (Mo-Kα) = 4.87 cm⁻¹: crystal dimensions $0.08 \times 0.72 \times 0.85$ mm. The structure was solved by the heavy atom method and anisotropically refined for all the non-H atoms. All the hydrogen atoms were located from difference Fourier mags and introduced as fixed contributors in the last stage of refinement ($U_{\rm iso} = 0.05$ Å²). For 4428 unique observed reflections [$I > 2\sigma(I)$] collected at T = 133 K on a Rigaku AFC6S diffractometer ($5 < 2\theta < 50^\circ$) and corrected for absorption the final R is 0.059 ($wR_2 = 0.163$ for the 5293 reflections having I > 0 used in the refinement). All calculations were carried out on a QUANSAN Personal Computer equipped with an INTEL PENTIUM processor. CCDC 182/644.

 Transition Metal Carbene Complexes, VCH, Weinheim, 1983;
W. D. Wulff, in Advances in Metal–Organic Chemistry, ed. L. S. Liebeskind, Jay, London, vol. 1, 1989, pp. 209; J. W. Herndon, S. U. Tumer, L. A. McMullen, J. J. Matasi and W. F. K. Schnatter, in Advances in Metal–Organic Chemistry, ed. L. S. Liebeskind, Jay, London, vol. 3, 1994, pp. 51–95; Comprehensive Organometallic Chemistry, ed. G. Wilkinson, F. G. A. Stone and E. W. Abel, Pergamon, Oxford, vol. 12, 1995, ch. 5.1–5.5.

- 2 (a) D. Mansuy, M. Lange, J. C. Chottard, J. F. Bartoli, B. Chevrier and R. Weiss, Angew. Chem., 1978, 90, 828; (b) P.J. Brothers and J. P. Collman, Acc. Chem. Res., 1986, 19, 209; (c) D. Mansuy, Pure Appl. Chem., 1987, 59, 759; (d) I. Artaud, N. Gregoire, J.-P. Battioni, D. Dupre and D. Mansuy, J. Am. Chem. Soc., 1988, 110, 8714; (e) I. Artaud, N. Gregoire, P. Leduc and D. Mansuy, J. Am. Chem. Soc., 1990, 112, 6899; (f) D. Mansuy and J. P. Mahy, in Metalloporphyrins Catalyzed Oxidations, ed. F. Montanari and L. Casella, Kluwer, Dordrecht, 1994, p. 175.
- 3 J. P. Collman, P. J. Brothers, L. McElwee-White, E. Rose and L. J. Wright, *J. Am. Chem. Soc.*, 1985, **107**, 4570; J. P. Collman, P. J. Brothers, L. McElwee-White and E. Rose, *J. Am. Chem. Soc.*, 1985, **107**, 6110.
- 4 H. Brand and J. Arnold, *Coord. Chem. Rev.*, 1995, **140**, 137; T. Aida and S. Inoue, *Acc. Chem. Res.*, 1996, **29**, 39.
- 5 J. P. Collman, L. S. Hegedus, J. R. Norton and R. G. Finke, *Principles and Applications of Organotransition Metal Chemistry*, University Science Books, Mill Valley, CA, 1987.
- 6 (a) T. Boschi, S. Licoccia, R. Paolesse and P. Tagliatesta, Organometallics, 1989, 8, 330; (b) J.-P. Djukic, D. A. Smith, V. G. Young, Jr. and L. K. Woo, Organometallics, 1994, 13, 3020.
- 7 Some leading references to the use of [tmtaa]²⁻ as an ancillary ligand in the organometallic chemistry of the early transition metals are: (a) C. H. Jang, J. A. Ladd and V. L. Goedken, J. Coord. Chem., 1988, 18, 317; (b) S. De Angelis, E. Solari, E. Gallo, C. Floriani, A. Chiesi-Villa and C. Rizzoli, *Inorg. Chem.*, 1992, 31, 2520; (c) L. Giannini, E. Solari, C. Floriani, A. Chiesi-Villa and C. Rizzoli, *Angew. Chem., Int. Ed. Engl.*, 1994, 33, 2204; (d) L. Giannini, E. Solari, S. De Angelis, T. R. Ward, C. Floriani, A. Chiesi-Villa and C. Rizzoli, *J. Am. Chem. Soc.*, 1995, 117, 5801; (e) D. G. Black, D. C. Swenson and R. F. Jordan, *Organometallics*, 1995, 14, 3539; (f) H. Schumann, *Inorg. Chem.*, 1996, 35, 1808; (g) G. I. Nikonov, A. J. Blake and P. Mountford, *Inorg. Chem.*, 1997, 36, 1107.
- (a) V. L. Goedken, J. J. Pluth, S. M. Peng and B. Bursten, J. Am. Chem. Soc., 1976, 98, 8014; (b) M. C. Weiss, B. Bursten, S. M. Peng and V. L. Goedken, J. Am. Chem. Soc., 1976, 98, 8021; (c) V. L. Goedken, S. M. Peng, J. A. Molin-Norris and Y. Park, J. Am. Chem. Soc., 1976, 98, 8391; (d) M. C. Weiss and V. L. Goedken, Inorg. Chem., 1979, 18, 819; (e) P. Berno, C. Floriani, A. Chiesi-Villa and C. Guastini, J. Chem. Soc., Dalton Trans., 1988, 1409; (f) P. Berno, C. Floriani, A. Chiesi-Villa and C. Guastini, J. Chem. Soc., Dalton Trans., 1989, 551.
- 9 P. E. Riley, R. E. Davis, N. T. Allison and W. M. Jones, *Inorg. Chem.*, 1982, **21**, 1321; D. Bauer, P. Härter and E. Herdtweck, *J. Chem. Soc.*, *Chem. Commun.*, 1991, 829.

Received in Basel, Switzerland, 4th July 1997; 7/04736C