

# Chemical Science

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: J. Fox, Y. Fang, H. Zhang, Z. Huang, S. Scinto, J. Yang, C. W. A. Ende, O. Dmitrenko and D. S. Johnson, *Chem. Sci.*, 2018, DOI: 10.1039/C7SC04773H.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

## Photochemical Syntheses, Transformations, and Bioorthogonal Chemistry of *trans*-Cycloheptene and Sila *trans*-Cycloheptene Ag(I) Complexes

Yinzhi Fang,<sup>a</sup> Han Zhang,<sup>a,†</sup> Zhen Huang,<sup>b,†</sup> Samuel S. L. Scinto,<sup>a</sup> Jeffrey C. Yang,<sup>b</sup> Christopher W. Am Ende,<sup>c</sup> Olga Dmitrenko,<sup>a</sup> Douglas S. Johnson,<sup>b</sup> Joseph M. Fox<sup>a,\*</sup>

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

A photochemical synthesis of AgNO<sub>3</sub> complexes of *trans*-cycloheptene (TCH) and *trans*-1-sila-4-cycloheptene (Si-TCH) derivatives is described. A low temperature flow photoreactor was designed to enable the synthesis of carbocyclic TCH derivatives due to their thermal sensitivity in the absence of metal coordination. Unlike the free carbocycles, TCH•AgNO<sub>3</sub> complexes can be handled at rt and stored for weeks in the freezer (−18 °C). Si-TCH•AgNO<sub>3</sub> complexes are especially robust, and are bench stable for days at rt, and for months in the freezer. X-ray crystallography was used to characterize a Si-TCH•AgNO<sub>3</sub> complex for the first time. With decomplexation of AgNO<sub>3</sub> in situ, metal-free TCO and Si-TCH derivatives can engage in a range of cycloaddition reactions as well as dihydroxylation reactions. Computation was used to predict that Si-TCH would engage in bioorthogonal reactions that are more rapid than the most reactive *trans*-cyclooctenes. Metal-free Si-TCH derivatives were shown to display good stability in solution, and to engage in the fastest bioorthogonal reaction reported to date ( $k_2$  1.14 × 10<sup>7</sup> M<sup>−1</sup>s<sup>−1</sup> in 9:1 H<sub>2</sub>O:MeOH). Utility in bioorthogonal protein labeling in live cells is described, including labeling of GFP with an unnatural tetrazine-containing amino acid. The reactivity and specificity of the Si-TCH reagents with tetrazines in live mammalian cells was also evaluated using the HaloTag platform. The cell labeling experiments show that Si-TCH derivatives are best suited as probe molecules in the cellular environment.

### Introduction

For nearly seventy years,<sup>1–3</sup> the unusual bonding, reactivity and planar chirality of *trans*-cycloalkenes have captured the imagination of scientists. The unique reactivity of *trans*-cycloalkenes has produced an impressive collection of applications in synthesis, including reactions with dienes,<sup>4–6</sup> 1,3-dipoles<sup>7</sup> and ketenes.<sup>8–10</sup> Additionally, strained *trans*-cycloalkenes can serve as excellent ligands for transition metals. In the field of bioorthogonal chemistry,<sup>11–19</sup> *trans*-cycloalkenes hold special significance due to their particularly fast kinetics in cycloaddition reactions.<sup>20–23</sup> Relative to *trans*-cyclooctene, the chemistry of the homolog *trans*-cycloheptene is less explored. *trans*-Cycloheptene was first trapped with diphenylisobenzofuran by Corey and Winter from *trans*-1,2-cycloheptenethionocarbonate through treatment with P(OMe)<sub>3</sub>.<sup>24</sup> Studies by Marshall,<sup>25</sup> Kropp<sup>26</sup> and Beauchemin<sup>27</sup> on the

photoprotonation reactions of cyclic alkenes, including cycloheptene, have shown that *cis/trans* equilibria could be driven by selective addition reactions of *trans*-cycloalkenes.

*trans*-Cycloheptene was first spectroscopically characterized by Inoue via singlet sensitized photoisomerization of *cis*-cycloheptene at −35 °C.<sup>28,29</sup> Unlike *trans*-cyclooctene, which is stable at room temperature, *trans*-cycloheptene undergoes rapid isomerization under ambient conditions via a proposed ‘interrupted dimerization’ mechanism.<sup>30</sup> *trans*-Cycloheptene has also been prepared via ligand exchange from a *trans*-cycloheptene•CuOTf complex.<sup>31</sup> 1-Phenyl-*trans*-cycloheptene and *trans*-cycloheptenone derivatives are also known to be thermally unstable at ambient temperature, but can be trapped *in situ*.<sup>32–38</sup>

While the parent *trans*-cycloheptene is thermally labile, it has been demonstrated in several studies that metal complexes can be isolated. CuOTf has been proposed to catalyze photodimerization reactions of cyclic olefins via photoinduced *cis-trans* isomerization,<sup>39</sup> with predominant formation of a cyclotrimer from cycloheptene.<sup>40</sup> A stable *trans*-cycloheptene•CuOTf complex has been prepared through irradiation of *cis*-cycloheptene•CuOTf, but a yield for the process was not reported.<sup>41</sup> A pybox-RuCl<sub>2</sub> complex of *trans*-cycloheptene has been prepared by irradiation of the

<sup>a</sup> Brown Laboratories, Department of Chemistry and Biochemistry, University of Delaware, Newark, DE 19716.

<sup>b</sup> Pfizer Worldwide Research and Development, Cambridge, Massachusetts 02139.

<sup>c</sup> Pfizer Worldwide Research and Development, Groton, Connecticut 06340.

<sup>†</sup> equal contributions.

Electronic Supplementary Information (ESI) available: full procedures, computational details and characterization data. See DOI: 10.1039/x0xx00000x



## ARTICLE

## Journal Name

corresponding ethylene complex in the presence of *cis*-cycloheptene under singlet sensitized conditions.<sup>42</sup> Jendralla described the preparation of AgOTf and AgClO<sub>4</sub> complexes of 3-methoxy-*trans*-cycloheptene and 6-Methoxy-(*Z*),4(*E*)-cycloheptadiene.<sup>43,44</sup> These compounds can be prepared through the Ag-mediated ring opening of a nitrosoarene derivative of bicyclo[4.1.0]heptane. In unspecified yields, the AgClO<sub>4</sub>•3-methoxy-*trans*-cycloheptene complex was combined with a number of dienes to give the products of metal decomplexation and [4+2] cycloaddition.<sup>43,44</sup>

Because C-Si bonds are long, the inclusion of silicon into the cyclic backbone can alleviate olefinic strain and impart stability to *trans*-cycloalkenes.<sup>45–52</sup> In 1997, (*E*)-1,1,3,3,6,6-hexamethyl-1-sila-4-cycloheptene was synthesized, resolved, and characterized crystallographically.<sup>46,47</sup> Here, the exhaustive allylic substitution imparts a high degree of stability to the *trans*-alkene. Recent studies by Woerpel<sup>48–51</sup> and Tomooka<sup>52</sup> have provided demonstrations of the utility of backbone heteroatom-containing *trans*-cycloalkene derivatives in synthesis. Woerpel has elegantly developed a general method for the preparation of *trans*-oxasilacycloheptenes—seven membered rings that contain *trans*-alkenes and siloxy bonds in the backbone.<sup>48–51</sup>

There has been little investigation into the reaction chemistry of *trans*-cycloheptene derivatives. In 1980, Jendralla showed that silver complexes of 3-methoxy-*trans*-cycloheptene are isolable, and can reversibly dissociate and in unspecified yields undergo cycloaddition reactions.<sup>43,44</sup> Woerpel has described selective addition reactions and difunctionalization reactions of *trans*-oxasilacycloheptenes, and recently has reported the Diels-Alder reactions of *trans*-oxasilacycloheptenes with furan and tetrazine derivatives.<sup>48–51</sup> At 25 °C in benzene, an oxasilacycloheptene was 7-fold more reactive toward 2,5-diphenylisobenzofuran than a conformationally strained *trans*-cyclooctene ('s-TCO') derivative.<sup>48</sup> An oxasilacycloheptene derivative was shown to react with 3,6-diphenyl-s-tetrazine in benzene at rt in less than 10 min and in 90% NMR yield.<sup>48</sup>


Our group has described a closed-loop flow reactor for the synthesis of *trans*-cyclooctene derivatives, whereby selective complexation with AgNO<sub>3</sub> is used to drive the formation of *trans*-isomer from *cis*-cyclooctene.<sup>21,53–55</sup> Described herein is an approach to the synthesis of *trans*-cycloheptene (TCH) and *trans*-1-sila-4-cycloheptene (Si-TCH) derivatives via flow photochemical synthesis. The derivatives described are especially stable as their AgNO<sub>3</sub> metal complexes, which can be stored neat or in solution for long periods. With decomplexation of AgNO<sub>3</sub> *in situ*, metal-free TCH and Si-TCH derivatives can engage in a range of cycloaddition reactions as well as dihydroxylation reactions. Unlike the carbocycles, Si-TCH derivatives display good stability in solution and are shown to engage in the fastest bioorthogonal reaction reported to date. Decomplexation of AgNO<sub>3</sub> can be carried out

*in situ* directly in cell media for bioorthogonal protein labeling in live cells.

## Results and discussion


For the photochemical synthesis of *trans*-cyclooctene derivatives, the *trans*-cycloalkene is first scavenged on AgNO<sub>3</sub>/SiO<sub>2</sub> and then liberated through treatment with aqueous or methanolic ammonia.<sup>21,53–55</sup> However, our attempts to directly apply this procedure to the synthesis of carbocyclic TCHs was unsuccessful, most likely due to the susceptibility of carbocyclic TCHs to readily isomerize to their *cis*-isomers.<sup>30</sup> Recently, we demonstrated that the shelf-life of conformationally strained *trans*-cyclooctenes could be enhanced by storing the cycloalkenes as their AgNO<sub>3</sub> complexes, and that the free alkenes could be liberated *in situ* through treatment with NaCl in aqueous solution or cell media.<sup>56</sup> We reasoned that TCH and Si-TCH derivatives may also be isolable and better stored as their Ag-complexes, and that the corresponding free-alkenes could be liberated at later time points as required.

Photoisomerizations to form Si-TCH•AgNO<sub>3</sub> derivatives were carried out at rt using the previously described flow-photoisomerization apparatus,<sup>21,53,54</sup> with the modification that Si-TCH•AgNO<sub>3</sub> complexes were directly isolated from SiO<sub>2</sub> without Ag-decomplexation. The metal complexes that were obtained were stable in neat form for >1 month in the freezer. However, it was necessary to alter our reactor design for the synthesis of carbocyclic *trans*-cycloheptenes due to their thermal lability (Figure 1). As with the standard photoreactor, this system uses methyl benzoate as a sensitizer and a



**Figure 1** The apparatus for photoisomerization of carbocyclic *trans*-cycloheptene derivatives was designed to minimize loss of *trans*-cycloheptene due to thermal isomerization by using FEP tubing and inline cooling. For the synthesis of *trans*-1-sila-4-cycloheptenes, photoisomerizations could be carried out at rt using a conventional flow photoisomerization setup.



**Scheme 1.** Flow-photochemical synthesis of AgNO<sub>3</sub> complexes of *trans*-cycloheptenes and *trans*-1-sila-4-cycloheptenes.

TCH•AgNO<sub>3</sub> and Si-TCH•AgNO<sub>3</sub> complexes were isolated as semisolids that contained 20–30% free AgNO<sub>3</sub>. Yields are the average of two runs, and were corrected by measuring the <sup>1</sup>H NMR against an internal standard. <sup>a</sup>Photoisomerization of carbocycles were conducted at low temperature using the flow apparatus described in Figure 1. Photoisomerizations of silacycles were conducted at room temperature using the flow apparatus previously described for *trans*-cyclooctene synthesis.

metering pump to pass a solution of substrate at a rate of 100 mL/min through a photowell and then through a column of AgNO<sub>3</sub>•SiO<sub>2</sub>, where the *trans*-cycloalkene is selectively captured as a AgNO<sub>3</sub> complex. In the reactor for carbocyclic TCH synthesis, a reservoir of solvent chilled in a cold bath (–50 °C) was positioned before the photowell, and the photoisomerization was conducted in a coil of optically transparent FEP tubing.<sup>57</sup> The fluoropolymer tubing provides a high surface area and minimal volume (only 30 mL for 8 m tubing) thereby minimizing the residence time before product adsorption on AgNO<sub>3</sub>/SiO<sub>2</sub>. In our standard setup, an inline thermometer was included to measure the temperature for

the flowing mixture either before or just after the UV lamp. The temperature was measured as 0 °C before entering the Rayonet photoreactor, and as 20 °C after exiting the photoreactor. With this apparatus, TCH•AgNO<sub>3</sub> complexes were eluted from the column, and isolated as semisolids that are moderately stable at r.t. but stable for weeks in the freezer.

The scope of TCH and Si-TCH synthesis is shown in Scheme 1. With the exception of cycloheptene itself, the carbocyclic and sila-cycloheptene precursors were prepared in 2–7 steps using olefin metathesis as a key step (Supporting information). Silver nitrate complexes of *trans*-cycloheptene (**1a**) and *trans*-5-hydroxymethylcycloheptene (**1b**) were prepared in 53% and 64% yields, respectively. These TCH•AgNO<sub>3</sub> complexes are stable enough to handle on the bench for modest periods (hours), and to longer-term storage in the freezer (–18 °C). NMR monitoring showed 90% fidelity for a CD<sub>3</sub>OD solution of **1a** after 10 days storage in freezer (–18 °C), and 92% fidelity for a CD<sub>3</sub>OD solution of **1a** after 10 hours at rt on the bench.


The silver complexes of Si-TCH are much more stable. NMR monitoring showed 93% fidelity for a CD<sub>3</sub>OD solution of **2a** after 8 days at rt, and 96% fidelity for **2a** after storage for 1 month in the freezer (–18 °C, neat). The photoisomerization method could be used to produce diphenyl (**2a**) or dialkyl (**2b–g**) substituted silacycles. Cyano (**2b**) and hydroxyl (**2c–e**) groups were tolerated, as were NHS ester (**2f**) and chloroalkane (**2g**) groups that could be used to enable conjugation to fluorophores and HaloTag<sup>58,59</sup> fusion proteins, respectively. TCH•AgNO<sub>3</sub> and Si-TCH•AgNO<sub>3</sub> complexes were isolated as semisolids that contained 20–30% free AgNO<sub>3</sub>. The isolated yields were corrected by measuring the <sup>1</sup>H NMR against an internal standard.

Previously, a crystal structure showed that *trans*-cyclooctene coordinates with AgNO<sub>3</sub> as a 1:1 complex.<sup>60</sup> X-ray quality crystals of silver complex **2a** were grown from ethyl acetate/methanol. Selected bond lengths and angles are displayed in Figure 2. The coordination environment at silver is distorted trigonal, with bridging coordination of nitrite in an extended polymeric structure in the solid state. For comparison, we also grew crystals of the silver (I) nitrate complex of the equatorial diastereomer of 5-hydroxy-*trans*-cyclooctene **3** (Figure 2). Here, the coordination environment at silver is distorted tetrahedral due to the ability of the hydroxyl to serve as a second bridging ligand. The C–Ag bond lengths and bond angles were similar for **2a** and **3**. As expected, the C=C–C–C dihedral angle for **3** (136.7 °) is smaller than metal-free TCO **4** (139.1 °).<sup>53</sup> Similarly, the C=C–C–C dihedral angle for **2a** (126.3 °) was smaller than that of metal free Si-*trans*-cycloheptene **5** (130.9 °),<sup>46</sup> but comparable to **6** (126.1 °)<sup>50</sup> — a compound with additional strain due to relatively short C–O and Si–O bonds in the cyclic backbone. Reflecting the low level of metal backbonding that is common for Ag(I) alkene complexes, the C=C bond length for the Ag(I)



## ARTICLE

complexes (1.329 Å for **2a**, 1.366 Å for **3**) was very similar to that of the metal-free complexes **4–6** (1.331–1.335 Å).




**Figure 2** X-ray crystal structures of SiTCH•AgNO<sub>3</sub> and TCO•AgNO<sub>3</sub> complexes, with comparisons to known metal-free *trans*-cycloalkenes that have been crystallographically characterized.

Silver-free Si-TCH compounds could be prepared by treating their corresponding silver nitrate complexes with an excess of aq. NH<sub>4</sub>OH or aq. NaCl, followed by extraction with organic solvent. For example, silver complex **2a** upon treatment with aq. NH<sub>4</sub>OH was extracted with C<sub>6</sub>D<sub>6</sub> to give a solution of Si-TCH **7a** (98% *trans* isomer). Consistent with previous reports on a biomolecular mechanism for TCH isomerization,<sup>30</sup> variable amounts (20–30%) of the *cis*-isomer of **7a** was observed when **7a** was concentrated to dryness on the rotovap. However, **7a** displayed high stability when maintained in solution, with only 8% isomerization observed for a 100 mM solution of **7a** that was stored for 24 hours at room temperature, and <5% isomerization for a similar solution that was stored for 24 hours in a freezer (−20 °C).

Ag(I)-complex **2d** could also be freed of metal to give alkene **7d** as a mixture of diastereomers (Figure 3A). The allylic substituents of **7d** protect the alkene from biomolecular chemistry, and unlike most other metal-free Si-TCH compounds, **7d** is stable when stored in neat form and can be characterized by FT-IR (Figure 3B). The weak C=C double bond stretch of **7d** at 1624 cm<sup>−1</sup> is shifted to 1559 cm<sup>−1</sup> for the Ag(I)

complex **2d**. This 65 cm<sup>−1</sup> shift is consistent both in magnitude and direction for a Ag(I) alkene complex.<sup>61</sup> Finally, we noted



**Figure 3** (a) Decomplexation of AgNO<sub>3</sub> with NH<sub>4</sub>OH gives metal-free Si-TCH **7a** which is stable to overnight storage in solution. Aq. NaCl can also be used to liberate metal from SiTCH•AgNO<sub>3</sub> complexes. The metal-free complexes are stable in solution, but give variable amounts of isomerization when concentrated to dryness. (b) FT-IR spectrum of the C=C stretches of metal complex **2d** and free alkene **7d**. (c) Diagnostic shifts in alkene peaks in the <sup>1</sup>H and <sup>13</sup>C NMR spectra of metal complex **2b** and metal-free **7b**.

that Ag(I)-complexation leads to signature shifts of alkene resonances in both the <sup>1</sup>H and <sup>13</sup>C NMR spectra (Figure 3C). For example, alkene resonances in the <sup>1</sup>H NMR spectrum of metal complex **2b** were shifted downfield relative to metal-free **7b** by 0.15 ppm, while in <sup>13</sup>C NMR spectra alkene resonances of **2b** were shifted up field by ~16 ppm. In the <sup>1</sup>H NMR spectra, complexity arises due to the higher order effects for a ddd couplet.


TCH **1a** and Si-TCH **2a** were shown to engage in a range of reactions as shown in Scheme 2. Metal complex **1a** was directly combined with 3,6-diphenyl-1,2,4,5-tetrazine to give pyridazine **8** — the product of metal dissociation, Diels-Alder/retro-Diels-Alder, and oxidation, in 98% yield. Cyclopenta-1,3-diene was also used to trap *trans*-



cycloheptene, delivering the [4+2] cycloaddition adduct **9** in 81% yield as a single diastereomer. We also investigated the vicinal dihydroxylation of **1a**, and found that catalytic OsO<sub>4</sub> and NMO gave **10** in 82% yield as a single diastereomer. The observation that the dihydroxylation of **1a** is stereospecific is in line with earlier observations by Cope with *trans*-cyclooctene.<sup>2</sup>

Si-TCH•AgNO<sub>3</sub> complex **2a** could be freed from silver by treatment with NH<sub>4</sub>OH (Scheme 2), and subsequently combined with cyclopentadiene, diazomethane, dichloroketene, and benzyl azide to provide the cycloadducts **11–14** in 76%–96% yields (Scheme 2). In each case, a single diastereomer was obtained. Attempts to combine *cis*-1-diphenylsila-4-cycloheptene (the *cis*-isomer of **7a**) with dichloroketene, benzylazide or diazomethane only returned unreacted starting material. An X-ray structure was obtained for the dichloroketene adduct **13** (Supporting Information).


Scheme 2. Reactions of TCH **1a** and Si-TCH **2a**.



<sup>a</sup>Yields represent isolated yields as average of two runs


We sought to demonstrate that Si-TCH cycloadducts could be oxidized to give formal cycloadducts of 1,2-dialkyloléfins—which are recalcitrant substrates in intermolecular Diels-Alder reactions.<sup>62</sup> With the cycloadduct **11**, we demonstrated that the diphenylsila- group could be oxidized to a diol-product. Thus **11** was subjected to Tamao-Fleming reaction under Woerpel's conditions<sup>63</sup> to give the unknown diol product **15** in 76% yield (Scheme 3).

Scheme 3. Fleming-Tamao Oxidation



We further demonstrated that NHS-ester **2f** could be modified through coupling to a BODIPY-fluorophore conjugate. As shown in Scheme 4, NHS ester **2f** could be decomplexed from AgNO<sub>3</sub> by treatment with brine and extraction into CH<sub>2</sub>Cl<sub>2</sub>. The resulting free Si-TCH was then conjugated to an aminohexyl BODIPY **16**, and the resulting conjugate was isolated and stored as a AgNO<sub>3</sub> complex (**AgSiTCH-BODIPY**). As discussed below, this fluorophore conjugate finds utility for bioorthogonal labeling in live cells. We also demonstrated that the equatorial allylic alcohol **17**(eq), derived from Ag-complex **2e**, could be elaborated to the carbamate **18** through treatment with benzylisocyanate (Scheme 4). *trans*-Cyclooctenes with allylic carbamate leaving groups have been used by Robillard<sup>64,65</sup> and Chen<sup>66,67</sup> for the tetrazine-ligation initiated decaging of doxorubicin and other cargo molecules. The 7-membered analog **18** is particularly stable as the Ag-free *trans*-cycloalkene, and can be handled neat and stored without AgNO<sub>3</sub> in the freezer for long periods. Efforts to synthesize and explore the ability of analogs of **18** to function for payload release is a topic of ongoing study.

Scheme 4. Transformations of Si-TCH derivatives



Anticipating that Si-TCH compounds may be useful in bioorthogonal chemistry, computation was used to study the




## ARTICLE

## Journal Name


structure and Diels-Alder reactivity of Si-TCH **19** (Scheme 5). We compared **19** to the parent *trans*-cyclooctene and a conformationally strained 's-TCO' derivative **20** that was previously studied in our labs.<sup>21,54</sup> We reasoned that the 7-membered ring in the backbone of **19** would augment the olefinic strain of the *trans*-cycloalkene, and thereby increase the reactivity in tetrazine ligation. Previously, thiacycloheptynes and dibenzoselenocycloheptynes have been studied in bioorthogonal chemistry.<sup>67</sup> As shown in Scheme 5A, ground state calculations were carried out for Si-TCH **19** at the M06L/6-311+G(d,p) level, and compared to previous calculations on *trans*-cyclooctene and s-TCO **20**.<sup>21,54</sup> The calculated C=C-C dihedral angle for **19** (128.8°) is smaller than that for *trans*-cyclooctene (137.7°) or **20** (131.9°). M06L/6-311+G(d,p) calculations were also carried out to compare the Diels-Alder reactivity of Si-TCH **19** to *trans*-cyclooctene and s-TCO **20** (Scheme 5B). These calculations were carried out with diphenyl-s-tetrazine so that they could be benchmarked against previous calculations.<sup>21,54</sup> For Si-TCH **19**, the calculated barriers relative to a pre-reaction complex for the Diels-Alder reaction with 3,6-diphenyl-s-tetrazine are  $\Delta G^\ddagger$  12.48 kcal/mol;  $\Delta E^\ddagger$  9.27 kcal/mol,  $\Delta E^\ddagger(\text{ZPE})$  9.91 kcal/mol and  $\Delta H^\ddagger$  8.87 kcal/mol. The barrier is significantly lower than that of *trans*-cyclooctene with 3,6-diphenyl-s-tetrazine ( $\Delta\Delta G^\ddagger$  -3.61 kcal/mol,  $\Delta\Delta E^\ddagger$  -4.02 kcal/mol,  $\Delta\Delta E^\ddagger(\text{ZPE})$  -3.99 kcal/mol,  $\Delta\Delta H^\ddagger$  4.06 kcal/mol). The barrier is also lower than that calculated for s-TCO **20** ( $\Delta\Delta G^\ddagger$  -0.26 kcal/mol,  $\Delta\Delta E^\ddagger$  -0.92 kcal/mol,  $\Delta\Delta E^\ddagger(\text{ZPE})$  -0.59 kcal/mol,  $\Delta\Delta H^\ddagger$  -0.72 kcal/mol).

**Scheme 5.** (A) Ground state calculations on Si-TCH **19** show that the <C=C-C dihedral angle is more distorted than that of *trans*-cyclooctene or the conformationally constrained **20**. (B) Transition state calculations predict that Si-TCH **19** would be more reactive than *trans*-cyclooctene and **20**.

## A Ground state calculations (M06L/6-311+G(d,p))




## B Transition state calculations (M06L/6-311+G(d,p))



The computations prompted us to conduct experimental investigations into the utilization of Si-TCH complexes for applications in bioorthogonal chemistry.

Previously, we had studied the reaction of diphenyl-s-tetrazine in MeOH at 25 °C with s-TCO,<sup>21,54</sup> d-TCO,<sup>22,21</sup> and *trans*-cyclooctene,<sup>69</sup> with second order rate constants of 3100 (+/- 50), 520 (+/- 3) and 19.1 (+/- 0.2) M<sup>-1</sup>s<sup>-1</sup>, respectively. Si-TCH **7c** was liberated from AgNO<sub>3</sub>, and in agreement with computational prediction, was found to react with diphenyl-s-tetrazine with a rate constant of 4360 (+/- 430) M<sup>-1</sup>s<sup>-1</sup> — 1.4 times faster than sTCO, 8.4 times faster than dTCO, and 228 times faster than *trans*-cyclooctene itself (Scheme 6A).

Under aqueous conditions, tetrazine ligations are accelerated due to the hydrophobic effect. We previously had studied the



**Scheme 6.** Second order rate constants ( $k_2$ ) were determined with a stopped-flow spectrophotometer under pseudo-first order conditions using excess *trans*-cycloalkene and limiting tetrazine. (A) The kinetics of the cycloaddition of 3,6-diphenyl-s-tetrazine in MeOH at 25 °C with Si-TCH **7c** was measured by stopped flow spectroscopy with UV-vis monitoring at 295 nm, and compared to previously reported rate constants for s-TCO **21**, d-TCO **22** and *trans*-cyclooctene. (B) Stopped flow kinetics with a TAMRA-3,6-dipyridyl-s-tetrazine conjugate were monitored in 9:1 H<sub>2</sub>O:MeOH with monitoring by fluorescence (Ex: 556 nm; Em: 576 nm). Data points are shown in red, and the fit is shown in blue. (C) Second order rate constants ( $k_2$ ) were determined by plotting  $k_{\text{obs}}$  vs. the concentration of Si-TCH




reactions of s-TCO, d-TCO and *trans*-cyclooctene with a water soluble dipyrindyl-s-tetrazine derivative under stopped flow conditions in water with UV-vis monitoring.<sup>21</sup> In the most rapid example, a rate constant of  $3.3 \times 10^6 \text{ M}^{-1}\text{s}^{-1}$  was measured for an s-TCO derivative.<sup>21</sup> However, our attempts to conduct a similar measurement using Si-TCH **7c** were complicated because the reaction was complete before we could collect data even with stopped flow monitoring. To enable the measurement, we synthesized a fluorescent tetrazine-TAMRA conjugate **23**, and used fluorescence 'turn-

on'<sup>70</sup> to monitor reaction progress. The fluorogenic reaction enabled reaction monitoring at much lower concentrations (down to  $3 \mu\text{M}$  in tetrazine). As shown in Scheme 6B and 6C, the reaction of **23** with Si-TCH **7c** in 9:1 water:MeOH proceeds at  $25^\circ\text{C}$  with a second order rate constant  $k_2$   $1.14 \times 10^7$  ( $\pm 5 \times 10^5$ )  $\text{M}^{-1}\text{s}^{-1}$ . This is the fastest rate constant reported to date for a bioorthogonal reaction.

We also studied the in vitro and in vivo cycloaddition of SiTCH and a green fluorescent protein with an unnatural tetrazine-containing amino acid (sfGFP-150Tet-v.2.0, referred to as **GFP-Tet**), encoded via the procedure of Mehl and coworkers.<sup>71</sup> Thus, 4-(6-methyl-s-tetrazin-3-yl)phenylalanine was site-specifically introduced into a C-terminally hexahistidine-tagged GFP (sfGFP-150TAG-His6) via orthogonal translation using the evolved aminoacyl-tRNA synthetase MjRS/tRNACUA pair. Co-expression of these components in *E. coli* resulted in the amino acid-dependent synthesis of full-length recombinant **GFP-Tet** (Scheme 7A). The reaction of **GFP-Tet** with dienophiles is fluorogenic, and it is therefore possible to determine the reaction kinetics by monitoring the increase in GFP fluorescence.

Although **GFP-Tet** is highly reactive, with in vitro rates as fast as  $87,000 \text{ M}^{-1}\text{s}^{-1}$  toward sTCO, **GFP-Tet** is not as rapid as **23** due to the less reactive nature of the tetrazine. Kinetic measurements were carried out with silver free Si-TCH **7c**, which was obtained by treating Ag-complex **2c** with  $\text{NH}_4\text{OH}$  and extracting with ether. The second order rate constant of the reaction between Si-TCH **7c** and **GFP-Tet** was determined to be  $250,000 \pm 15,000 \text{ M}^{-1}\text{s}^{-1}$  in PBS at rt (Scheme 7B). The reaction was quantitative under these conditions as determined by ESI-MS (Scheme 7C), and is the fastest rate measured to date for **GFP-Tet**—2.9 times faster than previously measured rate constant for sTCO.<sup>71</sup> Si-TCH **18** was also shown to display very rapid labeling of **GFP-Tet** when carried out in live bacteria. The kinetics of the in vivo tetrazine ligation were monitored in a suspension (PBS) of *E. coli* overexpressing **GFP-Tet** by measuring the increase in whole-cell fluorescence upon addition of **7c**. At room temperature, a second-order rate constant of  $155,000 \pm 20,000 \text{ M}^{-1}\text{s}^{-1}$  was measured for the in vivo reaction, which is 62% as rapid as the in vitro ligation. The modest reduction in rate is in line with previous observations with TCO-based dienophiles.<sup>21,71,72</sup> Quantitative determination of the bioorthogonal reaction was verified by cell washing, lysis, purification by IMAC, and analysis by ESI-MS. Thus, Si-TCH **7c** is capable of crossing the bacterial cell membrane and engaging in rapid, high yielding conjugation inside a living cell.

The reactivity and specificity of the SiTCH reagent with tetrazines in live mammalian cells was evaluated using the HaloTag platform<sup>56</sup>, which we have previously used to benchmark the efficiency of various bioorthogonal reactions in the cellular environment. We have previously shown that *trans*-cycloalkene- $\text{AgNO}_3$ -complexes liberate  $\text{AgNO}_3$  immediately in cell media due to the high NaCl content, and




**Scheme 7.** (A) **GFP-Tet** reacts with **7c** to give a Diels-Alder adduct. Because the tetrazine quenches GFP fluorescence, the reaction progress can be measured by monitoring fluorescence increase. (B) Stopped-flow kinetic data for the reaction of **GFP-Tet** (50 nM) with **7c** (1.96  $\mu\text{M}$ ) at  $25^\circ\text{C}$  in PBS with fluorescence monitoring (Ex: 488 nm; Em: 506 nm). Data points are shown in green, and the fit is shown in black. A second order rate constant ( $k_2$ ) was determined by plotting  $k_{\text{obs}}$  vs. the concentration of Si-TCH **7c**. (C) Quantitative determination of the cycloadduct was confirmed by ESI-MS. Deconvoluted Mass spectra of **GFP-Tet** (grey) and the cycloadduct (green) are displayed and show the expected mass shift of 170 Da. Smaller peaks in each spectrum correspond to proteins that are truncated by a methionine.



perform identically to their metal-free analogs in cell labeling experiments.<sup>56</sup> Thus, we synthesized chloroalkane derivatives of SiTCH (**AgSiTCH-Halo**) and methyl-tetrazine (**MeTz-Halo**) and used these clickable HaloTag ligands to covalently label HaloTag protein expressed in HEK 293T cells with the clickable tag. In a competitive pulse-chase experiment, it was shown that 10  $\mu$ M of these HaloTag ligands completely blocked incorporation of BODIPY-Halo substrate (Fig S25), and therefore this concentration was used for subsequent experiments. Next we evaluated the tetrazine ligation of SiTCH in mammalian cells by reacting the **AgSiTCH-Halo** and **MeTz-Halo** protein conjugates with the corresponding **MeTz-BODIPY** or **AgSiTCH-BODIPY** fluorescent probes (300 nM) for different times (2–90 min) (Scheme 8, Fig S26). The reaction


was quenched at the various time points by chasing with excess non-fluorescent tetrazine-amine (method 1) or TCO-amine (method 2) (Fig S27), and in-gel fluorescence was used to quantify conversion vs time.

We found that the reaction of **SiTCH-Halo** with **MeTz-BODIPY** was complete within 15 minutes (Scheme 8B, Method 2). Interestingly, the reaction appeared to be slower with the reverse pairing where **SiTCH-BODIPY** was reacted with **MeTz-Halo** and did not reach saturation until 90 minutes (Scheme 8B, Method 1). We believe this is due to the suboptimal permeability and nonspecific protein binding of **AgSiTCH-BODIPY** which results in lower free concentrations available for the reaction. Compared to **MeTz-BODIPY** or fluorescent 5-



**Scheme 8.** Reactions of AgSiTCH with 3-methyl-6-s-tetrazine derivatives in live HEK 293T cells expressing HaloTag. (A) Cells were treated with 10  $\mu$ M **MeTz-Halo** (method 1) or **AgSiTCH-Halo** (method 2) for 30 min, followed by two 30 min washings and treatment with 300 nM **AgSiTCH-BODIPY** (method 1) or 300 nM **MeTz-BODIPY** (method 2) for the indicated time. (B) Analyzed by in-gel fluorescence and Western blot for each method: in-gel fluorescence is shown on the top panel, and Western blot with an anti-HaloTag antibody is on the bottom panel. (C) Time course for the two reactions in B. In-gel fluorescence intensity from  $n = 3$  independent replicates was normalized by the corresponding Western blot signal, and was fit to a one-phase exponential equation. Data were plotted as mean  $\pm$  SEM.





**Scheme 9. Live cell stability and imaging studies of SiTCH-Halo.** (A) In the stability experiment, HEK293T cells expressing HaloTag were treated with 10  $\mu$ M **Ag-sTCO-Halo** or **AgSiTCH-Halo** for 30 min, followed by two 30 min wash periods. Control cells were treated with 2  $\mu$ M **TAMRA-Halo**. At 0, 7, and 24 h time points, 2  $\mu$ M **TAMRA-Tz** was added and allowed to react for 1 h before analysis. (B) Cellular stability of **Ag-sTCO-Halo** and **AgSiTCH-Halo**. In-gel fluorescent intensities were normalized by the corresponding Western blot signals, and were subsequently normalized by the value of **TAMRA-Halo** at time 0. Data were plotted as mean  $\pm$  SEM. (C) Live cell images of HeLa cells expressing Halo-H2B-GFP. Cells were labeled with 10  $\mu$ M **Ag-sTCO-Halo** or **AgSiTCH-Halo** for 30 min, followed by two 30 min wash periods, before the treatment with 1  $\mu$ M **TAMRA-Tz** for 5 min. The reactions were quenched with 100  $\mu$ M TCO-amine for 10 min, followed by two 30 min wash periods prior to addition of 8  $\mu$ M Hoechst 33342 to visualize the nuclei and confocal imaging. Signals of Hoechst, GFP, and TAMRA were shown in blue, green, and red, respectively. Colocalization was demonstrated in composite images of GFP and TAMRA. Scale bar = 25  $\mu$ m.

hydroxy-*trans*-cyclooctenes,<sup>56</sup> **AgSiTCH-BODIPY** also resulted in more nonspecific protein labeling (Fig S26). However, compared to in-gel fluorescence with a TAMRA-labeled bicyclononyne (BCN),<sup>56</sup> **AgSiTCH-BODIPY** appears to be more selective.

To investigate in vivo stability of incorporated SiTCH probes in the intracellular environment, we compared HaloTag protein tagging by **AgSiTCH-Halo** to the previously described **Ag-sTCO-Halo**.<sup>56</sup> Subsequently the **TAMRA-Tz** fluorescent probe was attached via the tetrazine ligation (Scheme 9A). As a benchmark, we also compared directly incorporated **TAMRA-Halo** without a second bioorthogonal step. In these experiments, HaloTag-transfected cells were treated with 10  $\mu$ M HaloTag ligands for 30 min, followed by two 30 min wash periods. After this initial 90 min exposure to cells, time course experiments were carried out to test the intracellular stability of the sTCO- and SiTCH-labeled HaloTag proteins. Thus, cells were allowed to incubate for up to 24 additional hours, and cells initially tagged by **AgSiTCH-Halo** or **Ag-sTCO-Halo** were then treated with **TAMRA-Tz** at the indicated time points. Loss of fluorescence intensity relative to the **TAMRA-Halo** benchmark indicated instability of the bioorthogonal protein tag.

Compared to the **TAMRA-Halo** ligand, the **Ag-sTCO-Halo** tagged protein displayed 84% fluorescence intensity when the **TAMRA-Tz** probe was attached immediately after the 30 min HaloTag-labeling and two 30 min wash periods (Scheme 9B, Fig S28 and S29). By contrast, **AgSiTCH-Halo** showed only 50% fluorescence intensity at this initial timepoint. After incubation for 7 h and 24 h, **Ag-sTCO-Halo** tagged protein showed 61% and 43% fluorescence intensity—very similar to the **TAMRA-Halo** benchmark. However, the **AgSiTCH-Halo** tagged protein displayed only 10% fluorescence after 7 h, and negligible fluorescence after 24 h. The reduced cellular labeling efficiency with **AgSiTCH-Halo** could also be observed in live cell images according to the published protocol (Scheme 9C, Fig S30).<sup>56</sup> Specifically, HeLa cells overexpressing Halo-H2B-GFP were labeled with **AgSiTCH-Halo** or **Ag-sTCO-Halo**, and after two 30 min wash periods, treated with **TAMRA-Tz** for imaging. In both cases, specific fluorescent signals were observed in the nucleus with good colocalization with GFP signals. When **AgSiTCH-Halo** was used, however, the fluorescent intensity was only about half of that achieved with **Ag-sTCO-Halo**, confirming that the reduced stability of **SiTCH** in cells resulted in lower labeling efficiency. The lower labelling efficiency with **AgSiTCH-Halo** is likely due to *trans*-to-*cis* deactivation of the cycloalkane in the intracellular environment. Previously, we have studied the isomerization of *trans*-cyclooctenes under various conditions, and have shown that isomerization of *trans*-cyclooctenes likely occurs via radical mechanism that can be promoted by high thiol concentrations. Consistent with the faster *in cellulo* deactivation of **SiTCH-Halo** relative to **sTCO-Halo**, we observe that **SiTCH 7b** (30 mM) completely isomerizes in 2 h at 22  $^{\circ}$ C in CD<sub>3</sub>OD with mercaptoethanol (30 mM), whereas as control sample without thiol was >98% stable under these conditions.



## ARTICLE

## Journal Name

At  $-17\text{ }^{\circ}\text{C}$ , **7b** isomerizes more slowly in the presence of 30 mM mercaptoethanol, with 47% isomerization after 24 h. By comparison, the *trans*-cyclooctenes d-TCO, s-TCO, and oxo-TCO are much more stable toward thiol promoted isomerization. In  $\text{CD}_3\text{OD}$  with mercaptoethanol (30 mM) at room temperature, s-TCO (30 mM) isomerized only after an 8 hour induction period, with complete conversion to the *cis*-isomer after 4 additional hours.<sup>54</sup> In a similar experiment with d-TCO (30 mM), the induction period was 10 hours. After the induction period, there was 42% isomerization after 4 hours, and 92% isomerization after 14 hours.<sup>56</sup> 5-OxoTCO (25 mM) in the presence of mercaptoethanol (25 mM) showed only 8% isomerization in  $\text{CD}_3\text{OD}$  over a 22 hour period at room temperature.<sup>72</sup>

Together, the labeling experiments in bacteria and HEK 293T cells show that SiTCH derivatives can serve as useful probe molecules in the cellular environment, as the unprecedented speed of the bioorthogonal reactions of SiTCH are much more rapid than competing deactivation pathways. However, the utility of SiTCH derivatives as protein tagging molecules, where extended incubation in the cellular environment takes place prior to bioorthogonal reactivity, appears much more limited in utility plausibly due to alkene isomerization in the cellular environment.

## Conclusions

In conclusion,  $\text{AgNO}_3$  complexes of *trans*-cycloheptene and *trans*-1-sila-4-cycloheptene derivatives have been prepared via a flow photochemical synthesis, using a new low temperature flow photoreactor to enable the synthesis of carbocyclic TCH derivatives.  $\text{TCH}\cdot\text{AgNO}_3$  complexes can be handled for brief periods at rt and stored for weeks in the freezer ( $-18\text{ }^{\circ}\text{C}$ ). Si-TCH $\cdot\text{AgNO}_3$  complexes are especially stable, and can be stored on the bench stable for >1 week at rt, and for months in the freezer. X-ray crystallography was used to characterize a Si-TCH $\cdot\text{AgNO}_3$  complex for the first time. With decomplexation of  $\text{AgNO}_3$  in situ, metal-free TCO and Si-TCH derivatives can engage in a range of cycloaddition reactions as well as dihydroxylation reactions. Computation predicted that Si-TCH would display faster bioorthogonal reactions toward tetrazines than even the most reactive *trans*-cyclooctenes. Metal-free Si-TCH derivatives were shown to display good stability in solution, and to engage in the fastest bioorthogonal reaction reported to date ( $k_2\ 1.14 \times 10^7\ \text{M}^{-1}\text{s}^{-1}$  in 9:1  $\text{H}_2\text{O}:\text{MeOH}$ ). Utility in bioorthogonal protein labeling in live cells is described, including labeling of GFP with an unnatural tetrazine-containing amino acid. The reactivity and specificity of the Si-TCH reagents with tetrazines in live mammalian cells was also evaluated using the HaloTag platform. The cell labeling experiments show that Si-TCH derivatives are suitable as highly reactive probe molecules in the cellular environment.

## Conflicts of interest

There are no conflicts to declare

## Acknowledgements

This work was supported by NIH R01EB014354, R01DC014461, and NSF DMR-1506613. Spectra were obtained with instrumentation supported by NIH grants P20GM104316, P30GM110758, S10RR026962, S10OD016267 and NSF grants CHE-0840401, CHE-1229234, and CHE-1048367. We thank Joshua Judkins for preliminary cellular bioorthogonal labeling experiments, Dr. Glenn Yap and Casey Rowland for x-ray crystallography, Shuyu Xu and He Zhang for IR(ATR) characterization and Prof. Dr. Colin Thorpe for insightful discussions.

## Notes and references


‡ Footnotes relating to the main text should appear here. These might include comments relevant to but not central to the matter under discussion, limited experimental and spectral data, and crystallographic data.

- 1 K. Ziegler and H. Wilms, *Liebigs Ann. Chem.*, 1950, **567**, 1.
- 2 A. C. Cope, R. A. Pike and C. F. Spencer, *J. Am. Chem. Soc.*, 1953, **75**, 3212.
- 3 A. C. Cope, C. R. Ganellin, H. W. Johnson, T. V. Van Auken and H. J. S. Winkler, *J. Am. Chem. Soc.*, 1963, **85**, 3276.
- 4 F. Palacios, I. Perez de Heredia and G. Rubiales, *Tetrahedron Lett.*, 1993, **34**, 4377.
- 5 H. Jendralla, *Tetrahedron*, 1983, **39**, 1359.
- 6 U. Thallhammer, U. Wallfahner and J. Sauer, *Tetrahedron Lett.*, 1990, **31**, 6851.
- 7 K.-J. Shea, J.-S. Kim, *J. Am. Chem. Soc.*, 1992, **114**, 4846.
- 8 W. Weyler Jr., L. R. Byrd, M. C. Caserio and H. W. Moore, *J. Am. Chem. Soc.*, 1972, **94**, 1027.
- 9 H. Jendralla and K. Laumen, *Chem. Ber.*, 1983, **116**, 2136.
- 10 H. Jendralla, *Synthesis*, 1983, 111.
- 11 E. M. Sletten and C. R. Bertozzi, *Angew. Chem. Int. Ed.*, 2009, **48**, 6974.
- 12 D. M. Patterson, L. A. Nazarova and J. A. Prescher, *ACS Chem. Biol.*, 2014, **9**, 592.
- 13 K. Lang and J. W. Chin, *ACS Chem. Biol.*, 2014, **9**, 16.
- 14 C. S. McKay and M. G. Finn, *Chem. Biol.*, 2014, **21**, 1075.
- 15 C. P. Ramil and Q. Lin, *Chem. Commun.*, 2013, **49**, 11007.
- 16 R. Rossin and M. S. Robillard, *Curr. Opin. Chem. Biol.*, 2014, **21**, 161.
- 17 J.-P. Meyer, P. Adumeau, J. S. Lewis and B. M. Zeglis, *Bioconjug. Chem.*, 2016, **27**, 2791.
- 18 M. Vrabel and T. Carell, *Cycloadditions in Bioorthogonal Chemistry*, 2016.
- 19 I. Nikic and E. A. Lemke, *Curr. Opin. Chem. Biol.*, 2015, **28**, 164.
- 20 R. Selvaraj and J. M. Fox, *Curr. Opin. Chem. Biol.*, 2013, **17**, 753.
- 21 A. Darko, S. Wallace, O. Dmitrenko, M. M. Machovina, R. A. Mehl, J. W. Chin and J. M. Fox, *Chem. Sci.*, 2014, **5**, 3770.
- 22 H. Wu and N. K. Devaraj, *Top. Curr. Chem.*, 2015, **374**, 3.
- 23 N. K. Devaraj and R. Weissleder, *Acc. Chem. Res.*, 2011, **44**, 816.
- 24 E. J. Corey, F. A. Carey and R. A. E. Winter, *J. Am. Chem. Soc.*, 1965, **87**, 934.
- 25 J. A. Marshall, *Acc. Chem. Res.*, 1969, **2**, 33.




- 26 P. J. Kropp, *Mol Photochem*, 1978, **9**, 39.
- 27 J. Moran, P. H. Cebrowski and A. M. Beauchemin, *J. Org. Chem.*, 2008, **73**, 1004.
- 28 Y. Inoue, T. Ueoka, T. Kuroda and T. Hakushi, *J. Chem. Soc. Chem. Commun.*, 1981, 1031.
- 29 R. Hoffmann and Y. Inoue, *J. Am. Chem. Soc.*, 1999, **121**, 10702.
- 30 M. E. Squillacote, J. DeFellipis and Q. Shu, *J. Am. Chem. Soc.*, 2005, **127**, 15983.
- 31 G. M. Wallraff and J. Michl, *J. Org. Chem.*, 1986, **51**, 1794.
- 32 (a) R. Bonneau, P. Forner de Violet and J. Jousset-Dubien, *Nouv. J. Chim.*, 1977, **1**, 31. (b) R. Bonneau, J. Jousset-Dubien, J. Yarwood and J. Pereyre, *Tetrahedron Lett.*, 1977, 235.
- 33 X.-N. Wang, E. H. Krenske, R. C. Johnston, K. N. Houk and R. P. Hsung, *J. Am. Chem. Soc.*, 2015, **137**, 5596.
- 34 E. J. Corey, M. Tada, R. LaMahieu and L. Libit, *J. Am. Chem. Soc.*, 1965, **87**, 2051.
- 35 P. E. Eaton and K. Lin, *J. Am. Chem. Soc.*, 1965, **87**, 2052.
- 36 J. Moran, P. Dornan and A. M. Beauchemin, *Org. Lett.*, 2007, **9**, 3893.
- 37 H. Dorr and V. H. Rawal, *J. Am. Chem. Soc.*, 1999, **121**, 10229.
- 38 H. M. L. Davies, Ø. Loe and D. G. Stafford, *Org. Lett.*, 2005, **7**, 5561.
- 39 R. G. Salomon, K. Folting, W. E. Streib and J. K. Kochi, *J. Am. Chem. Soc.*, 1974, **96**, 1145.
- 40 T. Spee and A. Mackor, *J. Am. Chem. Soc.*, 1981, **103**, 6901.
- 41 J. T. M. Evers and A. Mackor, *Recl. des Trav. Chim. des Pays-Bas*, 1979, **98**, 423.
- 42 H. Nishiyama, T. Naitoh, Y. Motoyama and K. Aoki, *Chem. - Eur. J.*, 1999, **5**, 3509.
- 43 H. Jendralla, *Angew. Chem. Int. Ed.*, 1980, **19**, 1032.
- 44 H. Jendralla, *Chem. Ber.*, 1980, **113**, 3557.
- 45 T. Shimizu, K. Shimizu and W. Ando, *J. Am. Chem. Soc.*, 1991, **113**, 354.
- 46 A. W. Krebs, K.-I. Pforr, W. Raffay, B. Thölke, W. A. König, I. Hardt and R. Boese, *Angew. Chem. Int. Ed.*, 1997, **36**, 159.
- 47 A. W. Krebs, B. Thölke, K.-I. Pforr, W. A. König, K. Scharwächter, S. Grimme, F. Vögtle, A. Sobanski, J. Schramm and J. Holmes, *Tetrahedron: Asymmetry*, 1999, **10**, 3483.
- 48 J. Santucci, J. R. Sanzone and K. A. Woerpel, *Eur. J. Org. Chem.*, 2016, **2016**, 2933.
- 49 M. A. Greene, M. Prévost, J. Tolopilo and K. A. Woerpel, *J. Am. Chem. Soc.*, 2012, **134**, 12482.
- 50 B. Hurlocker, C. Hu and K. A. Woerpel, *Angew. Chem. Int. Ed.*, 2015, **54**, 4295.
- 51 (a) J. R. Sanzone, C. T. Hu and K. A. Woerpel, *J. Am. Chem. Soc.*, 2017, **139**, 8404. (b) J. R. Sanzone and K. A. Woerpel, *Synlett.*, 2017, DOI: 10.1055/s-0036-1588451
- 52 K. Tomooka, S. Miyasaka, S. Motomura and K. Igawa, *Chem. Eur. J.*, 2014, **20**, 7598.
- 53 M. Royzen, G. P. A. Yap and J. M. Fox, *J. Am. Chem. Soc.*, 2008, **130**, 3760.
- 54 M. T. Taylor, M. L. Blackman, O. Dmitrenko and J. M. Fox, *J. Am. Chem. Soc.*, 2011, **133**, 9646.
- 55 M. Royzen, M. T. Taylor, A. DeAngelis and J. M. Fox, *Chem. Sci.*, 2011, **2**, 2162.
- 56 H. E. Murrey, J. C. Judkins, C. W. Am Ende, T. E. Ballard, Y. Fang, K. Riccardi, L. Di, E. R. Guilmette, J. W. Schwartz, J. M. Fox and D. S. Johnson, *J. Am. Chem. Soc.*, 2015, **137**, 11461.
- 57 B. D. A. Hook, W. Dohle, P. R. Hirst, M. Pickworth, M. B. Berry and K. I. Booker-Milburn, *J. Org. Chem.*, 2005, **70**, 7558.
- 58 G. V. Los, L. P. Encell, M. G. McDougall, D. D. Hartzell, N. Karassina, C. Zimprich, M. G. Wood, R. Learish, R. F. Ohana, M. Urh, D. Simpson, J. Mendez, K. Zimmerman, P. Otto, G. Vidugiris, J. Zhu, A. Darzins, D. H. Klaubert, R. F. Bulleit and K. V. Wood, *ACS Chem. Biol.*, 2008, **3**, 373.
- 59 D. B. Janssen, *Curr. Opin. Chem. Biol.*, 2004, **8**, 150.
- 60 I. Rencken, J. C. A. Boeyens and S. W. Orchard, *J. Cryst. Spec. Res.*, 1988, **18**, 293.
- 61 M. Tsutsui, M. N. Levy and A. Nakamura, *Introduction to Metal  $\pi$ -complex chemistry*; Springer: New York, 1970; p 40.
- 62 A. A. Vasil'ev, N. A. Donskaya, G. V. Cherkaev and N. M. Yur'eva, *J. Org. Chem. (USSR)*, 1991, **27**, 273.
- 63 J. H. Smitrovich and K. A. Woerpel, *J. Org. Chem.*, 1996, **61**, 6044.
- 64 R. Rossin, S. M. J. Van Duijnhoven, W. Ten Hoeve, H. M. Janssen, L. H. J. Kleijn, F. J. M. Hoebe, R. M. Versteegen and M. S. Robillard, *Bioconjug. Chem.*, 2016, **27**, 1697.
- 65 R. M. Versteegen, R. Rossin, W. Ten Hoeve, H. M. Janssen and M. S. Robillard, *Angew. Chem. Int. Ed.*, 2013, **52**, 14112.
- 66 J. Li, S. Jia and P. R. Chen, *Nat. Chem. Biol.*, 2014, **10**, 1003.
- 67 X. Fan, Y. Ge, F. Lin, Y. Yang, G. Zhang, W. S. C. Ngai, Z. Lin, S. Zheng, J. Wang, J. Zhao, J. Li and P. R. Chen, *Angew. Chem. Int. Ed.*, 2016, **55**, 14046.
- 68 (a) G. de Almeida, E. M. Sletten, H. Nakamura, K. K. Palaniappan and C. R. Bertozzi, *Angew. Chem. Int. Ed.*, 2012, **51**, 2443. (b) G. de Almeida, L. C. Townsend and C. R. Bertozzi, *Org. Lett.*, 2013, **15**, 3038. (c) M. Martinek, L. Filipova, J. Galeta, L. Ludvikova and P. Klan, *Org. Lett.*, 2016, **18**, 4892. For earlier work on heteroatom containing cycloheptynes: (d) H. Kimling and A. Krebs *Liebigs Ann. Chem.*, 1974, 2074. (e) A. Krebs and H. Kimling *Tetrahedron Lett.*, 1970, **11**, 761.
- 69 M. L. Blackman, M. Royzen and J. M. Fox, *J. Am. Chem. Soc.*, 2008, **130**, 13518.
- 70 N. K. Devaraj, S. Hilderbrand, R. Upadhyay, R. Mazitschek and R. Weissleder, *Angew. Chem. Int. Ed.*, 2010, **49**, 2869.
- 71 R. J. Blizzard, D. R. Backus, W. Brown, C. G. Bazewicz; Y. Li; R. A. Mehl, *J. Am. Chem. Soc.*, 2015, **137**, 10044.
- 72 W. D. Lambert, S. L. Scinto, O. Dmitrenko, S. J. Boyd, R. Magboo, R. A. Mehl, J. W. Chin, J. M. Fox and S. Wallace, *Org. Biomol. Chem.*, 2017, **15**, 6640.





stable metal complexes



✓ ~~Chemical Synthesis~~ **flow synthesis**

Page 12 of 12

✓ x-ray structure of silacycle

✓ applications in synthesis

✓ bioorthogonal reactions  $>10^7 \text{ M}^{-1}\text{s}^{-1}$

✓ protein labeling in live cells