View Article Online View Journal



Organic & Biomolecular Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: S. K. Woo and D. Goo, *Org. Biomol. Chem.*, 2015, DOI: 10.1039/C5OB01952D.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/obc

Organic & Biomolecular Chemistry

Deuk-young Goo and Sang Kook Woo*

ARTICLE



One-Pot Synthesis of Carbazoles via Tandem C-C Cross-Coupling and Reductive Amination

Received 00th January 20xx, Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

We have developed a highly efficient synthetic route to carbazoles that employs sequential C-C/C-N bond formation via Suzuki cross-coupling and Cadogan cyclization using commercially available or easily preparable starting materials. The developed method is compatible with electron neutral, rich or deficient substrates. The synthetic utility of this method was demonstrated by the concise syntheses of four natural products (glycozoline, glycozolicine, glycozolidine and clausenalene).

Introduction

The Carbazole is an ubiquitous structural motif in a large number of biologically active natural products and pharmaceuticals.¹ In addition, carbazole derivatives have been widely used as functional organic materials.² Thus, development of efficient synthetic routes to structurally diverse carbazoles represents an important objective in organic synthesis. Many methods have been developed for the preparation of carbazole.³ The most widely used approaches for synthesis of the carbazole framework are C-N or C-C bond forming cyclizations from substituted biaryls or N,N-biaryl amines. Representative C-N bond forming reactions for synthesis of carbazoles are summarized in Scheme 1 (a): (A) transition metal free⁴ or palladium catalyzed Buchwald-Hartwig⁵ amination from 2-amino-2'-halo-biphenyls; (B) dehydrogenative C-H/N-H coupling of 2-amino-biphenyls⁶; (C) iron-mediated oxidative amination of iron $complex^{7}$; (D) transition metal mediated double amination of 2,2'dihalobiphenyls or cyclic diphenyleneiodoniums⁸; (E) transition metal free⁹ or rhodium catalyzed¹⁰ nitrene insertion of 2-azidobiphenyls; (F) reductive amination of 2-nitro-biphenyls¹¹. Similarly, dehydrogenative C-C bond formation of N,N-biaryl amines leads to carbazoles (Scheme (b)).¹² Although these methods proceed in good yields and are tolerant of most organic functional groups. They require the preparation of complex starting materials, such as substituted biaryls or N,Nbiaryl amines for the synthesis of diverse carbazoles. Therefore, we investigated the development of a one-pot C-C and C-N bond coupling reaction that utilizes commercially available or

easily preparable starting materials.

Previous work

(a) C-N bond formation for the synthesis of carbazoles





(b) C-C bond formation for the synthesis of carbazoles



(c) One-pot synthesis of carbazoles



Scheme 1. Various synthetic approaches to carbazoles.

Published on 28 October 2015. Downloaded by UNIVERSITY OF OTAGO on 02/11/2015 10:29:46.

Department of Chemistry, University of Ulsan, 93 Daehak-Ro, Nam-Gu, Ulsan 44610, Korea. E-mail: <u>woosk@ulsan.ac.kr;</u> Fax: +82-52-259-2348; Tel: +82-52-259-2338

⁺Electronic Supplementary Information (ESI) available: Details of experimental procedures and spectroscopic data for all new com-pounds (¹H NMR, ¹³C NMR, IR, MS), including images of NMR spectra. See DOI: 10.1039/x0xx00000x

Published on 28 October 2015. Downloaded by UNIVERSITY OF OTAGO on 02/11/2015 10:29:46.

Organic & Biomolecular Chemistry





Entry 1, X Catalyst (mol%) Reductant T (°C) time (h) Y (%) ^b 1 1a, I Pd(OAc) (2) PPh ₃ 150 16 24 2 1b, Br Pd(OAc) (2) PPh ₃ 150 16 56 3 1c, CI Pd(OAc) (2) PPh ₃ 150 16 31 4 1b, Br Pd(OAc) (2) PEt ₃ 150 16 12 5 1b, Br Pd(OAc) (2) PC(Et ₃) 150 16 21 6 1b, Br Pd ₂ (dba) ₃ (2) PPh ₃ 150 16 21 6 1b, Br Pd ₂ (dba) ₃ (2) PPh ₃ 150 16 21 6 1b, Br Pd(OAc) (1) PPh ₃ 150 16 29 8 1b, Br Pd(OAc) (2) PPh ₃ 150 16 13 10 1b, Br Pd(OAc) (2) PPh ₃ 180 16 72 11 1b, Br Pd								
1 1a, I Pd(OAc) (2) PPh ₃ 150 16 24 2 1b, Br Pd(OAc) (2) PPh ₃ 150 16 56 3 1c, CI Pd(OAc) (2) PPh ₃ 150 16 31 4 1b, Br Pd(OAc) (2) PEt ₃ 150 16 12 5 1b, Br Pd(OAc) (2) P(OEt) ₃ 150 16 21 6 1b, Br Pd(OAc) (2) P(OEt) ₃ 150 16 41 7 1b, Br Pd(DAc) (1) PPh ₃ 150 16 29 8 1b, Br Pd(OAc) (1) PPh ₃ 150 16 17 9 1b, Br Pd(OAc) (2) PPh ₃ 150 16 13 10 1b, Br Pd(OAc) (2) PPh ₃ 180 16 72 11 1b, Br Pd(OAc) (2) PPh ₃ 180 24 78 12 ^c 1b, Br Pd(OAc) (2) PPh ₃ Reflux 24 66 13 ^d 1b, Br Pd(OAc) (2)	Entry	1, X	Catalyst (mol%)	Reductant	T (°C)	time (h)	Y (%) ^b	
2 1b, Br Pd(OAc) (2) PPh ₃ 150 16 56 3 1c, Cl Pd(OAc) (2) PPh ₃ 150 16 31 4 1b, Br Pd(OAc) (2) PEt ₃ 150 16 12 5 1b, Br Pd(OAc) (2) P(OEt) ₃ 150 16 21 6 1b, Br Pd(OAc) (2) P(OEt) ₃ 150 16 41 7 1b, Br Pd(OAc) (1) PPh ₃ 150 16 29 8 1b, Br Pd(OAc) (1) PPh ₃ 150 16 13 9 1b, Br Pd(OAc) (2) PPh ₃ 150 16 13 10 1b, Br Pd(OAc) (2) PPh ₃ 180 16 72 11 1b, Br Pd(OAc) (2) PPh ₃ 180 24 78 12 ^c 1b, Br Pd(OAc) (2) PPh ₃ Reflux 24 66 \Rightarrow 13 ^d 1b, Br Pd(OAc) (2) PPh ₃ 180 24 86	1	1a, I	Pd(OAc) (2)	PPh_3	150	16	24	
3 1c, Cl Pd(OAc) (2) PPh ₃ 150 16 31 4 1b, Br Pd(OAc) (2) PEt ₃ 150 16 12 5 1b, Br Pd(OAc) (2) P(OEt) ₃ 150 16 21 6 1b, Br Pd ₂ (dba) ₃ (2) PPh ₃ 150 16 41 7 1b, Br Pd(PPh ₃) ₄ (2) PPh ₃ 150 16 29 8 1b, Br Pd(OAc) (1) PPh ₃ 150 16 17 9 1b, Br Pd(OAc) (2) PPh ₃ 150 16 13 10 1b, Br Pd(OAc) (2) PPh ₃ 180 16 72 11 1b, Br Pd(OAc) (2) PPh ₃ 180 24 78 12 ^c 1b, Br Pd(OAc) (2) PPh ₃ Reflux 24 66 \Rightarrow 13 ^d 1b, Br Pd(OAc) (2) PPh ₃ 180 24 86	2	1b, Br	Pd(OAc) (2)	PPh ₃	150	16	56	
4 1b, Br Pd(OAc) (2) PEt ₃ 150 16 12 5 1b, Br Pd(OAc) (2) P(OEt) ₃ 150 16 21 6 1b, Br Pd ₂ (dba) ₃ (2) PPh ₃ 150 16 41 7 1b, Br Pd(PPh ₃) ₄ (2) PPh ₃ 150 16 29 8 1b, Br Pd(OAc) (1) PPh ₃ 150 16 17 9 1b, Br Pd(OAc) (2) PPh ₃ 150 16 13 10 1b, Br Pd(OAc) (2) PPh ₃ 180 16 72 11 1b, Br Pd(OAc) (2) PPh ₃ 180 24 78 12 ^c 1b, Br Pd(OAc) (2) PPh ₃ Reflux 24 66 ⇒ 13 ^d 1b, Br Pd(OAc) (2) PPh ₃ 180 24 86	3	1c, Cl	Pd(OAc) (2)	PPh ₃	150	16	31	
5 1b, Br Pd(OAc) (2) P(OEt)_3 150 16 21 6 1b, Br Pd ₂ (dba) ₃ (2) PPh ₃ 150 16 41 7 1b, Br Pd(PPh ₃) ₄ (2) PPh ₃ 150 16 29 8 1b, Br Pd(OAc) (1) PPh ₃ 150 16 17 9 1b, Br Pd(OAc) (2) PPh ₃ 150 16 13 10 1b, Br Pd(OAc) (2) PPh ₃ 180 16 72 11 1b, Br Pd(OAc) (2) PPh ₃ 180 24 78 12 ^c 1b, Br Pd(OAc) (2) PPh ₃ Reflux 24 66 ⇒ 13 ^d 1b, Br Pd(OAc) (2) PPh ₃ 180 24 86	4	1b, Br	Pd(OAc) (2)	PEt ₃	150	16	12	
6 1b, Br Pd ₂ (dba) ₃ (2) PPh ₃ 150 16 41 7 1b, Br Pd(PPh ₃) ₄ (2) PPh ₃ 150 16 29 8 1b, Br Pd(OAc) (1) PPh ₃ 150 16 17 9 1b, Br Pd(OAc) (2) PPh ₃ 150 16 13 10 1b, Br Pd(OAc) (2) PPh ₃ 180 16 72 11 1b, Br Pd(OAc) (2) PPh ₃ 180 24 78 12 ^c 1b, Br Pd(OAc) (2) PPh ₃ Reflux 24 66 ⇒ 13 ^d 1b, Br Pd(OAc) (2) PPh ₃ 180 24 86	5	1b, Br	Pd(OAc) (2)	P(OEt) ₃	150	16	21	
71b, BrPd(PPh_3)_4 (2)PPh_3150162981b, BrPd(OAc) (1)PPh_3150161791b, BrPd(OAc) (4)PPh_31501613101b, BrPd(OAc) (2)PPh_31801672111b, BrPd(OAc) (2)PPh_31802478 12^c 1b, BrPd(OAc) (2)PPh_3Reflux2466 \Rightarrow 13^d1b, BrPd(OAc) (2)PPh_31802486	6	1b, Br	Pd ₂ (dba) ₃ (2)	PPh ₃	150	16	41	
8 1b, Br Pd(OAc) (1) PPh ₃ 150 16 17 9 1b, Br Pd(OAc) (4) PPh ₃ 150 16 13 10 1b, Br Pd(OAc) (2) PPh ₃ 180 16 72 11 1b, Br Pd(OAc) (2) PPh ₃ 180 24 78 12 ^c 1b, Br Pd(OAc) (2) PPh ₃ Reflux 24 66 ⇒ 13 ^d 1b, Br Pd(OAc) (2) PPh ₃ 180 24 86	7	1b, Br	Pd(PPh ₃) ₄ (2)	PPh ₃	150	16	29	
9 1b, Br Pd(OAc) (4) PPh3 150 16 13 10 1b, Br Pd(OAc) (2) PPh3 180 16 72 11 1b, Br Pd(OAc) (2) PPh3 180 24 78 12^c 1b, Br Pd(OAc) (2) PPh3 Reflux 24 66 ⇒ 13 ^d 1b, Br Pd(OAc) (2) PPh3 180 24 86	8	1b, Br	Pd(OAc) (1)	PPh ₃	150	16	17	
10 1b, Br Pd(OAc) (2) PPh ₃ 180 16 72 11 1b, Br Pd(OAc) (2) PPh ₃ 180 24 78 12 ^c 1b, Br Pd(OAc) (2) PPh ₃ Reflux 24 66 \Rightarrow 13 ^d 1b, Br Pd(OAc) (2) PPh ₃ 180 24 86	9	1b, Br	Pd(OAc) (4)	PPh ₃	150	16	13	
11 1b, Br Pd(OAc) (2) PPh ₃ 180 24 78 12^c 1b, Br Pd(OAc) (2) PPh ₃ Reflux 24 66 \Rightarrow 13 ^d 1b, Br Pd(OAc) (2) PPh ₃ 180 24 86	10	1b, Br	Pd(OAc) (2)	PPh ₃	180	16	72	
12^{c} 1b, Br Pd(OAc) (2) PPh ₃ Reflux 24 66 \Rightarrow 13 ^d 1b, Br Pd(OAc) (2) PPh ₃ 180 24 86	11	1b, Br	Pd(OAc) (2)	PPh ₃	180	24	78	
\Rightarrow 13 ^d 1b, Br Pd(OAc) (2) PPh $_3$ 180 24 86	12 ^c	1b, Br	Pd(OAc) (2)	PPh ₃	Reflux	24	66	
	≎ 13 ^d	1b, Br	Pd(OAc) (2)	PPh ₃	180	24	86	

^aReaction conditions: **1** (0.5 mmol), **2a** (0.6 mmol), catalyst (quantity noted), reductant (250 mol%), K_2CO_3 (200 mol%), *o*-DCB (1.0 mL) under air in pressure tubes. ^bIsolated yield by flash column chromatography. ^cUsing reflux condenser. ^d**2a** (130 mol%).

More recently, one-pot syntheses of carbazoles have been developed (Scheme 1 (c)).¹³ Bedford^{13a} and Ackermann^{13b, 13d} reported a palladium catalyzed domino Buchwald-Hartwig amination and C-H activation of 2-chloroanilines with aryl bromides or anilines with dichloroarenes. Jean, Jr^{13c} reported the Pd-catalyzed tandem Suzuki cross-coupling and S_NAr reaction of aniline-derived boronic esters with 2-fluoro-3-halobenzenes. Studer^{13f} reported a transition metal-free cascade reaction of nitrosoarenes with *in situ* generated arynes.

We envision a one-pot reaction sequence of C-C crosscoupling and C-N bond forming cyclization to develop a stepeconomic method to prepare carbazoles. We considered the Suzuki reaction¹⁴ and Cadogan cyclization¹⁵ for the one-pot reaction. Both reactions are widely used methods for C-C cross-coupling or reductive amination and use similar phosphine reagents for ligand or reductant^{15d}. Herein, we present a highly efficient synthetic method to carbazoles that consists of palladium-catalyzed tandem C-C/C-N bond formation via Suzuki reaction and reductive Cadogan cyclization (Scheme 1).

Results and discussion

Initially, we explored the domino reaction of 2iodonitrobenzene **1a** with phenylboronic acid **2a** in the presence of $Pd(OAc)_2$ and triphenylphosphine in *o*-DCB at 150 °C, which delivered the carbazole **3a** and 2-nitrobiphenyl in 24% and 20% yield (Table 1, entry 1). Based on this result, other 2halonitrobenzenes were assayed, and it was found that 2bromonitrobenzene **1b** was the most efficient (Table 1, entry 2).



^aAs given in Table 1, entry 13. ^b48h. ^cPd(OAc) (5 mol%), 48h.; yields of isolated material. See ESI[†] for details.

Thus chosen for further studies. When other reductants such as PEt₃ and P(OEt)₃^{15e} were used, the yield of the reaction did not improve (Table 1, entries 4 and 5). Moreover, other catalysts such as $Pd_2(dba)_3$ and $Pd(PPh_3)_4$ were not effective in this transformation (Table 1, entries 6 and 7). Reductant and catalyst screening indicated that $Pd(OAc)_2$ and PPh_3 were the best choices. To improve the yield, the catalyst loading, as well as reaction temperature and time were surveyed. Increasing and decreasing the catalyst loading were not efficient (Table 1, entries 8 and 9). However, the yield did increase with a higher temperature and longer reaction time (Table 1, entries 10 and 11). Coupling reaction was not efficient in atmospheric pressure (Table 1, entry 12). Using 130 mol% phenylboronic acid **2a**, gave the best result (Table 1, entry 13).

We next examined the substrate scope with respect to the substituted arylboronic acid (Table 2). Under the optimized conditions, *o*-bromonitrobenzene **1b** was coupled to a diverse range of arylboronic acids **2** to form carbazoles **3a-3o** in moderate to excellent yields.



Scheme 2. Further functionalization of 3-chlorocarbazole (3i). ^aSee ESI⁺ for details.

Notably, electron neutral and electron rich arylboronic acids possessing alkyl, aryl, and alkoxy substituents were tolerated in the one-pot reaction (3a-3h, 3m-o), as o-, m- and psubstituted arylboronic acids and 3,4-disubstitued arylboronic acids afforded the corresponding products (3a-3h, 3m-o) in good to high yields. Aryl boronic acids possessing bulky groups at the para position such as t-butyl and phenyl were also converted to carbazoles in excellent yields. As expected, coupling of *m*-substituted arylboronic acids with obromonitrobenzene 1b gave a mixture of corresponding 1- and 3-substituted carbazoles as regioisomers. In these experiments, 3-substituted carbazoles were obtained as major regioisomer except with 3-tolylboronic acid. In particular, 3.4-(methylenedioxy)phenylboronic acid gave a 5.9:1 mixture of 3n and 3n', demonstrating moderate regioselectivity, while 3tolylboronic acid and 4-methoxy-3-methylphenylboronic acid gave mixtures with low regioselectivity (3m¹⁶/m', 3o¹⁷/o').

In addition, we explored the scope of electron deficient arylboronic acids possessing fluorine, chlorine, ketone, and ester moieties and found they also gave the desired products (**3i-3i**). Slightly lower yields were observed for electron deficient arylboronic acids except for 4-fluorophenylboronic acid.

Table 3. Substrate scope of various o-bromonitrobenzenes^a



^aAs given in Table 1, entry 13. ^b48h. ^cPd(OAc) (5 mol%), 48h.; yields of isolated material. See ESI[†] for details.

In particular, 4-fluoro and 4-chloroboronic _{Viacidscle} were tolerated under these coupling conditions? Halogen Containing carbazoles are useful for the introduction of additional functionalization. Thus, we examined the further functionalization of 3-chlorocarbazole **3i** (Scheme 2). Under modified Suzuki cross-coupling conditions¹⁸ for unprotected *N*heterocycles, 3-chlorocarbazole **3i** was coupled to a diverse range of arylboronic acids to form functionalized carbazoles (**3e, 3ia-3id**) in good to excellent yields.

To demonstrate the efficiency of this method, we conducted the gram scale synthesis of carbazole **3a**. Coupling reaction of 2-bromonitrobenzene **1b** (1 g, 5 mmol) with phenylboronic acid **2a** (750 mg, 6.5 mmol) afforded the carbazole **3a** in 75% yields.

Next, we turned our attention to the domino reaction of phenylboronic acid with a broad range of substituted *o*-bromonitrobenzenes (Table 3). Under the optimized conditions, the one-pot reaction of substituted *o*-bromonitrobenzenes **1** with phenylboronic acid **2a** were converted to the corresponding carbazoles in good yields, and similar results with respect to substrate scope of various aryl boronic acids were observed (Table 3, **3g**, **3i**, **3j**, **3m** and **3n**). These approaches constitute an alternative route that avoids the regioselectivity issue in the synthesis of 1- or 3-substituted carbazoles (**3m** – **3q**).

Table 4. Substrate scope of substituted o bromonitrobenzenes with aryl boronic acids^a



^aAs given in Table 1, entry 13. ^b48h.; yields of isolated material. See ESI¹ for details.

This journal is © The Royal Society of Chemistry 2015

The scope and limitations of the domino reaction with respect to various substituted o-bromonitrobenzenes and arylboronic acids were next investigated under the optimized conditions(Table 4). Coupling reactions of 1-bromo-4-methoxy-2-nitrobenzene with 3-tolylboronic acid and 4fluorophenylboronic acid afforded the corresponding carbazoles 3s and 3t in 71% and 66% yields. Likewise, obromonitrobenzenes possessing methyl and fluoro substituents were coupled to arylboronic acids to form carbazoles 3u and 3v in 70% and 53% yield. This tandem reaction was applied to the syntheses of three bioactive carbazole alkaloids, glycozoline **3w**¹⁹, glycozolicine **3w'**²⁰, glycozolidine $3x^{13a, 21}$ and clausenalene $3y^{22}$, in good yields and moderate regioselectivity.

Conclusions

In conclusion, we have developed highly efficient synthetic methods for carbazoles that are palladium-catalyzed sequential C-C/C-N bond formations via Suzuki cross-coupling and reductive Cadogan cyclization. Electron neutral, rich and deficient substrates were tolerated in this method. The synthetic utility of this method was demonstrated by the concise syntheses of four naturally occurring bioactive carbazole alkaloids, glycozoline **3w**, glycozolicine **3w'**, glycozolidine **3x** and clausenalene **3y** from commercially available simple starting materials. Future studies will focus on further applications of this one-pot strategy to the syntheses of complex natural products.

Experimental

General Information

All reactions were run under an atmosphere of air under anhydrous conditions otherwise indicated. unless Dichloromethane $(CH_2Cl_2),$ tetrahydrofuran (THF), dimethylformamide (DMF) and toluene (PhMe) were obtained from Pure-Solv MD-5 Solvent Purification System (Innovative Technology). Pressure tubes (13 x 100 mm, PYREXPLUS) were dried in oven for overnight and cooled under a stream of nitrogen prior to use. All commercial reagents were used directly without further purification. The progress of reaction was checked on TLC plates (Merck 5554 Kiesel gel 60 F₂₅₄). Column chromatography was performed on silica gel (Merck 9385 Kiesel gel 60) using hexanes-EtOAc (v/v) or hexanesacetone (v/v). Infrared spectra were recorded on a Varian 2000 FT-IR. High-resolution mass spectra (EI) were obtained on a Jeol JMS700 HRMS at the Korea Basic Science Center(KBSI), Daegu, Korea and Low-resolution mass spectra (EI) Varian 450-GC/Varian 220-MS and are reported as m/z (relative intensity). Accurate masses are reported for the molecular ion ([M+H]⁺ or [M+Na]⁺). Nuclear magnetic resonance spectra (¹H NMR and ¹³C NMR) were recorded with a Bruker (300 MHz) spectrometer. Chemical shift values were recorded as parts per million relative to tetramethylsilane as an internal standard unless otherwise indicated, and coupling constants in Hertz. The following abbreviations are used: m (multiplet) ins (singlet), d (doublet), t (triplet), q (quartet)). ଏସି ଏସି ଅଧିକାର୍ଯ୍ୟ କେମ୍ବାର୍ ସେମ୍ବର ସେମ୍ବର ସେମ୍ବର କେମ୍ବର କେମ୍ବ doublets), etc.

Organic & Biomolecular Chemistry

General Procedure for Carbazole Synthesis. To a re-sealable pressure tube (13 x 100 mm) equipped with magnetic stir bar were added *o*-bromonitrobenzene **1** (0.5 mmol, 100 mol%), aryl boronic acid **2** (0.65 mmol, 130 mol%), Pd(OAc) (0.01 mmol, 2 mol%), PPh₃ (1.25 mmol, 250 mol %), K₂CO₃ (1 mmol, 200 mol%) and *o*-DCB (1 mL, 0.5 M concentration with respect to *o*-bromonitrobenzene **1**. The mixture was heated at 180 °C (oil bath temperature) for 24-48 hr, at which point the reaction mixture was allowed to cool to ambient temperature. The reaction mixture was filtered through a pad of celite and the resulting liquor was concentrated *in vacuo* and purified by flash column chromatography (SiO₂) under the conditions noted to furnish the corresponding product.

9H-carbazole (3a)^{6e}. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3a** as a white solid (72 mg, 86% yield); ¹H NMR (300 MHz, DMSO) δ 11.33 (s, 1H), 8.12 (d, *J* = 7.8 Hz, 3H), 7.56 (d, *J* = 8.1 Hz, 2H), 7.45 – 7.39 (m, 3H), 7.21 – 7.16 (m, 3H); ¹³C NMR (75 MHz, DMSO) δ 139.79, 125.54, 122.48, 120.18, 118.53, 110.98; FTIR (neat): υ 3379, 2946, 1450, 1033 cm⁻¹; LRMS *m/z* (EI) 166.8 (M⁺-1), 139.0, 83.4, 62.8.

2-Methyl-9H-carbazole (**3b**)^{23a}. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3b** as a white solid (76 mg 84% yield); ¹H NMR (300 MHz, CDCl₃) δ 8.03 (d, *J* = 7.8 Hz, 1H), 7.95 (d, *J* = 7.9 Hz, 2H), 7.42 – 7.35 (m, 2H), 7.24 – 7.18 (m, 2H), 7.07 (d, *J* = 8.0 Hz, 1H), 2.53 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 140.09, 139.59, 136.15, 125.41, 123.58, 121.19, 121.12, 120.14, 120.13, 119.46, 110.86, 110.60, 22.21; FTIR (neat) u 3394, 2920, 1652, 1530, 1365 cm⁻¹; LRMS *m/z* (EI) 179.8 (M⁺-1), 151.8, 76.8, 62.8.

4-Methyl-9H-carbazole (3c)¹⁰. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3c** as a white solid (79 mg 87% yield); ¹H NMR (300 MHz, CDCl₃): δ 8.19 (d, J = 7.9 Hz, 1H), 8.09 (br s, 1H), 7.50 – 7.39 (m, 2H), 7.37 – 7.31 (m, 1H), 7.32 – 7.26 (m, 2H), 7.02 (dd, J = 7.3, 0.8 Hz, 1H), 2.89 (s, 3H); ¹³C NMR (75 MHz, CDCl₃): δ 139.54, 139.51, 133.49, 125.77, 125.29, 123.93, 122.68, 121.90, 120.99, 119.44, 110.49, 108.26, 20.92; FTIR (neat) u 3396, 3200, 2924, 1653, 1521, 1267, 1048 cm⁻¹; LRMS *m/z* (EI) 181.3 (M⁺), 182.3, 180.4, 177.5, 155.3, 125.8, 93.7.

2-(Tert-butyl)-9H-carbazole (3d)^{11d}. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3d** as a white solid (94 mg 84% yield); ¹H NMR (300 MHz, CDCl₃) δ 8.05 (d, J = 7.0 Hz, 1H), 8.00 (d, J = 7.5 Hz, 1H), 7.94 (s, 1H), 7.44 (d, J = 1.6 Hz, 1H), 7.41 – 7.35 (m, 2H), 7.32 (dd, J = 8.3, 1.4 Hz, 1H), 7.25 – 7.20 (m, 1H), 1.44 (s, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 149.72, 139.89, 139.82, 125.40, 123.49, 121.03, 120.21, 119.90, 119.41, 117.65, 110.57, 107.25, 35.25, 31.90; FTIR (neat) \cup 3194, 2916, 1650, 1525, 1267 cm⁻¹; LRMS m/z (EI) 222.7 (M⁺-1), 207.8, 180.0, 167.0, 89.9.

2-Phenyl-9H-carbazole $(3e)^{6d}$. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3e** as a white solid (110 mg 90% yield); ¹H NMR (300

Published on 28 October 2015. Downloaded by UNIVERSITY OF OTAGO on 02/11/2015 10:29:46.

Organic & Biomolecular Chemistry

MHz, Acetone-d₆) δ 10.50 (s, 1H), 8.19 (d, J = 8.1 Hz, 1H), 8.14 (d, J = 7.7 Hz, 1H), 7.80 – 7.71 (m, 3H), 7.57 – 7.32 (m, 6H), 7.21 (td, J = 8.2, 4.3 Hz, 1H); ¹³C NMR (75 MHz, DMSO): δ 141.26, 140.43, 140.35, 137.88, 128.99, 127.12, 127.06, 125.70, 122.23, 121.88, 120.68, 120.32, 118.76, 117.89, 111.06, 108.93; FTIR (neat) \cup 3377, 3268, 1692, 1528, 1367, 1227 cm⁻¹; LRMS m/z (EI): 242.8 (M⁺-1), 120.3, 76.7, 50.9.

7H-Benzo[c]carbazole (3f)^{23b}. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3f** as a white solid (91 mg 84% yield); ¹H NMR (300 MHz, CDCl₃) δ 8.79 (d, *J* = 8.3 Hz, 1H), 8.58 (d, *J* = 7.8 Hz, 1H), 8.44 (s, 1H), 8.01 (d, *J* = 8.1 Hz, 1H), 7.87 (d, *J* = 8.8 Hz, 1H), 7.75 – 7.68 (m, 1H), 7.65 (d, *J* = 8.8 Hz, 1H), 7.59 (d, *J* = 7.8 Hz, 1H), 7.52 – 7.43 (m, 2H), 7.43 – 7.36 (m, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 138.52, 137.16, 130.03, 129.32, 127.55, 127.01, 124.46, 124.09, 123.38, 123.13, 122.16, 120.36, 115.54, 112.71, 111.25; FTIR (neat) \cup 3396, 2920, 1876, 1465, 1267, 1120 cm⁻¹; LRMS *m/z* (EI): 217.7 (M⁺+1), 216.8, 188.8, 108.2, 95.3.

2-Methoxy-9H-carbazole (3g)^{16c}. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give 3g as a white solid (76 mg, 77% yield); ¹H NMR (300 MHz, CDCl₃) δ 7.99 (s, 1H), 7.96 (d, J = 3.4 Hz, 1H), 7.94 (d, J = 8.6 Hz, 1H), 7.41 – 7.31 (m, 1H), 7.24 – 7.18 (m, 1H), 6.92 (d, J = 2.1 Hz, 1H), 6.86 (dd, J = 8.5, 2.2 Hz, 1H), 3.91 (s, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 159.21, 140.93, 139.63, 124.69, 123.67, 121.19, 119.70, 119.63, 117.39, 110.44, 108.29, 94.83, 55.77; FTIR (neat) \cup 3395, 1336, 1200, 1030 cm⁻¹; LRMS m/z (EI): 197.7 (M⁺+1), 196.7, 181.8, 153.8, 127.8.

4-Methoxy-9H-carbazole (3h)^{13f}. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3h** as a white solid (66 mg, 67% yield); ¹H NMR (300 MHz, CDCl₃) δ 8.34 (d, J = 7.8 Hz, 1H), 8.02 (s, 1H), 7.43 – 7.38 (m, 2H), 7.34 (d, J = 8.0 Hz, 1H), 7.30 – 7.21 (m, 1H), 7.04 (d, J = 8.1 Hz, 1H), 6.70 (d, J = 8.0 Hz, 1H), 4.09 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 156.32, 140.92, 138.72, 126.77, 125.01, 123.12, 122.67, 119.70, 112.60, 110.07, 103.62, 100.42, 55.51; FTIR (neat) u 3386, 2938, 1405, 1248, 1184, 1045 cm⁻¹; LRMS *m/z* (EI): 197.8 (M⁺+1), 196.7, 181.9, 153.9, 128.0.

2-Chloro-9H-carbazole (3i)^{6e}. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3i** as a white solid (71 mg, 70% yield); ¹H NMR (300 MHz, DMSO) δ 11.43 (s, 1H), 8.12 (d, *J* = 8.2 Hz, 2H), 7.54 – 7.49 (m, 2H), 7.40 (t, *J* = 7.5 Hz, 1H), 7.20 – 7.14 (m, 2H); ¹³C NMR (75 MHz, DMSO) δ 140.30, 140.08, 129.88, 125.98, 121.81, 121.57, 121.31, 120.35, 119.09, 118.68, 111.23, 110.61; FTIR (neat) \cup 3393, 3128, 1636, 1406, 1321, 1178, 754 cm⁻¹; LRMS *m/z* (EI): 200.7 (M⁺-1), 165.7, 138.8, 100.3.

2-Fluoro-9H-carbazole (*3j*)^{11d}. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3j** as a white solid (75 mg, 81% yield); ¹H NMR (300 MHz, CDCl₃) δ 8.07 (s, 1H), 8.03 – 7.97 (m, 2H), 7.44 – 7.39 (m, 2H), 7.26 – 7.21 (m, 1H), 7.11 (dd, *J* = 9.6, 2.2 Hz, 1H), 7.02 – 6.93 (m, 1H); ¹³C NMR (75 MHz, DMSO) δ 161.18 (d, J = 238.6 Hz), 140.41, 140.28, 125.25, 122.07, 121.46 (d, J = 10.6 Hz), 119.97, 119.20, 119.00, 111.02, 106.49 (d, J = 24.1 Hz), 97.34 (d, J = 26.1 Hz); FTIR (neat) \cup 3411, 2919, 1362, 1263 cm⁻¹; LRMS *m/z* (EI): 186.2 (M⁺+1), 185.2, 184.3, 157.3, 92.8.

1-(9H-Carbazole-2-yl)ethan-1-one (**3k**)^{11d}. Purified Abyle flash column chromatography (SiO₂) using 7% acetone an factor for the state of th

Methyl-9H-carbazole-2-carboxylate (3I)^{6e}. Purified by flash column chromatography (SiO₂) using 7% acetone in hexanes as eluant to give **3I** as a white solid (69 mg, 61% yield); ¹H NMR (300 MHz, DMSO) δ 11.58 (s, 1H), 8.21 (t, J = 8.8 Hz, 2H), 8.12 (s, 1H), 7.78 (d, J = 8.2 Hz, 1H), 7.57 (d, J = 8.1 Hz, 1H), 7.47 (t, J = 7.5 Hz, 1H), 7.21 (t, J = 7.4 Hz, 1H), 3.89 (s, 3H); ¹³C NMR (75 MHz, DMSO) δ 166.95, 141.10, 139.05, 127.06, 126.30, 126.20, 121.60, 121.10, 120.14, 119.25, 119.16, 112.35, 111.42, 52.05; FTIR (neat) u 3337, 2937, 1738, 1365, 1216, 888 cm⁻¹; LRMS m/z (EI): 226.2 (M⁺+1), 225.2, 194.2, 166.2, 139.2, 97.2, 83.2, 69.7.

3-Methyl-9H-carbazole (3m) and 1-Methyl-9H-carbazole (3m'). Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3m/3m'** as a white solid (80 mg, 88% yield, **3m:3m'**=1:1.2 (determined by ¹H NMR and GC-Mass spectrometer)). GC (Varian 450-GC, SPBTM-680 (30 m x 0.25 mm x 0.25 µm), T₀ : 60 °C (hold 4 min), T₁ : Ramp, 250 °C at 10 °C/min, Carrier: He (1.0 mL/min), Total time 60 min), t_{3m'} = 21.95 min, t_{3m} = 22.32 min, Ratio (**3m:3m'** = 1:1.2).

3-Methyl-9H-carbazole (3m)^{6e. 1}H NMR (300 MHz, DMSO) δ 11.10 (s, 1H), 8.05 (d, J = 7.7 Hz, 1H), 7.89 (s, 1H), 7.45 (d, J = 8.1 Hz, 1H), 7.39 – 7.32 (m, 2H), 7.20 (d, J = 8.0 Hz, 1H), 7.12 (t, J = 7.4 Hz, 1H), 2.46 (s, 3H): ¹³C NMR (75 MHz, DMSO) δ 139.98, 137.96, 127.08, 126.82, 125.29, 122.54, 122.22, 120.02, 119.90, 118.23, 110.85, 110.63, 21.10; FTIR (neat) \cup 3442, 1701, 1371, 1207 cm⁻¹; LRMS *m/z* (EI): 182.2 (M⁺+1), 181.2, 180.3, 152.3, 90.2, 77.2, 63.2, 51.1.

1-Methyl-9H-carbazole (3m')^{13f}. ¹H NMR (300 MHz, DMSO) δ 11.16 (s, 1H), 8.08 (d, J = 7.8 Hz, 1H), 7.93 (d, J = 7.7 Hz, 1H), 7.50 (d, J = 8.1 Hz, 1H), 7.38 – 7.34 (m, 1H), 7.20 – 7.10 (m, 2H), 7.10 – 7.02 (m, 1H), 2.55 (s, 3H); ¹³C NMR (75 MHz, DMSO) δ 139.79, 139.04, 125.94, 125.29, 122.79, 121.95, 120.17, 120.05, 118.62, 118.46, 117.60, 111.03, 17.00; FTIR (neat) \cup 3442, 1701, 1371, 1207 cm⁻¹; LRMS *m/z* (EI): 181.2 (M⁺), 180.2, 152.3, 135.2, 77.2, 63.2, 51.1.

5*H*-[1,3]dioxolo[4,5-b]carbazole (3*n*) and 5*H*-[1,3]dioxolo[4,5-a]carbazole (3*n*'). Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3n/3n'** as a white solid (90 mg, 87% yield, **3n:3n'**=5.9:1 (determined by ¹H NMR and GC-Mass spectrometer)). GC: (Varian 450-GC, SPBTM-680 (30 m x 0.25 mm x 0.25 μm), T₀ : 60 °C (hold 4 min), T₁ : Ramp, 250 °C at 10 °C/min, Carrier: He (1.0 mL/min), Total time 60 min), t_{3n'} = 25.38 min, t_{3n} = 26.90 min, Ratio (**3n:3n'** = 5.9:1).

5H-[1,3]dioxolo[4,5-b]carbazole (3n)^{23c}. ¹H NMR (300 MHz, DMSO) δ 11.08 (s, 1H), 7.96 (d, J = 7.7 Hz, 1H), 7.62 (s, 1H), 7.40 (d, J = 8.0 Hz, 1H), 7.24 (t, J = 7.5 Hz, 1H), 7.09 – 7.02 (m,

Page 6 of 9

ARTICLE

2H), 6.03 (s, 2H); ¹³C NMR (75 MHz, DMSO) δ 146.72, 141.52, 139.56, 135.14, 123.76, 122.83, 119.26, 118.20, 115.40, 110.79, 100.69, 99.36, 92.16; FTIR (neat) \cup 3397, 1710, 1266, 1043, 936, 748 cm⁻¹; LRMS *m/z* (EI): 212.0 (M⁺+1), 211.2.

5H-[1,3]dioxolo[4,5-a]carbazole (3n'). ¹H NMR (300 MHz, DMSO) δ 11.35 (s, 1H), 8.01 (d, J = 7.7 Hz, 1H), 7.67 (d, J = 15.0 Hz, 1H), 7.42 (d, J = 8.3 Hz, 1H), 7.38 – 7.30 (m, 1H), 7.16 – 7.06 (m, 1H), 6.86 (d, J = 8.2 Hz, 1H), 6.13 (s, 2H); FTIR (neat) υ 3397, 1710, 1266, 1043, 936, 748 cm⁻¹; LRMS *m/z* (EI): 211.9 (M⁺+1), 211.1, 154.2.

3-Methoxy-2-methyl-9Hcarbazole (30)^{12k}. Purified by flash column chromatography (SiO₂) using 7% acetone in hexanes as eluant to give **30** as a yellow solid (54mg, 51% yield); ¹H NMR (300 MHz, Acetone-d₆) δ 10.10 (s, 1H), 7.96 (d, *J* = 7.7 Hz, 1H), 7.81 (s, 1H), 7.42 (d, *J* = 8.1 Hz, 1H), 7.28 – 7.23 (m, 1H), 7.13 – 7.08 (m, 1H), 7.02 (s, 1H), 3.90 (s, 3H), 2.31 (s, 3H); ¹³C NMR (75 MHz, Acetone-d₆) δ 158.18, 140.73, 140.68, 124.60, 124.12, 121.94, 119.79, 119.42, 118.89, 116.68, 111.28, 93.30, 55.67, 16.92; FTIR (neat) \cup 3313, 3069, 1783, 1689, 1404, 1270, 1169 cm⁻¹; LRMS *m/z* (EI): 212.2(M⁺+1), 211.3, 196.6, 195.7, 167.5.

1-Methoxy-2-methyl-9Hcarbazole (**30**')^{23e}. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **30**' as a yellow solid (29mg, 27% yield); ¹H NMR (300 MHz, Acetone-d₆) δ 10.16 (s, 1H), 7.99 (d, *J* = 7.7 Hz, 1H), 7.89 (d, *J* = 8.5 Hz, 1H), 7.43 (d, *J* = 8.0 Hz, 1H), 7.32 – 7.25 (m, 1H), 7.19 – 7.10 (m, 1H), 6.90 (d, *J* = 8.5 Hz, 1H), 3.91 (s, 3H), 2.40 (s, 3H); ¹³C NMR (75 MHz, Acetone-d₆) δ 156.86, 141.66, 141.42, 125.16, 124.60, 120.13, 119.56, 118.53, 118.01, 111.33, 107.65, 104.63, 56.46, 10.15; FTIR (neat) u 3313, 3069, 1783, 1689, 1404, 1270, 1169 cm⁻¹; LRMS *m/z* (EI): 212.3 (M⁺+1), 211.4, 196.6, 195.7, 167.7.

3-Methoxy-9H-carbazole (**3***p*)^{6e}. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3p** as a brown solid (59 mg, 60% yield); ¹H NMR (300 MHz, CDCl₃) δ 8.02 (d, *J* = 7.8 Hz, 1H), 7.85 (s, 1H), 7.54 (d, *J* = 2.4 Hz, 1H), 7.44 – 7.30 (m, 2H), 7.28 – 7.12 (m, 2H), 7.05 (dd, *J* = 8.7, 2.4 Hz, 1H), 3.91 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 153.88, 140.32, 134.42, 125.89, 123.76, 123.34, 120.34, 119.10, 115.16, 111.46, 110.88, 103.13, 56.14; FTIR (neat) u 3405, 1788, 1685, 1400, 1260, 1031 cm⁻¹; LRMS *m/z* (EI): 198.3 (M⁺+1), 197.4, 182.7, 181.7, 154.3, 101.5.

2,3-Dimethoxy-9H-carbazole (3q)^{23d}. Purified by flash column chromatography (SiO₂) using 7% acetone in hexanes as eluant to give **3q** as a white solid (70 mg, 62% yield); ¹H NMR (300 MHz, DMSO) δ 10.98 (s, 1H), 7.99 (d, *J* = 7.7 Hz, 1H), 7.66 (s, 1H), 7.43 (d, *J* = 8.0 Hz, 1H), 7.26 (t, *J* = 7.6 Hz, 1H), 7.08 (t, *J* = 7.4 Hz, 1H), 7.04 (s, 1H), 3.86 (s, 3H), 3.85 (s, 3H); ¹³C NMR (75 MHz, DMSO₃) δ 149.18, 143.71, 139.56, 134.63, 123.67, 123.00, 119.26, 118.12, 114.30, 110.72, 103.33, 94.62, 56.21, 55.63; FTIR (neat) u 3337, 2917, 1402, 1186, 1090 cm⁻¹; LRMS *m/z* (EI): 228.2 (M⁺+1), 227.5, 212.7, 211.7, 184.5.

2-Methoxy-7-methyl-9H-carbazole (3s)^{20d}. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3s** as a white solid (75 mg, 71% yield); ¹H NMR (300 MHz, DMSO) δ 10.98 (s, 1H), 7.88 (d, *J* = 8.5 Hz, 1H), 7.84 (d, *J* = 7.9 Hz, 1H), 7.21 (s, 1H), 6.96 – 6.88 (m, 2H), 6.73 (dd, *J* = 8.5, 2.2 Hz, 1H), 3.82 (s, 3H), 2.44 (s, 3H); ¹³C NMR (75 MHz

DMSO) δ 158.14, 141.07, 140.22, 133.50, 120,54_{Art}120,444, 120.03, 119.02, 116.32, 110.73, 107.42,^D94.53,^O55/25,^D21,65,^DFTIR (neat) \cup 3394, 1726, 1366, 1219, 1091 cm⁻¹; LRMS *m/z* (EI): 212.5 (M⁺+1), 211.6, 196.6, 195.7, 168.5.

Organic & Biomolecular Chemistry

2-Fluoro-7-methoxy-9H-carbazole (3t). Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3t** as a white solid (71 mg, 66% yield); ¹H NMR (300 MHz, CDCl₃) δ 7.97 (br s, 1H), 7.87 (d, J = 8.5 Hz, 1H),, 7.85 (d, J = 8.3 Hz, 1H), 7.06 (dd, J = 9.5, 2.3 Hz, 1H), 6.98 – 6.93 (m, 1H), 6.90 (d, J = 2.2 Hz, 1H), 6.86 (dd, J = 8.5, 2.2 Hz, 1H), 3.90 (s, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 158.87, 141.31, 120.78, 120.38, 120.25, 116.91, 108.55, 107.82, 107.50, 97.63, 97.27, 95.02, 55.78; FTIR (neat) u 3192, 2843, 1539, 1406, 1155 cm⁻¹; LRMS *m/z* (EI): 216.5 (M⁺+1), 215.5, 200.6, 199.7, 172.5, 171.5; HRMS *m/z* (EI): calcd. For C₁₃H₁₀FNO (M⁺) 215.0747, found 215.0746.

10-Methyl-7H-benzo[c]carbazole (3u)^{23f}. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3u** as a white solid (81 mg , 70% yield); ¹H NMR (300 MHz, CDCl₃) δ 8.79 (d, J = 8.3 Hz, 1H), 8.37 (s, 1H), 8.31 (s, 1H), 8.01 (d, J = 8.1 Hz, 1H), 7.84 (d, J = 8.8 Hz, 1H), 7.72 (t, J = 7.6 Hz, 1H), 7.59 (d, J = 8.8 Hz, 1H), 7.52 – 7.44 (m, 2H), 7.29 (d, J = 8.4 Hz, 1H), 2.64 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 137.41, 136.78, 130.09, 129.60, 129.27, 127.27, 126.90, 125.87, 124.27, 123.39, 122.98, 122.06, 115.22, 112.79, 110.88, 21.96; FTIR (neat) \cup 3470, 2897, 1696, 1542, 1267, 1152 cm⁻¹; LRMS *m/z* (EI): 232.7 (M⁺+1), 231.2, 230.2, 228.8, 197.2, 118.8.

2-(Tert-butyl)-7-fluoro-9H-carbazole (3v). Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3v** as a yellow solid (64 mg, 53% yield); ¹H NMR (300 MHz, DMSO) δ 11.24 (s, 1H), 8.04 (dd, *J* = 8.5, 5.7 Hz, 1H), 7.97 (d, *J* = 8.3 Hz, 1H), 7.43 (s, 1H), 7.23 (dd, *J* = 10.3, 2.1 Hz, 2H), 6.99 – 6.91 (m, 1H), 1.36 (s, 9H); ¹³C NMR (75 MHz, DMSO) δ 160.94 (d, *J* = 237.8 Hz), 148.29, 140.61, 140.44, 121.08 (d, *J* = 10.5 Hz), 119.71, 119.50, 119.20, 117.01, 107.26, 106.27 (d, *J* = 24.1 Hz), 97.30 (d, *J* = 26.0 Hz), 34.76, 31.60; FTIR (neat) u 3475, 3187, 1741, 1598, 1266 cm⁻¹; LRMS *m/z* (EI): 242.1 (M⁺+1), 241.1, 227.4, 226.5, 198.2; HRMS *m/z* (EI): calcd. For C₁₆H₁₆FN (M⁺) 241.1268, found 241.1267.

Glycozoline (3w)^{19d}. Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3w** as a yellow solid (37 mg, 35% yield); ¹H NMR (300 MHz, CDCl₃) δ 7.85 (s, 1H), 7.77 (s, 1H), 7.54 (d, J = 2.4 Hz, 1H), 7.31 – 7.21 (m, 3H), 7.06 (dd, J = 8.7, 2.5 Hz, 1H), 3.94 (s, 3H), 2.54 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 153.87, 138.69, 134.89, 128.45, 127.32, 123.76, 123.64, 120.25, 115.01, 111.41, 110.56, 103.23, 56.19, 21.54.; FTIR (neat) υ 3410, 2894, 1706, 1492, 1266, 1131 cm⁻¹; LRMS *m*/*z* (EI): 212.1 (M⁺+1), 211.2, 197.2, 196.3, 168.2, 167.2. (3w')^{20d}. Glycozolicine Purified flash column by chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3w'** as a yellow solid (41 mg, 39% yield); ¹H NMR (300 MHz, CDCl₃) δ 8.17 (s, 1H), 7.87 (s, 1H), 7.67 (d, J = 7.8 Hz, 1H), 7.34 (d, J = 8.2 Hz, 1H), 7.24 (d, J = 8.2 Hz, 1H), 7.16 (t, J = 7.8 Hz, 1H), 6.89 (d, J = 7.8 Hz, 1H), 4.00 (s, 3H), 2.54 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 145.79, 137.57, 130.24, 128.76, 127.19, 124.32, 123.96, 120.54, 119.61, 112.94, 110.71, 105.87, 55.59, 21.56; FTIR (neat) v 3410, 2894, 1706, 1492, 1266, 1131 cm⁻¹;

Published on 28 October 2015. Downloaded by UNIVERSITY OF OTAGO on 02/11/2015 10:29:46.

LRMS *m/z* (EI): 212.2 (M⁺+1), 211.3, 197.2, 196.3, 169.1, 168.2, 142.1.

 $(3x)^{20d}$. Glycozolidine Purified by flash column chromatography (SiO₂) using 7% acetone in hexanes as eluant to give **3x** as a white solid (56mg, 46% yield); ¹H NMR (300 MHz, CDCl₃) δ 7.76 (s, 1H), 7.56 (s, 1H), 7.46 (d, J = 2.2 Hz, 1H), 7.15 (d, J = 8.7 Hz, 1H), 6.96 (dd, J = 8.6, 2.3 Hz, 1H), 6.63 (s, 1H), 3.92 (s, 3H), 3.83 (s, 3H), 2.37 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 157.50, 153.90, 140.10, 134.23, 123.96, 121.47, 118.98, 116.17, 113.10, 111.10, 102.53, 92.53, 56.14, 55.52, 16.87; FTIR (neat) u 3389, 3066, 1787, 1637, 1509, 1365, 1270 cm⁻¹; LRMS *m/z* (EI): 242.3 (M⁺+1), 241.3, 227.2, 226.3, 198.2, 183.2, 154.1.

2,6-Dimethoxy-1-methyl-9H-carbazole (3x'). Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3x'** as a white solid (27mg, 22% yield). ¹H NMR (300 MHz, DMSO) δ 10.76 (s, 1H), 7.85 (d, *J* = 8.5 Hz, 1H), 7.56 (d, *J* = 2.4 Hz, 1H), 7.31 (d, *J* = 8.7 Hz, 1H), 6.91 (dd, *J* = 8.7, 2.5 Hz, 1H), 6.84 (d, *J* = 8.6 Hz, 1H), 3.85 (s, 3H), 3.82 (s, 3H), 2.33 (s, 3H); ¹³C NMR (75 MHz, DMSO). δ 155.44, 153.04, 141.29, 135.08, 123.59, 117.97, 116.79, 113.20, 111.23, 106.54, 103.49, 102.67, 56.08, 55.56, 10.24; FTIR (neat) v 3389, 3066, 1787, 1637, 1509, 1365, 1270 cm⁻¹; LRMS *m/z* (EI): 242.3 (M⁺+1), 241.3, 227.2, 226.3, 198.2, 197.2, 183.2, 154.1.

Clausenalene (3y) and 7-Methyl-10H-[1,3]dioxolo[4,5a]carbazole (3y'). Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give 3y/3y' as a white solid (61 mg, 80% yield, 3y:3y'=5.3:1 (determined by ¹H NMR and GC-Mass spectrometer)). GC: (Varian 450-GC, SPBTM-680 (30 m x 0.25 mm x 0.25 µm), T₀ : 60 °C (hold 4 min), T₁ : Ramp, 250 °C at 10 °C/min, Carrier: He (1.0 mL/min), Total time 60 min), t_{3y'} = 26.98 min, t_{3y} = 28.69 min, Ratio (**3y:3y'** = 5.3:1).

Clausenalene (3y)^{22c}. Purified by recrystallization from dichloromethane and hexanes to give **3y** as a white solid. ¹H NMR (300 MHz, Acetone-d₆) δ 10.09 (s, 1H), 7.76 (s, 1H), 7.50 (s, 1H), 7.31 (d, J = 8.2 Hz, 1H), 7.09 (dd, J = 8.2, 1.2 Hz, 1H), 6.98 (s, 1H), 6.00 (s, 2H), 2.45 (s, 3H); ¹³C NMR (75 MHz, Acetone-d₆) δ 147.94, 142.82, 139.11, 136.64, 128.26, 126.07, 124.45, 119.77, 116.76, 111.26, 101.66, 99.82, 92.85, 21.48; FTIR (neat) : υ 3400, 3009, 1728, 1366, 1219, 1093, 533 cm⁻¹; LRMS *m/z* (EI): 225.9 (M⁺+1), 225.2.

7-Methyl-10H-[1,3]dioxolo[4,5-a]carbazole (3y'). ¹H NMR (300 MHz, Acetone-d₆) δ 10.17 (s, 1H), 7.81 (s, J = 0.6 Hz, 2H), 7.61 – 7.57 (m, 1H), 7.36 (d, J = 8.2 Hz, 1H), 7.17 (dd, J = 8.2, 1.2 Hz, 1H), 6.80 (d, J = 8.2 Hz, 1H), 6.07 (s, 2H), 2.46 (s, 3H); FTIR (neat) υ 3400, 3009, 1728, 1366, 1219, 1093, 533 cm⁻¹; LRMS m/z (EI): 225.9 (M⁺+1), 225.2.

General Procedure for Suzuki coupling of 3-chlorocarbazole (3i). To a re-sealable pressure tube (13 x 100 mm) equipped with magnetic stir bar were 2-chloro-9H-carbazole 3i (0.1 mmol, 100 mol%), aryl boronic acid 2 (0.15 mmol, 150 mol%), Pd₂(dba)₃ (0.01 mmol, 10 mol%), PCy₃ (0.25 mmol, 25 mol %), K₃PO₄ (1.27 M in water, 0.34 mmol) and dioxane (2 mL, 0.05 M concentration with respect to 2-chloro-9H-carbazole 3i). The slurry was sealed and purged with Argon gas for 5 minutes. The mixture was heated at 100 °C (oil bath temperature) for 2 hr, at which point the reaction mixture was allowed to cool to ambient temperature. The reaction mRturel, 0.366466016644 saturated sodium bicarbonate solution, and extracted with dichloromethane (3x). The combined organic layers were dried with magnesium sulfate, filtered, and concentrated *in vacuo* and purified by flash column chromatography (SiO₂) and recrystallization from dichloromethane and hexanes under the conditions noted to furnish the corresponding product.

2-([1,1'-Biphenyl]-4-yl)-9H-carbazole (3ia). Purified by recrystallization from dichloromethane and hexanes to give **3ia** as a white solid (23 mg, 72% yield); ¹H NMR (300 MHz, DMSO) δ 11.35 (s, 1H), 8.20 (d, J = 8.1 Hz, 1H), 8.14 (d, J = 7.8 Hz, 1H), 7.86 (d, J = 8.1 Hz, 1H), 7.85 (s, 1H), 7.82 – 7.71 (m, 5H), 7.55 – 7.46 (m, 4H), 7.44 – 7.36 (m, 2H), 7.18 (t, J = 7.4 Hz, 1H); ¹³C NMR (75 MHz, DMSO) δ 140.41, 140.33, 140.20, 139.70, 138.77, 137.20, 129.03, 127.50, 127.21, 126.56, 125.70, 122.19, 121.94, 120.70, 120.30, 118.74, 117.72, 111.03, 108.74; FTIR (neat) υ 3403, 3101, 1753, 1691, 1530, 1368, 1206, 763 cm⁻¹; HRMS *m/z* (EI): calcd. For C₂₄H₁₇N (M⁺) 319.1361, found 319.1360.

2-(4-Fluorophenyl)-9H-carbazole (3ib). Purified by flash column chromatography (SiO₂) using 5% acetone in hexanes as eluant to give **3ib** as a yellow solid (23 mg, 88% yield); ¹H NMR (300 MHz, CDCl₃) δ 8.12 (d, J = 7.8 Hz, 1H), 8.10 (s, 1H) 8.09 (d, J = 7.4 Hz, 1H), 7.68 – 7.61 (m, 2H), 7.58 (s, 1H), 7.46 – 7.40 (m, 3H), 7.30 – 7.22 (m, 1H), 7.19 – 7.11 (m, 2H); ¹³C NMR (75 MHz, Acetone-d₆) δ 163.17 (d, J = 243.9 Hz), 141.61, 141.56, 139.27 (d, J = 3.2 Hz), 138.56, 129.87 (d, J = 8.0 Hz), 126.59, 123.71, 123.33, 121.34, 121.00, 119.91, 119.11, 116.33 (d, J = 21.5 Hz), 111.80, 109.96; FTIR (neat) υ 3400, 1650, 1367, 1223, 724 cm⁻¹; LRMS *m/z* (EI): 262.2 (M⁺+1), 262.2; HRMS *m/z* (EI): calcd. For C₁₈H₁₂FN (M⁺) 261.0954, found 261.0953.

Methyl 4-(9H-carbazol-2-yl)benzoate (3ic). Purified by recrystallization from dichloromethane and hexanes to give 3ic as a white solid (21 mg, 70% yield); ¹H NMR (300 MHz, DMSO) δ 11.39 (s, 1H), 8.22 (d, J = 8.2 Hz, 1H), 8.15 (d, J = 7.7 Hz, 1H), 8.07 (d, J = 8.4 Hz, 2H), 7.92 (d, J = 8.4 Hz, 2H), 7.81 (d, J = 1.3 Hz, 1H), 7.54 (dd, J = 8.2, 1.5 Hz, 1H), 7.52 (d, J = 8.1 Hz, 1H), 7.41 (td, J = 8.1, 0.9 Hz, 1H), 7.18 (td, J = 7.7, 0.5 Hz, 1H), 3.89 (s, 3H); ¹³C NMR (75 MHz, DMSO) δ 166.18, 145.77, 140.47, 140.28, 136.26, 129.86, 128.01, 127.21, 125.98, 122.59, 122.03, 120.83, 120.47, 118.85, 117.91, 111.11, 109.26, 52.16; FTIR (neat) υ 3388, 2139, 1719, 1363, 1218, 1095, 702 cm⁻¹; HRMS *m/z* (EI): calcd. For C₂₀H₁₅NO₂ (M⁺) 301.1103, found 301.1103.

2-(2-Methoxyphenyl)-9H-carbazole (3id). Purified by recrystallization from dichloromethane and hexanes to give **3id** as a white solid (26 mg, 95% yield); ¹H NMR (300 MHz, Acetone-d₆) δ 10.34 (s, 1H), 8.12 (d, J = 8.2 Hz, 2H), 7.67 (d, J = 0.6 Hz, 1H), 7.52 (d, J = 8.1 Hz, 1H), 7.43 - 7.39 (m, 2H), 7.37 -7.30 (m, 2H), 7.22 – 7.16 (m, 1H), 7.11 (d, J = 8.0 Hz, 1H), 7.05 (td, J = 7.4, 0.9 Hz, 1H), 3.81 (s, 3H); ¹³C NMR (75 MHz, DMSO) δ 156.58, 140.45, 140.02, 136.01, 131.16, 131.07, 129.05, 125.98, 122.63, 121.60, 121.27, 120.82, 120.53, 119.99, 119.10, 112.23, 112.02, 111.32, 55.90; FTIR (neat) u 3409, 3012, 1733, 1599, 1247, 1032, 866, 748 cm⁻¹; LRMS m/z (EI): 273.2 (M⁺), 273.9, 258.3; HRMS m/z (EI): calcd. $C_{19}H_{15}NO$ (M^{+}) 273.1155, found 273.1154.

Published on 28 October 2015. Downloaded by UNIVERSITY OF OTAGO on 02/11/2015 10:29:46.

Acknowledgements

This work was supported by the 2013 Research Fund of University of Ulsan.

Notes and references

- (a) J. Cheng, K. Kamiya and I. Kodama, *Cardiovasc. Drug. Rev.*, 2001, **19**, 152-171; (b) M.-J. Hsu, Y. Chao, Y.-H. Chang, F.-M. Ho, L.-J. Huang, Y.-L. Huang, T.-Y. Luh, C.-P. Chen and W.-W. Lin, *Biochem. Pharmacol.*, 2005, **70**, 102-112; (c) M. K. Roy, V. N. Thalang, G. Trakoontivakorn and K. Nakahara, *Br. J. Pharmacol.*, 2005, **145**, 145-155; (d) S. Oishi, T. Watanabe, J.-i. Sawada, A. Asai, H. Ohno and N. Fujii, *J. Med. Chem.*, 2010, **53**, 5054-5058; (e) T. Takeuchi, S. Oishi, T. Watanabe, H. Ohno, J.-i. Sawada, K. Matsuno, A. Asai, N. Asada, K. Kitaura and N. Fujii, *J. Med. Chem.*, 2011, **54**, 4839-4846.
- (a) Y. Zhang, T. Wada and H. Sasabe, J. Mater. Chem., 1998, 8, 809-828;
 (b) J. L. Díaz, A. Dobarro, B. Villacampa and D. Velasco, Chem. Mater., 2001, 13, 2528-2536;
 (c) K. R. Justin Thomas, J. T. Lin, Y.-T. Tao and C.-W. Ko, J. Am. Chem. Soc., 2001, 123, 9404-9411;
 (d) J. V. Grazulevicius, P. Strohriegl, J. Pielichowski and K. Pielichowski, Prog. Polym. Sci., 2003, 28, 1297-1353;
 (e) J. Li and A. C. Grimsdale, Chem. Soc. Rev., 2010, 39, 2399-2410;
 (f) A. D. Finke, D. E. Gross, A. Han and J. S. Moore, J. Am. Chem. Soc., 2011, 133, 14063-14070;
 (g) D. Kim, V. Coropceanu and J.-L. Brédas, J. Am. Chem. Soc., 2011, 133, 17895-17900.
- (a) H.-J. Knölker and K. R. Reddy, *Chem. Rev.*, 2002, **102**, 4303-4428; (b)I. Bauer and H.-J. Knölker, *Top. Curr. Chem.*, 2012, **309**, 203-253; (c) A. W. Schmidt, K. R. Reddy and H.-J. Knölker, *Chem. Rev.*, 2012, **112**, 3193-3328; (d) N. Yoshikai and Y. Wei, *Asian J. Org. Chem.*, 2013, **2**, 466-478.
- (a) T. Noji, H. Fujiwara, K. Okano and H. Tokuyama, Org. Lett., 2013, 15, 1946-1949; (b) W. D. Guerra, R. A. Rossi, A. B. Pierini and S. M. Barolo, J. Org. Chem., 2015, 80, 928-941.
- C. P. Leslie, R. D. Fabio, F. Bonetti, M. Borriello, S. Braggio, G. D. Forno, D. Donati, A. Falchi, D. Ghirlanda, R. Giovannini, F. Pavone, A. Pecunioso, G. Pentassuglia, D. A. Pizzi, G. Rumboldt and L. Stasi, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 1043-1046.
- (a) J. A. Jordan-Hore, C. C. C. Johansson, M. Gulias, E. M. Beck and M. J. Gaunt, J. Am. Chem. Soc., 2008, 130, 16184-16186; (b) W. C. P. Tsang, R. H. Munday, G. Brasche, N. Zheng and S. L. Buchwald, J. Org. Chem., 2008, 73, 7603-7610; (c) S. H. Cho, J. Yoon and S. Chang, J. Am. Chem. Soc., 2011, 133, 5996-6005; (d) K. Takamatsu, K. Hirano, T. Satoh and M. Miura, Org. Lett., 2014, 16, 2892-2895; (e) C. Suzuki, K. Hirano, T. Satoh and M. Miura, Org. Lett., 2015, 17, 1597-1600.
- (a) H.-J. Knölker and M. Bauermeister, J. Chem. Soc., Chem. Commun., 1989, 1468-1470;
 (b) H.-J. Knölker and W. Fröhner, Tetrahedron Lett., 1997, **38**, 1535-1538;
 (c) O. Kataeva, M. P. Krahl and H.-J. Knölker, Org. Biomol. Chem., 2005, **3**, 3099-3101;
 (d) R. Czerwonka, K. R. Reddy, E. Baum and H.-J. Knölker, Chem. Commun., 2006, 711-713;
 (e) K. E. Knott, S. Auschill, A. Jager and H.-J. Knölker, Chem. Commun., 2009, 1467-1469;
 (f) K. K. Gruner, T. Hopfmann, K. Matsumoto, A. Jager, T. Katsuki and H.-J. Knölker, Org. Biomol. Chem., 2011, **9**, 2057-2061.
- (a) K. Nozaki, K. Takahashi, K. Nakano, T. Hiyama, H.-Z. Tang, M. Fujiki, S. Yamaguchi and K. Tamao, *Angew. Chem. Int. Ed.*, 2003, **42**, 2051-2053;
 (b) D. Zhu, Q. Liu, B. Luo, M. Chen, R. Pi, P. Huang and S. Wen, *Adv. Synth. Catal.*, 2013, **355**, 2172-2178.
- M. Pudlo, D. Csányi, F. Moreau, G. Hajós, Z. Riedl and J. Sapi, Tetrahedron, 2007, 63, 10320-10329.
- B. J. Stokes, B. Jovanović, H. Dong, K. J. Richert, R. D. Riell and T. G. Driver, J. Org. Chem., 2009, 74, 3225-3228.
- (a) I. C. F. R. Ferreira, M.-J. R. P. Queiroz and G. Kirsch, *Tetrahedron Lett.*, 2003, 44, 4327-4329; (b) J. H. Smitrovich and I. W. Davies, *Org. Lett.*, 2004, 6, 533-535; (c) P. Appukkuttan, E. Van der Eycken and W. Dehaen, *Synlett*, 2005, 127-133; (d) A. W. Freeman, M. Urvoy and M. E. Criswell, *J. Org. Chem.*, 2005, 70, 5014-5019; (e) R. Sanz, J. Escribano, M. R. Pedrosa, R. Aguado and F. J. Arnáiz, *Adv. Synth. Catal.*, 2007, 349, 713-718; (f) J. T. Kuethe and K. G. Childers, *Adv. Synth. Catal.*, 2008, 350, 1577-1586; (g) H. Gao, Q.-L. Xu, M. Yousufuddin, D. H. Ess and L. Kürti, *Angew. Chem. Int. Ed.*, 2014, 53, 2701-2705.

 (a) B. Akermark, L. Eberson, E. Jonsson and E. Pettersson, J. Org. Chem., 1975, 40, 1365-1367; (b) H.-J. Knölker and N.O. Sullivan, Cetaghedson, 1994, 50, 10893-10908; (c) H. J. Knölker and K. R. Reddy, Heterocycles, 2003, 60, 1049-1052; (d) R. Forke, M. P. Krahl, T. Krause, G. Schlechtingen and H. J. Knölker, Synlett, 2007, 268-272; (e) R. Forke, A. Jager and H.-J. Knölker, Org. Biomol. Chem., 2008, 6, 2481-2483; (f) K. K. Gruner and H.-J. Knölker, Org. Biomol. Chem., 2008, 6, 3902-3904; (g) Kn, ouml and H.-J. Iker, Chem. Lett., 2009, 38, 8-13; (h) M. Fuchsenberger, R. Forke and H. J. Knölker, Synlett, 2011, 2056-2058; (i) T. Gensch, M. Rönnefahrt, R. Czerwonka, A. Jäger, O. Kataeva, I. Bauer and H.-J. Knölker, Chem.-Eur. J., 2012, 18, 770-776; (j) L. Huet, R. Forke, A. Jager and H. J. Knölker, Synlett, 2012, 1230-1234; (k) R. Hesse, K. K. Gruner, O. Kataeva, A. W. Schmidt and H.-J. Knölker, Chem.-Eur. J., 2013, 19, 14098-14111; (l) V. P. Kumar, K. K. Gruner, O. Kataeva and H.-J. Knölker, Angew. Chem. Int. Ed., 2013, 52, 11073-11077.

Organic & Biomolecular Chemistry

- (a) R. B. Bedford and M. Betham, J. Org. Chem., 2006, 71, 9403-9410; (b)
 L. Ackermann and A. Althammer, Angew. Chem. Int. Ed., 2007, 46, 1627-1629; (c) D. J. St. Jean, S. F. Poon and J. L. Schwarzbach, Org. Lett., 2007, 9, 4893-4896; (d) L. Ackermann, A. Althammer and P. Mayer, Synthesis-Stuttgart, 2009, 3493-3503; (e) J. K. Laha, P. Petrou and G. D. Cuny, J. Org. Chem., 2009, 74, 3152-3155; (f) S. Chakrabarty, I. Chatterjee, L. Tebben and A. Studer, Angew. Chem. Int. Ed., 2013, 52, 2968-2971.
- (a) N. Miyaura and A. Suzuki, J. Chem. Soc., Chem. Commun., 1979, 866-867; (b) N. Miyaura, K. Yamada and A. Suzuki, Tetrahedron Lett., 1979, 20, 3437-3440; (c) N. Miyaura, T. Yanagi and A. Suzuki, Synth. Commun., 1981, 11, 513-519; (d) N. Miyaura and A. Suzuki, Chem. Rev., 1995, 95, 2457-2483; (e) A. Suzuki, J. Organomet. Chem., 1999, 576, 147-168.
- (a) J. I. G. Cadogan, *Quarterly Reviews, Chemical Society*, 1962, **16**, 208-239; (b) J. I. G. Cadogan and M. Cameron-Wood, *Proc. Chem. Soc.*, 1962, 361; (c) J. I. G. Cadogan, M. Cameron-Wood, R. K. Mackie and R. J. G. Searle, *J. Chem. Soc.*, 1965, 4831-4837; (d) J. I. G. Cadogan, *Quarterly Reviews, Chemical Society*, 1968, **22**, 222-251; (e) J. I. G. Cadogan, *Synthesis*, 1969, 11-17; (f) J. I. G. Cadogan and M. J. Todd, *J. Chem. Soc. C*, 1969, 2808-2813.
- (a) H. J. Knölker, M. Bauermeister and J. B. Pannek, *Chem. Ber.*, 1992, 125, 2783-2793; (b) M. Chakrabarty, A. C. Nath, S. Khasnobis, M. Chakrabarty, Y. Konda, Y. Harigaya and K. Komiyama, *Phytochemistry*, 1997, 46, 751-755; (c) T. Watanabe, S. Oishi, N. Fujii and H. Ohno, *J. Org. Chem.*, 2009, 74, 4720-4726.
- (a) R. Forke, M. P. Krahl, F. Däbritz, A. Jäger and H.-J. Knölker, *Synlett*, 2008, 1870-1876; (b) V. Sridharan, M. A. Martín and J. C. Menéndez, *Eur. J. Org. Chem.*, 2009, 4614-4621.
- J. Lim, B. Taoka, R. D. Otte, K. Spencer, C. J. Dinsmore, M. D. Altman, G. Chan, C. Rosenstein, S. Sharma, H.-P. Su, A. A. Szewczak, L. Xu, H. Yin, J. Zugay-Murphy, C. G. Marshall and J. R. Young, *J. Med. Chem.*, 2011, 54, 7334-7349.
- (a) D. P. Chakraborty, *Phytochemistry*, 1969, **8**, 769-772; (b) H. Peng, X. Chen, Y. Chen, Q. He, Y. Xie and C. Yang, *Tetrahedron*, 2011, **67**, 5725-5731; (c) S. W. Youn, J. H. Bihn and B. S. Kim, *Org. Lett.*, 2011, **13**, 3738-3741; (d) T. Pirali, F. Zhang, A. H. Miller, J. L. Head, D. McAusland and M. F. Greaney, *Angew. Chem. Int. Ed.*, 2012, **51**, 1006-1009; (e) R. Hesse, A. W. Schmidt and H.-J. Knölker, *Tetrahedron*, 2015, **71**, 3485-3490; (f) M. S. Naykode, V. T. Humne and P. D. Lokhande, *J. Org. Chem.*, 2015, **80**, 2392-2396.
- (a) A. Chakraborty, B. K. Chowdhury, S. S. Jash, G. K. Biswas, S. K. Bhattacharyya and P. Bhattacharyya, *Phytochemistry*, 1992, **31**, 2503-2505; (b) A. K. Chakravarty, T. Sarkar, K. Masuda, T. Takey, H. Doi, E. Kotani and K. Shiojima, *Indian J. Chem. B*, 2001, **40**, 484-489; (c) D. Ding, X. Lv, J. Li, G. Xu, B. Ma and J. Sun, *Chem. Asian J.*, 2014, **9**, 1539-1542; (d) C. Schuster, C. Börger, K. K. Julich-Gruner, R. Hesse, A. Jäger, G. Kaufmann, A. W. Schmidt and H.-J. Knölker, *Eur. J. Org. Chem.*, 2014, 4741-4752.
- (a) M. Schmidt and H.-J. Knölker, *Synlett*, 2009, 2421-2424; (b) C. Ito, M. Itoigawa, A. Sato, C. M. Hasan, M. A. Rashid, H. Tokuda, T. Mukainaka, H. Nishino and H. Furukawa, *J. Nat. Prod.*, 2004, **67**, 1488-1491; (c) M. Iwao, H. Takehara, S. Furukawa and M. Watanabe, *Heterocycles*, 1993, **36**, 1483-1488.
- (a) P. Bhattacharyya, G. K. Biswas, A. K. Barua, C. Saha, I. B. Roy and B. K. Chowdhury, *Phytochemistry*, 1993, **33**, 248-250; (b) T. L. Scott, X. Yu, S. P. Gorugantula, G. Carrero-Martínez and B. C. G. Söderberg,

Organic & Biomolecular Chemistry

Tetrahedron, 2006, **62**, 10835-10842; (c) S. Rasheed, D. N. Rao, K. R. Reddy, S. Aravinda, R. A. Vishwakarma and P. Das, *RSC Adv.*, 2014, **4**, 4960-4969.

 (a) B. Alcaide, P. Almendros, J. M. Alonso, M. T. Quirós and P. Gadziński, *Adv. Synth. Catal.*, 2011, **353**, 1871-1876; (b) C. Wang, W. Zhang, S. Lu, J. Wu and Z. Shi, *Chem. Commun.*, 2008, 5176-5178; (c) S. Chakraborty, G. Chattopadhyay and C. Saha, *J. Heterocycl. Chem.*, 2013, **50**, 91-98; (d) B. Liegault, D. Lee, M. P. Huestis, D. R. Stuart and K. Fagnou, *J. Org. Chem.*, 2008, **73**, 5022-5028; (e) S. Kureel, R. Kapil and S. Popli, *Journal of the Chemical Society D: Chemical Communications*, 1969, 1120-1121; (f) B.-Y. Lim, M.-K. Choi and C.-G. Cho, *Tetrahedron Lett.*, 2011, **52**, 6015-6017. View Article Online DOI: 10.1039/C5OB01952D