



Radical cyclization of β -aminoacrylates: synthesis of bicyclic pyrrolidine derivatives

Eun Lee,* Sang Kyun Kim, Ji Young Kim and Jaehong Lim

School of Chemistry and Molecular Engineering, Seoul National University, Seoul 151-742, South Korea

Received 8 May 2000; revised 5 June 2000; accepted 9 June 2000

Abstract


Radical cyclization of the β -aminoacrylate derived from 4-(2-bromoethyl)azetidin-2-one led to the stereoselective formation of a bicyclic β -lactam product, which may serve as a precursor for *cis*-2,5-disubstituted pyrrolidines. © 2000 Elsevier Science Ltd. All rights reserved.

Radical cyclization of β -aminoacrylates was shown to be a useful method in the preparation of azacyclic compounds.¹ β -Aminoacrylates prepared from primary amines carrying secondary alkyl groups were converted into pyrrolidine and piperidine derivatives, and a useful level of stereocontrol ($\sim$ 4:1) favoring *trans*-2,5-disubstituted pyrrolidine products was ascertained when the methanesulfonamide substrates were employed.^{1a} In our continuing search for stereoselective conversions, we considered the use of β -lactam templates in radical cyclization reactions.

The known azetidin-2-one derivative **3**² may be synthesized from the carboxylic acid **1**³ via the homologous alcohol **2**.⁴ Reaction of the azetidin-2-one **3** with ethyl propiolate in the presence of tributylphosphine afforded the β -aminoacrylate **4**. Radical cyclization of **4** under the standard high dilution conditions led to the formation of the novel bicyclic β -lactam **5**⁵ in 49% yield. An epimeric product **6** was isolated in 3% yield along with the simple reduction product 4-ethylazetidin-2-one (28%) (Scheme 1). The stereochemical assignment was confirmed by comparing proton NMR spectra of the *N*-benzyl derivatives of *cis*- and *trans*-2,5-bis(ethoxycarbonylmethyl)pyrrolidine, obtained from the reaction of **5** and **6** with sodium ethoxide in ethanol and then with benzyl bromide in acetone in the presence of K₂CO₃. The benzylic methylene protons exhibit a singlet signal (δ 3.78) for the *cis* isomer and an AB quartet (δ 3.64 and 3.85, J = 14.3 Hz) for the *trans* isomer.⁶


For further comparison studies, the β -aminoacrylate **8** was prepared from the known oxazolidinone **7**.⁷ Radical cyclization of **8** in the presence of tributylstannane and AIBN resulted in the formation of an inseparable 3.3:1 mixture of the products **9** and **10** (Scheme 2). The mixture was reacted with trimethylsilyl phenylselenide in toluene under reflux in the presence of a catalytic

* Corresponding author. Fax: +82-2-889-1568; e-mail: eunlee@snu.ac.kr



Scheme 1. (a) $(\text{COCl})_2$, DCM; CH_2N_2 , ether; Ag_2O , MeOH, 55°C , 90 min; (b) LAH, ether, 0°C , 40 min; (c) CBr_4 , Ph_3P , TEA, DCM, rt, 4 h; (d) KF, AcOH, MeOH, 0°C , 10 min; (e) 2.0 equiv. HCCCCO_2Et , 1.0 equiv. Bu_3P , DCM, 0°C to rt, 90 min; (f) 1.1 equiv. Bu_3SnH , 0.1 equiv. AIBN, benzene (0.025 M), reflux, 5 h (syringe pump, 4 h)

amount of ZnI_2 ,^{1a} and *cis*- and *trans*-2-(phenylseleno)methyl-5-(ethoxycarbonylmethyl)-pyrrolidine thus produced were separated by flash column chromatography. The benzylic methylene protons of the *N*-benzyl derivatives exhibit an AB quartet at δ 3.71 and 3.77 ($J = 14.2$ Hz) for the *cis* isomer and an AB quartet at δ 3.66 and 3.91 ($J = 14.4$ Hz) for the *trans* isomer.



Scheme 2. (a) *p*-TsCl, pyridine, DCM; (b) HCCCCO_2Et , NMM, DCM, reflux; (c) LiBr, acetone, reflux; (d) 1.3 equiv. Bu_3SnH , 0.15 equiv. AIBN, benzene (0.025 M), reflux, 5 h (syringe pump, 4 h)

It is apparent from these studies that *cis*-2,5-disubstituted pyrrolidine derivatives may be prepared with useful stereoselectivity when azetidinone and oxazolidinone templates are used in the radical cyclization reactions of β -aminoacrylates, and they complement the results of earlier studies in which *trans*-2,5-disubstituted pyrrolidine derivatives were obtained as major products.

Acknowledgements

This work was supported by a grant of the Korea Science and Engineering Foundation (98-0501-05-01).

References

- (a) Lee, E.; Kang, T. S.; Joo, B. J.; Tae, J. S.; Li, K. S.; Chung, C. K. *Tetrahedron Lett.* **1995**, 36, 417–420. (b) Lee, E.; Li, K. S.; Lim, J. *Tetrahedron Lett.* **1996**, 37, 1445–1446. (c) Lee, E.; Kang, T. S.; Chung, C. K. *Bull. Kor. Chem. Soc.* **1996**, 17, 212–214.
- Pfaendler, H. R.; Strasser, R. *Liebigs Ann. Chem.* **1987**, 911–919.
- Baldwin, J. E.; Adlington, R. M.; Gollins, D. W.; Schofield, C. J. *Tetrahedron* **1990**, 46, 4733–4748.
- Umezawa, H. et al. Eur. Pat. Appl. EP 91,239, 1983; *Chem. Abstr.* **1984**, 100, 103629.
- ^{13}C NMR (50.3 MHz, CDCl_3) δ 14.58, 29.91, 36.25, 37.05, 42.81, 52.69, 57.36, 61.03, 171.74, 174.89.
- Hill, R. K.; Chan, T.-H. *Tetrahedron* **1965**, 21, 2015–2019.
- Urban, D.; Skrydstrup, T.; Beau, J.-M. *Chem. Commun.* **1998**, 955–956.