

$\text{Pr}_3)_2$ by neutron diffraction,¹⁷ 0.890 Å for $\text{H}_2\text{-W}(\text{CO})_3(\text{PCy}_3)_2$ reported here by NMR, and 0.89 Å for *trans*- $[\text{Fe}(\eta^2\text{-H}_2)(\text{H})(\text{dppe})_2]\text{BF}_4$ by X-ray.³ The latter distance is likely to be in error on the low side as X-ray diffraction will tend to underestimate the distance between the nuclei. A systematic study is currently under way to quantitate this trend by the solid-state NMR method described here. In conclusion dynamic solid-state NMR studies as described here should prove to be a powerful tool for studying the structure and fluxional behavior of this interesting class of compounds.

Acknowledgment. This work was supported by a Camille and Henry Dreyfus Foundation grant for newly appointed young faculty in chemistry. Stimulating discussions with Prof. R. H. Crabtree and Dr. J. M. Millar are gratefully acknowledged. G.J.K. is supported by the U.S. Department of Energy, Division of Chemical Sciences, Office of Basic Energy Sciences.

Registry No. $\text{H}_2\text{-W}(\text{CO})_3(\text{PCy}_3)_2$, 104198-75-6.

(17) Vergamini, P. J.; Wasserman, H. J.; Koetzle, T.; Kubas, G. J., manuscript in preparation describing a better neutron structure than reported in ref 1.

Multiple Pathways in the Oxidation of *cis*-Stilbene by Fe-Bleomycin

David C. Heimbrook,^{1a} Robert L. Mulholland, Jr.,^{1b} and Sidney M. Hecht*^{1a,b}

Chemical Research and Development
Smith Kline & French Laboratories
Swedeland, Pennsylvania 19479
Departments of Chemistry and Biology
University of Virginia
Charlottesville, Virginia 22901

Received July 14, 1986

The bleomycins are clinically useful glycopeptide-derived antitumor antibiotics.² Their therapeutic effects are believed to derive from their ability to cleave DNA, an oxidative process that can be mediated *in vitro* by $\text{Fe}^{\text{III}}\text{-BLM}$ + an oxidant (e.g., $\text{C}_6\text{H}_5\text{IO}$ or H_2O_2)³ or $\text{Fe}^{\text{II}}\text{-BLM}$ + O_2 .⁴ Bleomycin derivatives so activated have also been shown to effect the oxygenation of olefinic substrates.^{3b,5} While the products of BLM-mediated oxidation of DNA and olefins have been characterized, their mechanism(s) of formation are less well-defined. Presently, we provide evidence that oxygenation of *cis*-stilbene by activated Fe-BLM involves at least two oxidative mechanisms.

Under aerobic conditions, the oxidation of *cis*-stilbene by $\text{Fe}^{\text{III}}\text{-BLM}$ + an oxidant⁶ yielded a variety of oxidation products




Figure 1. Ratio of aerobic/anaerobic product yields. Values represent the ratios of mean product yields obtained under air vs. under argon⁶ and were determined from at least five independent trials with each oxidant under each atmosphere. *trans*-Stilbene formation with $\text{C}_6\text{H}_5\text{IO}$ was found to be BLM-independent; the ratio was therefore omitted.




Figure 2. Multiple pathways postulated for *cis*-stilbene oxidation.

including *cis*-stilbene oxide and benzaldehyde.⁵ Exclusion of dioxygen had little effect on the production of *cis*-stilbene oxide, in accord with the hypothesis that O_2 plays no direct role in the formation of this product in exogenous oxidant-supported reactions.^{5b} In contrast, in all cases the yield of benzaldehyde was

(1) (a) Smith Kline & French Laboratories. (b) University of Virginia.

(2) (a) Umezawa, H. In *Bleomycin: Current Status and New Developments*; Carter, S. K., Crooke, S. T., Umezawa, H., Eds.; Academic Press: New York, 1978; p 15 ff. (b) Hecht, S. M. In *Bleomycin: Chemical Biochemical and Biological Aspects*; Hecht, S. M., Ed.; Springer-Verlag: New York, 1979; p 1 ff.

(3) (a) Burger, R. M.; Peisach, J.; Horwitz, S. B. *J. Biol. Chem.* **1981**, *256*, 11636. (b) Murugesan, N.; Ehrenfeld, G. M.; Hecht, S. M. *J. Biol. Chem.* **1982**, *257*, 8600.

(4) (a) Sausville, E. A.; Peisach, J.; Horwitz, S. B. *Biochemistry* **1978**, *17*, 2740. (b) Sausville, E. A.; Stein, R. W.; Peisach, J.; Horwitz, S. B. *Biochemistry* **1978**, *17*, 2746.

(5) (a) Aoyagi, Y.; Suguna, H.; Murugesan, N.; Ehrenfeld, G. M.; Chang, L.-H.; Ohgi, T.; Shekhani, M. S.; Kirkup, M. P.; Hecht, S. M. *J. Am. Chem. Soc.* **1982**, *104*, 5237. (b) Murugesan, N.; Hecht, S. M. *J. Am. Chem. Soc.* **1985**, *107*, 493 and references therein.

(6) Oxidants employed included H_2O_2 , NaIO_4 , $\text{C}_6\text{H}_5\text{IO}$, and cumene hydroperoxide. In a typical experiment, 15 μL of an aqueous solution of preformed 9.1 mM $\text{Fe}^{\text{III}}\text{-BLM}$ was added to a solution composed of 150 μL of 100 mM *cis*-stilbene in methanol and 285 μL of 8:2 methanol/water. In aerobic experiments, the reaction was initiated by the addition of 50 μL of a 40 mM solution of the oxidant in 8:2 methanol/water. In anaerobic experiments, the oxidant was placed in the top section of a Thunberg tube, with the $\text{Fe}^{\text{III}}\text{-BLM}$ + *cis*-stilbene solution in the bottom section. The tube was then degassed by three vacuum/argon cycles before mixing. Reactions were typically run for 30 min at 37 °C and diluted with 2 mL of aqueous 0.5% NaCl and the products extracted into CH_2Cl_2 . Products were quantitated by gas chromatography, using *m*-tolualdehyde and benzophenone as internal standards.

much lower in the absence of O₂ (Figure 1), and ¹⁸O₂-labeling experiments confirmed that the oxygen atom in the formed C₆H₅CHO was derived predominantly (>60%) from O₂ for three of the oxidants employed. Also detected by gas chromatographic analysis as a product of *cis*-stilbene treatment with activated Fe-BLM was *trans*-stilbene. While formed only to a limited extent in the presence of O₂, *trans*-stilbene was the major product with each oxidant under anaerobic conditions (Figure 1).

That benzaldehyde formation was favored under aerobic conditions, while *trans*-stilbene was formed primarily under anaerobic conditions, suggested that these two products might be derived from a single intermediate that partitioned down one of two pathways depending on the presence or absence of O₂. One candidate for this intermediate is the stilbene cation radical, which could react with O₂ to form benzaldehyde via a dioxetane intermediate or, in the absence of O₂, isomerize to the more stable *trans* configuration and undergo reduction to *trans*-stilbene (Figure 2). It is of interest that photoexcited dyes catalyzed similar chemistry from *cis*-stilbene, with the same O₂ dependence.⁷ In contrast, BLM-mediated stilbene epoxidation was O₂ independent and occurred without loss of stereochemistry. The *cis*-epoxide is apparently formed via a completely different reaction pathway, which may more closely resemble the monooxygenase chemistry of cytochrome P-450.^{5b,8}

Three additional experiments provide support for the scheme outlined in Figure 2. First, the finding that the ratio of C₆H₅CHO to *cis*-stilbene oxide formed under aerobic conditions was essentially constant over the time course (5–30 s) of the H₂O₂-supported reaction was consistent with the formation of both from either a single "activated BLM" intermediate or from two different metal-oxygen species which are kinetically indistinguishable on this time scale. The second finding involved the oxidation of *p*-nitro-*cis*-stilbene under aerobic conditions by Fe^{III}-BLM + H₂O₂. Consistent with the scheme outlined in Figure 2, the ratio of (*p*-nitro)benzaldehyde/(*p*-nitro)-*cis*-stilbene oxide formed was 0.95 for *cis*-stilbene and 0.27 for *p*-nitro-*cis*-stilbene, in parallel with the redox potentials for the two olefins (+1.54 V for *cis*-stilbene; +1.71 V for *p*-nitro-*cis*-stilbene).

While the foregoing results are consistent with the scheme in Figure 2, they do not exclude the possible intermediacy of a stilbene radical; this could form by initial H[•] abstraction in analogy with BLM-mediated DNA oxidation.⁹ Accordingly, the oxidation of *cis*-stilbene was carried out in D₂O/CD₃OD to test this possible mechanism. When activated in deuterated solvent with any of the oxidants in Figure 1, Fe^{III}-BLM catalyzed the conversion of *cis*-stilbene to *trans*-stilbene with no detectable (<1%) deuterium incorporation. While this finding does not exclude a radical intermediate,¹⁰ given the reactivity of the stilbene intermediate toward O₂, and the proposals that many cytochrome P-450-catalyzed reactions proceed via initial electron abstraction,^{8b} the data seem most consistent with the model outlined in Figure 2 and constitute a third line of support.

The nature of the metal-oxo species responsible for the products documented here is uncertain. The similarity in results with each of the oxidants employed suggests, however, that the same intermediate or intermediates are being formed with each oxidant. It is clearly of interest to determine whether the electron abstraction pathway suggested for *cis*-stilbene oxidation is applicable

to the Fe-BLM-mediated oxidation of alkenes *in vivo*.¹¹

Acknowledgment. This study was supported at the University of Virginia by P.H.S. Grant CA38544, awarded by the National Cancer Institute.

(11) Ekimoto, H.; Takahashi, K.; Matsuda, A.; Takita, T.; Umezawa, H. *J. Antibiot.* **1985**, *38*, 1077.

Binuclear Platinum(III) Complexes. Preparation, Structure, and $d\sigma \rightarrow d\sigma^*$ Spectrum of [Bu₄N]₂[Pt₂(P₂O₅H₂)₄(CH₃CN)₂]

Chi-Ming Che,*^{1a} Thomas C. W. Mak,*^{1b}
Vincent M. Miskowski,*^{1c} and Harry B. Gray*^{1c}

Department of Chemistry
University of Hong Kong, Hong Kong
Chinese University of Hong Kong
Shatin, New Territories, Hong Kong
Contribution No. 7427, Arthur Amos Noyes Laboratory
California Institute of Technology
Pasadena, California 91125

Received June 9, 1986

An intense $d\sigma \rightarrow d\sigma^*$ absorption system is the electronic spectroscopic signature of a d^7 - d^7 binuclear complex containing a metal-metal single bond.^{2,3} Previous theoretical and experimental work has indicated²⁻⁶ that the highest occupied σ orbital may acquire appreciable axial-ligand character, and, at least in the [Pt₂(P₂O₅H₂)₄X₂]²⁻ (X = axial ligand) series,^{4,5} the absorption band positions depend so strongly on X that it has not been possible to assign a $d\sigma \rightarrow d\sigma^*$ energy to the Pt-Pt bond itself. We have addressed this problem in the latter series by utilizing CH₃CN as the axial ligand, because its very stable $\sigma(N)$ donor orbital should not significantly contaminate the highest occupied $d\sigma$ Pt-Pt level. Thus the structure and the electronic spectrum of [Pt₂(P₂O₅H₂)₄(CH₃CN)₂]²⁻ should serve as benchmarks in attempts to elucidate axial σ interactions in binuclear platinum(III) complexes.

Dropwise addition of H₂O₂ (30%, 2 mL) to an acetonitrile solution of [Bu₄N]₄[Pt₂(P₂O₅H₂)₄] (0.3 g in 50 mL) containing excess PhSSPh (1 g) yielded a bright yellow solution. Upon addition of diethyl ether, a yellow solid precipitated (> 70% yield). Recrystallization of the crude yellow solid by slow diffusion of diethyl ether into acetonitrile solution gave orange prismatic [Bu₄N]₂[Pt₂(P₂O₅H₂)₄(CH₃CN)₂] crystals together with some starting material.

(1) (a) Department of Chemistry, University of Hong Kong. (b) Department of Chemistry, the Chinese University of Hong Kong. (c) Arthur Amos Noyes Laboratory, California Institute of Technology.

(2) (a) Levenson, R. A.; Gray, H. B. *J. Am. Chem. Soc.* **1975**, *97*, 6042. (b) Pöe, A. J.; Jackson, R. A. *Inorg. Chem.* **1978**, *17*, 2330.

(3) (a) Miskowski, V. M.; Schaefer, W. P.; Sadeghi, B.; Santarsiero, B. D.; Gray, H. B. *Inorg. Chem.* **1984**, *23*, 1154. (b) Miskowski, V. M.; Smith, T. P.; Loehr, T. M.; Gray, H. B. *J. Am. Chem. Soc.* **1985**, *107*, 7925.

(4) (a) Che, C.-M.; Butler, L. G.; Grunthaner, P. J.; Gray, H. B. *Inorg. Chem.* **1985**, *24*, 4662. (b) Che, C.-M.; Lee, W. M.; Mak, T. C. W.; Gray, H. B. *J. Am. Chem. Soc.* **1986**, *108*, 4446. (c) Che, C.-M.; Schaefer, W. P.; Gray, H. B.; Dickson, M. K.; Stein, P. B.; Roundhill, D. M. *J. Am. Chem. Soc.* **1982**, *104*, 4253. (d) Che, C.-M.; Mak, T. C. W.; Gray, H. B. *Inorg. Chem.* **1984**, *23*, 4386. (e) Che, C.-M.; Herbstein, F. H.; Schaefer, W. P.; Marsh, R. E.; Gray, H. B. *J. Am. Chem. Soc.* **1983**, *105*, 4604. (f) Alexander, K. A.; Bryan, S. A.; Fronczek, F. R.; Fultz, W. C.; Rheingold, A. L.; Roundhill, D. M.; Stein, P. B.; Watkins, S. F. *Inorg. Chem.* **1985**, *24*, 2803. (g) Hedden, D.; Walkinshaw, M. D.; Roundhill, D. M. *Inorg. Chem.* **1985**, *24*, 3146.

(5) Isci, H.; Mason, W. R. *Inorg. Chem.* **1985**, *24*, 1761.

(6) Bursten, B. E.; Cotton, F. A. *Inorg. Chem.* **1981**, *20*, 3042.

(7) (a) Eriksen, J.; Foote, C. S. *J. Am. Chem. Soc.* **1980**, *102*, 6083. (b) Lewis, F. D.; Petisce, J. R.; Oxman, J. D.; Nepras, M. J. *J. Am. Chem. Soc.* **1985**, *107*, 203.

(8) White, R. E.; Coon, M. J. *Annu. Rev. Biochem.* **1980**, *49*, 315 and references therein. (b) Guengerich, F. P.; Macdonald, T. L. *Acc. Chem. Res.* **1984**, *17*, 9.

(9) (a) Giloni, L.; Takeshita, M.; Johnson, F.; Iden, C.; Grollman, A. P. *J. Biol. Chem.* **1981**, *256*, 8608. (b) Wu, J. C.; Kozarich, J. W.; Stubbe, J. *J. Biol. Chem.* **1983**, *258*, 4694. (c) Wu, J. C.; Kozarich, J. W.; Stubbe, J. *Biochemistry* **1985**, *24*, 7562.

(10) It is possible that the stilbene radical intermediate might collapse with H[•] abstraction from Fe-BLM. However, the observation that the stilbene intermediate reacts with O₂ in solution, and not with the oxygen of the activated Fe-BLM, suggests that it diffuses away from the BLM "active site" before reacting with O₂.