www.rsc.org/obc

Synthesis of achiral and racemic catenanes based on terpyridine and a directionalized terpyridine mimic, pyridyl-phenanthroline

Jon C. Loren,^a Peter Gantzel,^a Anthony Linden^b and Jay S. Siegel*^b

^a The Department of Chemistry, University of California San Diego, 9500 Gilman Drive, La Jolla, California 92093-0358, USA

^b The Organic Chemistry Institute, University of Zürich, Winterthurerstrasse 190, 8057, Zürich, Switzerland. E-mail: jss@oci.unizh.ch

Received 29th April 2005, Accepted 12th July 2005

First published as an Advance Article on the web 2nd August 2005

Concatenated macrocycles containing manisyl-substituted tridentate ligands 2,2':6',2"-terpyridine and 2-pyridin-2-yl-1,10-phenanthroline (simply referred to as terpyridine and pyridyl-phenanthroline herein) have been prepared *via* dual cyclization procedures. The manisyl derivative (manisyl = 4-methoxy-2,6-dimethylphenyl) was chosen for its ability to improve solubility while simultaneously incorporating functionality. Deprotection of the methoxy groups provided a soluble ligand that was re-alkylated with an array of terminal alkyne and alkene linkers. The tridentate coordinating ability of these ligands enabled complexation with Ru(II) and Fe(II), generating achiral and racemic octahedral complexes for terpyridine and pyridyl-phenanthroline, respectively. Subsequent macrocyclization *via* olefin metathesis or copper-mediated alkyne coupling afforded the corresponding catenanes, and in some cases a figure-eight macrocycle. The difference in symmetry and the presence of the manisyl group allowed the distinction between the catenane and the undesired figure-eight to be made directly by ¹H NMR. Metal-free achiral and racemic catenanes were obtained by liberating Fe(II) from the octahedral bound title ligands by treatment with hydrogen peroxide.

Introduction

Molecular representations of complex topologies are aesthetically attractive constructs that present challenges in structural design and synthesis.^{1,2} Borromean Links,^{2a} catenanes,^{2b} and knots^{2b} are representative of this class of structures, and along with their precursors, have been noticed for their potential applications.³ Many of these assemblies rely on preorganized templates⁴ that include rigid complexes between chelating ligands and transition metals.⁵ This approach, termed metal-templated, requires a coordinating component, of which the polypyridines, 2,2'-bipyridine,⁶ 1,10-phenanthroline,⁷ and 2,2':6',2"-terpyridine⁸ are commonly employed and are of considerable interest to our research group.^{9,10}

The bidentate ligand 1,10-phenanthroline and its tetrahedral coordination complex with copper(1) is the pioneering model for metal-templated synthesis of molecular catenanes.¹¹ The ease of introducing alkyl- and aryl-substituents into the 2- and 9-positions of commercially available 1,10-phenanthroline makes this ligand synthetically desirable for scaffolding.¹² In contrast, there are fewer examples of catenanes where the template is an octahedral center,¹³⁻¹⁵ or that include terpyridine within their framework.^{16,17} The first catenane built solely around terpyridine precluded removal of the templateing metal,¹⁵ while subsequent attempts to prepare a template-free version resulted in exclusive formation of a figure-eight macrocycle.¹⁸

Chiral variants of a catenane can be made by introducing asymmetry in the tethering or ring-closing steps,^{14,19} or by incorporating directionality within the macrocycles.²⁰ For example, coordination complexes of the directionalized terpyridine mimic, pyridyl-phenanthroline,²¹ are chiral and make practical building blocks for concatenated species.

Advances in poly-pyridine ligand synthesis, *via* cross-coupling methods, makes terpyridine and pyridyl-phenanthroline readily accessible;^{9,22,23} and thus, prime candidates for the synthesis of achiral and racemic catenanes, respectively. To broaden the family of concatenated topological isomers and demonstrate proof-of-principle, a series of catenanes from octahedral Ru(II) and Fe(II) templates of manisyl-substituted terpyridine and pyridyl-phenanthroline were targeted.²⁴ Olefin metathesis²⁵ and

copper-mediated alkyne coupling²⁶ for macrocyclization were implemented, and the relative advantages of each method are discussed. Symmetry analysis for the identification of catenane *vs.* figure-eight isomeric products by counting manisyl signals in the ¹H or ¹³C NMR is presented as confirmation of the stereoselective synthesis. To our knowledge, the synthesis of metal-free achiral/racemic catenanes from these ligands has not been reported.¹⁷

Results and discussion

The metal-templated synthesis of catenanes commences with tridentate ligands 3a and 4a, previously prepared from commercially available and literature-known intermediates in two and three steps, respectively.²³ Deprotection of the methoxy groups with molten pyridinium hydrochloride provides soluble phenolic derivatives, 3b and 4b, that serve as backbones to the chiral and racemic catenanes presented here (Scheme 1). The terminal acetylene and olefin linkers used for macrocyclization, 21-24b, were prepared in two steps as indicated from commercially available materials (Scheme 2).²⁷



Ru(II)-templated catenanes

Coordination of ligand **3b** with Ru(II) was achieved by reaction with $RuCl_2(DMSO)_4$ in refluxing ethylene glycol.²⁸ Aqueous workup with potassium hexafluorophosphate provided the



Scheme 2 Reagents and conditions: a) NaH, 0 °C \rightarrow RT, 5-bromopentene, reflux, 12 h, 55%. b) Triethylamine, tosyl chloride, DMAP, dichloromethane, 0 °C \rightarrow RT, 87% for **21b** and 78% for **23b**. c) NaH, 0 °C \rightarrow RT, 6-chlorohexyne, reflux, 15 h, 56%. d) NaH, 0 °C \rightarrow RT, allyl bromide, reflux, 12 h, 76%. e) NaH, 0 °C \rightarrow RT, 1,2-dichloroethane, reflux, 15 h, 67%. f) NaH, 0 °C \rightarrow RT, propargyl bromide, reflux, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 1,2-dichloroethane, reflux, 15 h, 67%. f) NaH, 0 °C \rightarrow RT, propargyl bromide, reflux, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 1,2-dichloroethane, reflux, 15 h, 67%. f) NaH, 0 °C \rightarrow RT, propargyl bromide, reflux, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 1,2-dichloroethane, reflux, 15 h, 67%. f) NaH, 0 °C \rightarrow RT, propargyl bromide, reflux, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 1,2-dichloroethane, reflux, 15 h, 67%. f) NaH, 0 °C \rightarrow RT, propargyl bromide, reflux, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 1,2-dichloroethane, reflux, 15 h, 67%. f) NaH, 0 °C \rightarrow RT, propargyl bromide, reflux, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 1,2-dichloroethane, reflux, 15 h, 67%. f) NaH, 0 °C \rightarrow RT, propargyl bromide, reflux, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 1,2-dichloroethane, reflux, 15 h, 67%. f) NaH, 0 °C \rightarrow RT, propargyl bromide, reflux, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 1,2-dichloroethane, reflux, 15 h, 67%. f) NaH, 0 °C \rightarrow RT, propargyl bromide, reflux, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 1,2-dichloroethane, reflux, 15 h, 67%. f) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g) NaH, 0 °C \rightarrow RT, 10 h, 56%. g)



Scheme 3 *Reagents and conditions*: a) $RuCl_2(DMSO)_4$, ethylene glycol, 160 °C, 12 h, 91%. b) **21b**, Cs_2CO_3 , CH_3CN , reflux, 12 h, 66%. c) $RuCl_2(PCy_3)_2CHPh$, CH_2Cl_2 , 60 h, 74%. d) H_2 , 10% Pd/carbon, CH_2Cl_2 -EtOH, 12 h, 95%.

octahedral complex **5** as a deep red crystalline solid. As a result of the C_{2v} symmetry of **3b**, complex **5** has D_{2d} symmetry and is the template to the Ru(II) achiral catenanes. Alkylation of the tetra-phenolate of **5** with the terminal olefin linker **21b** using cesium carbonate in acetonitrile afforded pre-catenane **6**. Exposure of **6** to high dilution olefin metathesis conditions with Grubbs catalyst,²⁹ RuCl₂(PCy₃)₂CHPh, followed by immediate catalytic hydrogenation, generated both the catenane **7a** and the unexpected figure-eight macrocycle **7b** in a 1 : 2 ratio (Scheme 3).

Distinction between **7a** and **7b** was made directly by ¹H NMR due to the difference in molecular symmetry (D_{2d} and D_2 , respectively) and the orthogonal conformation of the manisyl-pyridyl plane. Comparison of the signal for the manisyl aromatic proton **f** (~6.6 ppm), in compounds **6**, **7a** and **7b** is displayed in Fig. 1. After cyclization, this signal remains a singlet of

isochronous enantiotopic signals for catenane **7a**, and splits into two diastereotopic singlets, **f** and **f'**, for figure-eight **7b** due to a reduced local symmetry, C_s to C_1 , in the manisyl group. This is also observed for the protons of the methyl groups on manisyl (not pictured here). Additionally, the slight up-field shift in the peripheral pyridyl proton **a** associated with catenane formation is also observed for **7a** while those signals for **7b** remain relatively unchanged.

An undesired feature of olefin metathesis was the lack of thermodynamic drive to produce products and consume starting materials or intermediates. Tedious chromatographic separation was required in order to remove unreacted 6, as well as the mono-ring-closed adducts of catenane as well as figureeight. To circumvent this matter and to eliminate the need for hydrogenation we opted for a copper-mediated acetylenic



11a/b R = -C₆H₄-C₆H₄O-

Scheme 4 Reagents and conditions: a) 23b or 24b, Cs_2CO_3 , CH_3CN , reflux, 12 h, 54% for 8 and 70% for 10. b) $Cu(OAc)_2$, CH_3CN , reflux, 12–15 h, 61% for 9a/b and 57% for 11a/b.

coupling using modified Eglington conditions as the cyclization step.

Pre-catenane 8 was prepared in an analogous manner to 6 using linker 23b, which replaces the terminal olefins by terminal acetylenes. High dilution acetylenic coupling with excess copper(II) acetate gave the catenane 9a and the figureeight 9b in approximately the same 1 : 2 ratio obtained for 7a and 7b by the olefin metathesis-hydrogenation procedure; however, precursor 8 was completely consumed to give 9a and 9b as monomeric products that are easily separable by column chromatography (Scheme 4). A similar symmetry analysis to that described above allowed the structural assignment.

Synthesis of the racemic modification of a chiral catenane was achieved using an analogous coupling strategy to that which afforded **9a** and **9b**. Coordination of the asymmetric tridentate

ligand 4b with Ru(II) using RuCl₂(DMSO)₄ in ethylene glycol gave complex 12 in racemic form (Scheme 5). Here the C_s symmetry of 4b renders complex 12 with C_2 symmetry and provides the template for a racemic catenane. Alkylation of the racemic tetra-phenolate of 12 with linker 23b gave precatenane 13. Exposure of 13 to excess copper(II) acetate under high dilution conditions provided the chiral catenane 14a and a set of diastereomeric figure-eights 14b/c (not reported in detail due to the difficulties in their separation) in a 1 : 2 ratio.

The chiral nature of the pyridyl-phenanthroline Ru(II) complex makes distinction between catenane and figure-eight more difficult. The local symmetry of the manisyl group is lost upon coordination to Ru(II) and requires the distinction to be made on the basis of chemical shift changes in the ¹H NMR after cyclization. As in the reaction to form **9a** and **9b**, this reaction



Scheme 5 Reagents and conditions: a) $RuCl_2(DMSO)_4$, ethylene glycol, 160 °C, 12 h, 87%. b) 23b, Cs_2CO_3 , CH_3CN , reflux, 12 h, 56%. c) $Cu(OAc)_2$, CH_3CN , reflux, 12 h, 51% for 14a and 14b/c.

yields two sets of products that are easily separable by column chromatography from each other and from polymeric material. The racemic catenane **14a** product displays a single set of ¹H NMR aromatic signals with the expected up-field shift in the pyridyl-phenanthroline peripheral protons; whereas product composed of the diastereomeric mixture **14b/c** displays two sets of ¹H NMR signals in the aromatic region with little change in the chemical shift of the peripheral protons. High-resolution MALDI mass spectroscopy confirms that **14a** and **14b/c** are isomers with the expected monomeric mass.

Selective formation of the catenane was attempted by incorporating a biphenyl linker with the hope of sterically discouraging figure-eight formation. Alkylation of **5** with linker **24b** provided precursor **10**.³⁰ Exposure of **10** to acetylenic coupling conditions unfortunately still provided catenane **11a** and figure-eight **11b**, albeit in an improved 1 : 1 ratio (Scheme 4). Incorporating the biphenyl linker coincided with formation of X-ray quality crystals of **11a**, providing proof of structure and confirmation to our ¹H NMR assignment of the catenanes (Fig. 2).³¹ Due to the thermodynamic stability of the octahedral Ru(II) complexes, removal of the ruthenium atom to form the metal-free catenanes was not effected.



Fig. 2 X-Ray crystal structure of Ru(II) [2]catenane 11a. Solvent molecules and PF_6^- counter-ion omitted for clarity.

Fe(II)-templated catenanes

Octahedral coordination complexes between Fe(II) and tridentate ligands produce complexes that are stable under mild conditions, but are labile under basic and oxidative conditions by way of Fe(III). To avoid complications possibly arising from a more labile metal system, alkylations were completed prior to coordination with Fe(II), and cyclization via copper-mediated actylenic coupling was not explored fully. Alkylation of 3b with biphenyl linker 22b using cesium carbonate gave ligand 15 (Scheme 6). Coordination of 15 with Fe(II) was achieved by treatment with iron(II) sulfate in acetone-water. Aqueous workup with potassium hexafluorophosphate gave pre-catenane 16 in its pure form. The mild conditions of olefin metathesis allowed for cyclization and preserved the Fe(II) complex. High dilution olefin metathesis of 16 surprisingly resulted in exclusive formation of the catenane and its mono-ring closed adduct as the only monomeric products; no figure-eight isomer was detected. The catenane was isolated as a mixture of cis/trans isomers in a 30% yield and immediately hydrogenated with catalytic Pt/alumina to afford 17. As in the analysis of 7a, 9a, and 11a, the symmetry of the manisyl group aids in the ¹H NMR identification of the catenane isomer. The expected up-field shifts in the pyridyl-proton aromatic signals are also observed. Removal of the Fe(II) template was achieved by the slow addition of aqueous hydrogen peroxide to a basic solution of 17 in acetonitrile-water. The purified achiral catenane 1 was isolated in 59% yield. The characteristic fragmentation pattern in the high resolution MALDI mass spectrum of 1 confirms formation of the catenane (Fig. 3).



Fig. 3 MALDI high resolution mass spectrum of the template-free catenane 1.



Scheme 6 Terpyridine = terpy; pyridyl-phenanthroline = pphen. *Reagents and conditions*: a) **22b**, Cs_2CO_3 , DMF, 80 °C, 12 h, 88% for **15** and 92% for **18**. b) $FeSO_4 \cdot 7H_2O$, acetone– H_2O , 98% for **16** and 91% for **19**. c) $RuCl_2(PCy_3)_2CHPh$, CH_2Cl_2 , 72 h, 30% for **17**-(*cis/trans*) and 21% for **20**-(*cis/trans*). d) Pt/alumina, CH_2Cl_2 –EtOH, 16 h, 91% for **17** and 96% for **20**. e) H_2O_2 , NaOH, CH_3CN – H_2O 1 : 1, 59% for **1** and 20% for **2**.

Racemic Fe(II)-templated catenane was prepared in an analogous fashion to that used to prepare 17. Alkylation of 4b with linker 22b afforded 18. Complex 19 was prepared by coordination between 18 and Fe(II) using iron(II) sulfate. High dilution olefin metathesis of 19 yields similar results to those seen for the reaction of 16; only the catenane is produced in a 21% yield as a mixture of cis/trans isomers. Unreacted starting material and the mono-ring closed adduct comprise the majority of recovered materials and could be resubmitted to the reaction conditions to produce more product. Immediate hydrogenation of the cis/trans mixture with catalytic Pt/alumina gave the catenane 20. Identification of 20 can be made by the characteristic up-field shifts in the aromatic region of the 1H NMR spectrum. Removal of the Fe(II) was achieved by oxidation with hydrogen peroxide under basic conditions to give the racemic catenane 2 in 20% yield.

We conjecture that the absence of the figure-eight macrocycle arises from the decreased bite angle that the tridentate ligands form around Fe(II) relative to Ru(II). CPK modeling of **16** indicates that with enough pinching of the ligand around the metal center, the terminal olefins on opposing ligands become distant enough to prevent figure-eight formation, resulting in the preferential formation of the catenane.

To test this conjecture, the parent homoleptic complexes $Ru(3a)_2 \cdot 2PF_6$ and $Fe(3a)_2 \cdot 2PF_6$ were prepared and suitable crystals for X-ray crystallography were grown from slow evaporation of a dichloromethane solution in the presence of benzene (Fig. 4).³²

X-Ray analysis reveals the N_1 –M– N_3 ligand–metal bite angles for $Ru(3a)_2 \cdot 2PF_6$ and $Fe(3a)_2 \cdot 2PF_6$ to be 158.4(2) and 161.7(2)° respectively. Furthermore, the distances between opposing manisyl oxygen atoms of the same ligand are 16.646(6) and 16.481(6) Å for $Ru(3a)_2 \cdot 2PF_6$ and $Fe(3a)_2 \cdot 2PF_6$ respectively, and indicate an increased pinching of the ligands around Fe(II)relative to the Ru(II) cognate. This is further supported by the difference in N_1 – N_3 distances, roughly 0.15 Å. Finally, the



Fig. 4 X-Ray crystal structure and selected data for $Ru(3a)_2 \cdot 2PF_6$ (left) and $Fe(3a)_2 \cdot 2PF_6$ (right). Counter-ions omitted for clarity.

Downloaded by North Carolina State University on 27 February 2013 Published on 02 August 2005 on http://pubs.rsc.org | doi:10.1039/B506101F smaller ligand proximity around Fe(II) is demonstrated by the distances between central nitrogen atoms on terpyridine, $N_2-N_{2'}$, which vary by 0.19 Å and are consistent with our notion about the basis for the absence of figure-eight macrocycle formation with the Fe(II) templates.

Conclusions

We have demonstrated successfully that achiral and racemic catenanes can be synthesized using a metal-templated approach with terpyridine and pyridyl-phenanthroline ligands with either Ru(II) or Fe(II). Moreover, we have debuted pyridyl-phenanthroline in its first racemic templated topological isomer synthesis. Macrocyclization *via* olefin metathesis or coppermediated alkyne coupling with Ru(II)-templated pre-catenanes resulted in both the catenane and the undesired figure-eight macrocycle, while olefin metathesis with Fe(II)-templated pre-catenanes resulted in preferential formation of a catenane. The labile Fe(II) core could easily be liberated to yield template-free achiral and racemic catenanes. Additionally, the aryl substituent manisyl served as an internal ¹H NMR probe for topological isomer identification of the achiral catenane.

Experimental

Materials and methods

¹H and ¹³C NMR spectra were recorded on Varian (Mercury 300/400 MHz and Unity 500 MHz) spectrometers with tetramethylsilane (TMS) as an internal standard. High-resolution mass spectral (HRMS) analyses were performed by the University California Riverside mass spectrometry facility in either MALDI or EI mode, as indicated for each compound. All experiments were carried out under argon in freshly distilled anhydrous solvents unless otherwise noted. Commercial chemicals were used as supplied. Column chromatography was performed on neutral aluminum oxide (Brockmann II) and silica gel (230-425 mesh) from Fisher Scientific Co. Centrifugal chromatography was performed on a Harrison Research Chromatotron with 1 mm silica gel plates. UV-Vis spectra were recorded on a Perkin-Elmer UV/VIS/NIR spectrometer Lambda 19. Melting points were recorded on a Mel-Temp Laboratory Device and are uncorrected. Overlapping, non-symmetry-equivalent ¹³C NMR signals for pyridyl-phenanthroline adducts are indicated by *.

5',5"-**Bis(4-hydroxy-2,6-dimethylphenyl)-2,2**'.**6**',2"-terpyridine (**3b**). To anhydrous molten pyridine hydrochloride (11.00 g, 90 mmol) at 150 °C was added **3a** (1.00 g, 2.1 mmol) and heated to 185 °C for 4 h. The hot solution was quenched with boiling water (20 mL). The resulting yellow precipitate was filtered, washed thoroughly with water, aqueous NH₄OH, and dried under vacuum overnight to yield a tan solid (0.92 g, 97%). Mp 212 °C (dec.). ¹H NMR (400 MHz, CDCl₃, δ): 8.75 (d, *J* = 8.0 Hz, 2H), 8.53 (d, *J* = 2.0 Hz, 2H), 8.52 (d, *J* = 8.0 Hz, 2H), 8.03 (t, *J* = 8.0 Hz, 1H), 7.70 (dd, *J* = 2.0, 8.0 Hz, 2H), 6.68 (s, 4H), 2.07 (s, 12H). ¹³C NMR (100 MHz, CH₃OD, δ): 158.43, 152.93, 151.27, 148.22, 144.49, 140.85, 140.64, 138.45, 128.12, 124.05, 124.02, 115.57, 21.21. EI HRMS *m/z*: found 473.2111 (M⁺); calcd (C₃₁H₂₇N₃O₂) 473.2103.

8-(4-Hydroxy-2,6-dimethylphenyl)-2-[5-(4-hydroxy-2,6-dimethylphenyl)pyridin-2-yl]-1,10-phenanthroline (4b). To anhydrous molten pyridine hydrochloride (11.00 g, 90 mmol) at 150 °C was added 4a (1.00 g, 2.0 mmol) and heated to 185 °C for 4 h. The hot solution was quenched with boiling water (20 mL). The resulting yellow precipitate was filtered, washed thoroughly with water, aqueous NH₄OH, and dried under vacuum overnight to yield a yellowish solid (0.84 g, 93%). Mp 264–266 °C (dec.). ¹H NMR (500 MHz, DMSO- d_6 , δ): 9.42 (bs, 1H), 9.37 (bs, 1H), 8.96 (d, J = 2.0 Hz, 1H), 8.91 (d, J = 8.0 Hz, 1H), 8.66 (d, J = 8.0 Hz, 1H), 8.54 (d, J = 2.0 Hz, 1H), 8.09 (d, J = 2.0 Hz, 1H), 8

9.0 Hz, 1H), 8.04 (d, J = 9.0 Hz, 1H), 7.87 (dd, J = 2.0, 8.0 Hz, 1H), 6.64 (s, 2H), 6.60 (s, 2H), 1.99 (s, 6H), 1.98 (s, 6H). ¹³C NMR (125 MHz, DMSO- d_6 , δ): 156.98, 156.84, 155.14, 153.86, 151.42, 150.12, 145.06, 143.89, 138.92, 137.56, 137.26, 137.09, 137.04, 137.01, 135.84, 128.79, 128.59, 128.38, 128.26, 127.22, 126.83, 121.42, 119.95, 114.57, 114.51, 20.85, 20.77. EI HRMS m/z: found 498.2192 (MH⁺); calcd (C₃₃H₂₈N₃O₂) 498.2182.

2-[2-(2-Pent-4-enyloxyethoxy)ethoxy]ethanol (21a). To a solution of triethylene glycol (15.0 g, 100 mmol) in THF (300 mL) at 0 °C was slowly added NaH (2.8 g, 70 mmol, 60% in mineral oil) as a solid. The resulting solution was allowed to reach room temperature while stirring over 1 h. Subsequently, 5bromopentene (11.2 g, 75 mmol) was added via syringe and the contents heated at reflux for 12 h. The reaction was cooled to room temperature and the solvent was reduced to one half on a rotary evaporator. The residue was extracted with ether (200 mL) and water (100 mL). The organic layer was separated, dried over magnesium sulfate, filtered, and solvent evaporated to give an oil. The crude oil was purified by column chromatography on silica gel with hexane-ethyl acetate 1 : 3 as the eluant to yield a clear yellow viscous oil, $R_f = 0.30$ (8.43 g, 55%). ¹H NMR $(400 \text{ MHz}, \text{CDCl}_3, \delta)$: 7.74 (m, 1H), 4.94 (d, J = 17.0 Hz, 1H), 4.88 (d, J = 10.0 Hz, 1H), 3.65 (bs, 2H), 3.60–3.58 (m, 6H), 3.56-3.52 (m, 4H), 3.40 (t, J = 8.0 Hz, 2H), 2.95 (bs, 1H), 2.04 (q, J = 7.0 Hz, 2H), 1.62 (q, J = 7.0 Hz, 2H). ¹³C NMR (100 MHz, CDCl₃, δ): 137.85, 114.39, 72.33, 70.49, 70.39, 70.34, 70.11, 69.83, 61.43, 30.08, 28.59. EI HRMS m/z: found 219.1593 (MH⁺); calcd (C₁₁H₂₃O₄) 219.1596.

Toluene-4-sulfonic acid 2-[2-(2-pent-4-enyloxyethoxy)ethoxy] ethyl ester (21b). To an anhydrous solution of 21a (8.43 g, 39 mmol), triethylamine (8.07 mL, 58 mmol) and 4-(dimethylamino)pyridine (0.47 g, 3.9 mmol) in dichloromethane (500 mL) at 0 °C was added *p*-toluenesulfonyl chloride (7.35 g, 39 mmol) as a solid. The reaction was slowly warmed to room temperature and stirred for an additional 8 h. The crude reaction mixture was quenched with water (150 mL), the organic layer separated, and the aqueous layer extracted with fresh dichloromethane (2×50 mL). The combined organic layers were dried over magnesium sulfate, filtered, and reduced to dryness to yield a viscous oil. The crude oil was purified by column chromatography on silica gel with hexane-ethyl acetate 1 : 1 as the eluant to yield a clear and colorless viscous oil, $R_{\rm f} = 0.50$ (3.50 g, 87%). ¹H NMR (400 MHz, CDCl₃, δ): 7.79 (d, J =8.0 Hz, 2H, 7.34 (d, J = 8.0 Hz, 2H), 5.80 (m, 1H), 4.99 (d, J = 100 Hz)17.0 Hz, 1H), 4.94 (d, J = 10.0 Hz, 1H), 4.15 (t, J = 5.0 Hz, 2H), 3.68 (t, J = 5.0 Hz, 2H), 3.62–3.54 (m, 8H), 3.46 (t, J =6.0 Hz, 2H), 2.44 (s, 3H), 2.10 (q, J = 7.0 Hz, 2H), 1.67 (q, J = 7.0 Hz, 2H). ¹³C NMR (100 MHz, CDCl₃, δ): 144.41, 137.85, 132.54, 129.47, 127.58, 114.37, 70.47, 70.41, 70.38, 70.27, 69.83, 69.09, 68.40, 30.09, 28.62, 21.52. EI HRMS m/z: found 390.1952 (MNH₄⁺); calcd (C₁₈H₃₂NO₆S) 390.1950.

4'-Allyloxybiphenyl-4-ol (22a). To a 0 °C solution of 4,4'biphenol (10.0 g, 54 mmol) in THF (125 mL) was added NaH (1.5 g, 38 mmol, 60% in mineral oil) as a solid. The resulting solution was allowed to reach room temperature while stirring over 1 h. Subsequently, allyl bromide (4.88 g, 40 mmol) was added via syringe and the contents heated at reflux for 12 h. The crude reaction mixture was cooled to room temperature, filtered, and reduced to dryness. The crude white solid was purified by column chromatography on silica gel with dichloromethanemethanol 50 : 1 as the eluant to yield a white crystalline solid, $R_{\rm f} = 0.25$ (5.72 g, 66%). Mp 172–173 °C. ¹H NMR (500 MHz, $CDCl_3, \delta$): 7.45 (d, J = 9.0 Hz, 2H), 7.43 (d, J = 9.0 Hz, 2H), 6.97 (d, J = 9.0 Hz, 2H), 6.88 (d, J = 9.0 Hz, 2H), 6.08 (m, 1H), 5.43(dd, J = 2.0, 18.0 Hz, 1H), 5.30 (dd, J = 2.0, 11.0 Hz, 1H), 4.75(s, 1H), 4.58 (d, J = 5.0 Hz, 2H).¹³C NMR (125 MHz, CDCl₃, δ): 157.83, 154.68, 133.78, 133.64, 133.38, 128.03, 127.76, 117.78,

Downloaded by North Carolina State University on 27 February 2013 "ublished on 02 August 2005 on http://pubs.rsc.org | doi:10.1039/B506101F 115.63, 115.05, 68.90. EI HRMS m/z: found 226.1000 (M⁺); calcd (C₁₅H₁₄O₂) 226.0994.

4'-Allyloxy-4-(2-chloroethoxy)biphenyl (22b). To an anhydrous solution of 22a (2.84 g, 13 mmol) in THF (150 mL) at 0 °C was added NaH (0.53 g, 13 mmol, 60% in mineral oil) as a solid. The resulting solution was stirred for 1 h while warming to room temperature. Subsequently, an excess of 1,2dichloroethane (50 mL) was added and the contents heated to reflux for 15 h. After cooling the reaction to room temperature, diethyl ether (100 mL), and water (50 mL) were added. The organic layer was separated and the aqueous layer extracted with fresh diethyl ether (2 \times 50 mL). The combined organic layers were dried over magnesium sulfate, filtered, and the solvent removed to yield a crude yellow solid. The crude product was purified by column chromatography on silica gel with hexanedichloromethane 1:3 as the eluant to yield a white crystalline solid, $R_{\rm f} = 0.75$ (1.70 g, 47%). Mp 142–144 °C. ¹H NMR $(500 \text{ MHz}, \text{CDCl}_3, \delta)$: 7.49 (d, J = 8.0 Hz, 2H), 7.46, (d, J =8.0 Hz, 2H), 6.98 (d, J = 8.0 Hz, 2H), 6.97 (d, J = 8.0 Hz, 2H), 6.08 (m, 1H), 5.44 (dd, J = 2.0, 17.0 Hz, 1H), 5.31 (d, J = 2.0, 10.0 Hz, 1H), 4.58 (d, J = 4.0 Hz, 2H), 4.26 (t, J = 6.0 Hz, 2H), 3.83 (t, J = 6.0 Hz, 2H). ¹³C NMR (125 MHz, CDCl₃, δ): 157.92, 157.39, 134.26, 133.47, 133.36, 127.87, 127.78, 117.74, 115.05, 115.04, 68.86, 68.10, 41.83. EI HRMS m/z: found 288.0927 (M^+) ; calcd $(C_{17}H_{17}O_2Cl)$ 288.0917.

2-[2-(2-Hex-5-ynyloxyethoxy)ethoxy]ethanol (23a). To a solution of triethylene glycol (6.22 g, 40 mmol) in THF (250 mL) at 0 °C was added NaH (1.58 g, 40 mmol, 60% in mineral oil) as a solid. The solution was stirred for 1 h and slowly warmed to room temperature. Subsequently, 6-chlorohexyne (4.81 g, 40 mmol) was added via syringe and the reaction was heated at reflux for 15 h. The reaction was cooled to room temperature and solvent was reduced to one half on a rotary evaporator. The residue was extracted with ether (300 mL) and water (100 mL). The organic layer was separated, dried over magnesium sulfate, filtered, and solvent evaporated. The crude oil was purified by column chromatography on silica gel with ethyl acetate-hexane 3 : 1 as the eluant to yield a clear viscous oil, $R_{\rm f} = 0.30$ (3.28 g, 36%). ¹H NMR (500 MHz, CDCl₃, δ): 3.73 (bs, 2H), 3.67–3.65 (m, 6H), 3.63-3.59 (m, 4H), 3.49 (t, J = 7.0 Hz, 2H), 2.71 (bs, 1H), 2.22 (dt, J = 2.0, 7.0 Hz, 2H), 1.95 (t, J = 2.0 Hz, 1H), 1.71 (q, J = 7.0 Hz, 2H), 1.60 (q, J = 7.0 Hz, 2H). ¹³C NMR (125 MHz, CHCl₃, δ): 84.31, 72.45, 70.72, 70.60, 70.55, 70.33, 70.02, 68.35, 61.69, 28.50, 25.04, 18.10. EI HRMS m/z: found 248.1853 (MNH₄⁺); calcd (C₁₂H₂₆NO₄) 248.1862.

Toluene-4-sulfonic acid 2-[2-(2-hex-5-ynyloxyethoxy)ethoxy] ethyl ester (23b). To an anhydrous solution of 23a (1.67 g, 7 mmol), triethylamine (1.52 mL, 11 mmol) and 4-(dimethylamino)pyridine (0.09 g, 0.7 mmol) in dichloromethane (200 mL) at 0 °C was added *p*-toluenesulfonyl chloride (1.39 g, 7 mmol) as a solid. The reaction was slowly warmed to room temperature and stirred for an additional 8 h. The crude reaction mixture was quenched with water (75 mL). The organic layer was separated and the aqueous layer extracted with fresh dichloromethane $(2 \times 20 \text{ mL})$. The combined organic layers were dried over magnesium sulfate, filtered, and reduced to dryness to yield a viscous oil. The crude oil was purified by column chromatography on silica gel with hexane-ethyl acetate 1:1 as the eluant to yield a clear and colorless oil, $R_{\rm f} = 0.50$ (2.29 g, 82%). ¹H NMR (400 MHz, CDCl₃, δ): 7.80 (d, J = 8.0 Hz, 2H), 7.34 (d, J = 8.0 Hz, 2H), 4.16 (t, J = 5.0 Hz, 2H), 3.69 (t, J = 5.0 Hz, 2H), 3.61-3.56 (m, 8H), 3.47 (t, J = 8.0 Hz,2H), 2.45 (s, 3H), 2.21 (dt, J = 2.0, 8.0 Hz, 2H), 1.94 (t, J =2.0 Hz, 1H), 1.68 (q, J = 7.0 Hz, 2H), 1.59 (q, J = 7.0 Hz, 2H). ¹³C NMR (100 MHz, CHCl₃, δ): 144.55, 132.93, 129.64, 127.82, 84.30, 70.77, 70.72, 70.65, 70.56, 70.08, 69.23, 68.70, 68.37, 28.74, 25.27, 21.74, 18.32. EI HRMS m/z: found 402.1938 (MNH₄⁺); calcd (C₁₉H₃₂NO₆S) 402.1950.

4'-Prop-2-ynyloxybiphenyl-4-ol (24a). To a 0 °C solution of 4,4'-biphenol (3.0 g, 16 mmol) in THF (100 mL) was added NaH (0.71 g, 17 mmol, 60% in mineral oil) as a solid. The resulting solution was allowed to reach room temperature while stirring over 1 h. Subsequently, propargyl bromide (2.11 g, 18 mmol) was added via syringe and the contents heated at reflux for 10 h. The crude reaction mixture was cooled to room temperature, filtered, and reduced to dryness. The crude solid was purified by column chromatography on silica gel with dichloromethane as the eluant to yield a white crystalline solid, $R_{\rm f} = 0.40$ (1.30 g, 36%). Mp 133–136 °C. ¹H NMR (300 MHz, CDCl₃, δ): 7.47 (d, J = 8.0 Hz, 2H), 7.43 (d, J = 8.0 Hz, 2H), 7.03 (d, J = 8.0 Hz, 2H), 6.88 (d, J = 8.0 Hz, 2H), 4.73 (d, J = 2.5 Hz, 2H), 2.54 (t, J = 2.5 Hz, 1H). ¹³C NMR (100 MHz, CDCl₃, δ): 156.39, 154.55, 134.18, 133.29, 127.87, 127.58, 115.50, 115.03, 78.55, 75.56, 55.92. EI HRMS m/z: found 224.0835 (M⁺); calcd (C₁₅H₁₂O₂) 224.0837.

4'-(2-Bromoethoxy)-4-prop-2-ynyloxybiphenyl (24b). To an anhydrous solution of 24a (0.572 g, 3 mmol) in THF (50 mL) at 0 °C was added NaH (0.15 g, 4 mmol, 60% in mineral oil) as a solid. The resulting solution was stirred for 1 h while warming to room temperature. Subsequently, an excess of 1,2-dibromoethane (6 mL) was added and the contents heated to reflux for 8 h. The reaction was cooled to room temperature, filtered, and solvent evaporated. The crude product was purified by column chromatography on silica gel with hexane-dichloromethane 1:3 as the eluant to yield a white crystalline solid, $R_{\rm f} = 0.60$ (0.60 g, 70%). Mp 118–119 °C. ¹H NMR (300 MHz, CDCl₃, δ): 7.49 (d, J = 8.7 Hz, 2H), 7.48 (q, J = 8.7 Hz, 2H), 7.03 (d, J = 8.7 Hz, 2H), 6.97 (d, J =8.7 Hz, 2H), 4.73 (d, J = 2.4 Hz, 2H), 4.33 (t, J = 6.0 Hz, 2H), 3.66 (t, J = 6.0 Hz, 2H), 2.54 (t, J = 2.4 Hz, 1H). ¹³C NMR (100 MHz, CDCl₃, δ): 157.05, 156.49, 134.02, 133.89, 127.76, 127.63, 115.05, 114.93, 78.54, 75.57, 67.96, 55.90, 29.23. EI HRMS m/z: found 330.0248 (M⁺); calcd (C₁₇H₁₅O₂Br) 330.0255.

Ru[5',5"-bis(4-hydroxy-2,6-dimethylphenyl)-2,2':6',2"-terpyri $dine_{12} \cdot 2PF_{6}$ (5). A degassed solution of ethylene glycol (30 mL), 3b (0.500 g, 1.1 mmol) and RuCl₂(DMSO)₄ (0.255 g, 0.5 mmol) was heated to 160 °C for 12 h to produce a deep red solution. The reaction was cooled to room temperature, poured into aqueous KPF₆ (50 mL) and filtered over Celite. The red precipitate was washed with water and dissolved into a clean flask with acetone. The resulting red solution was dried over magnesium sulfate, filtered, and the solvent evaporated. Purification by column chromatography on silica gel with CH_3CN-H_2O -aqueous KPF_6 96 : 4 : 0.04 as the eluant yielded a red solid, $R_{\rm f} = 0.40 \ (0.59 \text{ g}, 84\%)$. Mp >350 °C. ¹H NMR (300 MHz, CD₃CN, δ): 8.81 (d, J = 8.0 Hz, 4H), 8.63 (d, J =8.0 Hz, 4H, 8.43 (t, J = 8.0 Hz, 2H), 7.89 (dd, J = 2.0, 8.0 Hz, 2000 Hz)4H), 7.30 (d, J = 2.0 Hz, 4H), 7.08 (bs, 4H), 6.61 (s, 8H), 1.67 (s, 24H). ¹³C NMR (100 MHz, acetone- d_6 , δ): 157.88, 157.07, 155.93, 153.51, 141.29, 140.34, 137.47, 136.71, 126.91, 124.61, 124.19, 115.16, 20.49. MALDI HRMS m/z: found 1048.3250 (M⁺); calcd (C₆₂H₅₄N₆O₄ Ru) 1048.3250. UV-Vis (CH₃CN), $\lambda_{\text{max}}/\text{nm}$ (log ε): 201 (5.4), 272 (4.7), 312 (4.8), 478 (4.1).

Ru[8-(4-hydroxy-2,6-dimethylphenyl)-2-[5-(4-hydroxy-2,6-dimethylphenyl)pyridin-2-yl]-1,10-phenanthroline]₂**·2PF**₆ (12). A degassed solution of ethylene glycol (30 mL), 4b (0.500 g, 1.0 mmol) and RuCl₂(DMSO)₄ (0.243 g, 0.5 mmol was heated to 160 °C for 12 h to produce a deep red solution. The reaction was cooled to room temperature, poured into aqueous KPF₆ (50 mL) and filtered over Celite. The red precipitate was washed with water and dissolved into a clean flask with acetone. The resulting red solution was dried over magnesium sulfate, filtered, and the solvent removed. Purification by column chromatography on silica gel with CH₃CN–H₂O–aqueous KPF₆ 95 : 5 : 0.05 as the eluant yielded a red solid, $R_f = 0.40$ (0.61 g, 87%). Mp >350 °C. ¹H NMR (500 MHz, acetone- d_6 , δ): 9.20 (d, J = 8.5 Hz, 4H), 9.00 (d, J = 8.0 Hz, 4H), 8.94

(d, J = 8.5 Hz, 4H), 8.52 (d, J = 9.0 Hz, 4H), 8.46 (d, J = 2.0 Hz, 4H), 8.40 (s, 4H), 8.39 (s, 4H), 8.37 (d, J = 9.0 Hz, 4H), 7.98 (dd, J = 2.0, 8.0 Hz, 4H), 7.90 (d, J = 2.0 Hz, 4H), 6.49 (s, 4H), 6.46 (s, 4H), 6.42 (s, 4H), 6.39 (s, 4H), 1.57 (s, 12H), 1.56 (s, 12H), 1.34 (s, 12H), 1.32 (s, 12H). ¹³C NMR (125 MHz, acetone- d_6 , δ): 158.34*, 158.11, 156.35, 156.14, 155.41, 149.55, 147.30, 141.73, 140.92, 140.42, 138.29, 137.98, 137.90, 137.81, 137.72, 135.12, 131.76, 130.97, 129.97, 128.89, 127.65, 127.18, 125.60, 123.12, 115.44*, 115.38*, 20.59, 20.51, 20.33, 20.21. MALDI HRMS *m/z*: found 1096.3292 (M⁺); calcd (C₆₆H₅₄N₆O₄ Ru) 1096.3250. UV–Vis (CH₃CN), $\lambda_{max}/nm (\log \varepsilon)$: 201 (5.5), 239 (5.0), 296 (5.0), 343 (4.8), 499 (4.2).

Ru(5,5"-bis-(2,6-dimethyl-4-{2-[2-(2-pent-4-enyloxyethoxy)ethoxy]ethoxy]phenyl)-[2,2';6',2'']-terpyridine)₂·2PF₆ (6). To an anhydrous solution of 5 (0.222 g, 0.17 mmol) in acetonitrile (35 mL) was added anhydrous cesium carbonate (0.61 g, 1.9 mmol) and 21b (0.418 g, 1.1 mmol). The contents were heated at reflux for 10 h. The reaction was cooled to room temperature, poured into aqueous KPF₆ (50 mL) and filtered over Celite. The red oily precipitate was washed with water and transferred to a clean flask with dichloromethane. The resulting red solution was dried over magnesium sulfate, filtered, and the solvent removed. Purification by column chromatography on silica gel eluted with CH₃CN-H₂O-aqueous KPF₆ 96 : 4 : 0.04 afforded a red glassy solid, $R_{\rm f} = 0.70$ (0.23 g, 66%). Mp 49–52 °C. ¹H NMR (400 MHz, acetone- d_6 , δ): 8.94 (d, J =8.0 Hz, 4H), 8.86 (d, J = 8.0 Hz, 4), 8.37 (t, J = 8.0 Hz, 2H), 7.95 (dd, J = 2.0, 8.0 Hz, 4H), 7.64 (d, J = 2.0 Hz, 4H), 6.62 (s, 3.0 Hz, 4H), 6.62 (8H), 5.81 (m, 4H), 4.99 (dd, J = 2.0, 17.0 Hz, 4H), 4.91 (dd, J = 2.0, 10.0 Hz, 4H), 4.08 (t, J = 5.0 Hz, 8H), 3.78 (t, J =5.0 Hz, 8H), 3.65–3.57 (m, 24H), 3.52 (t, J = 5 Hz, 8H), 3.43 (t, J = 6.0 Hz, 8H), 2.14 (q, J = 7.0 Hz, 2H), 1.61 (s, 24H),1.60 (q, J = 7.0 Hz, 2H). ¹³C NMR (100 MHz, acetone- d_6, δ): 159.40, 157.17, 155.88, 153.36, 141.05, 140.29, 138.98, 137.46, 136.75, 128.01, 124.68, 124.25, 114.70, 114.28, 71.15, 71.05. 71.01, 70.68, 70.66, 70.15, 68.01, 30.93, 29.68, 20.61. MALDI HRMS m/z: found 1848.8969 (M⁺); calcd (C₁₀₆H₁₃₄N₆O₁₆Ru) 1848.8900. UV–Vis (CH₃CN), λ_{max} /nm (log ε): 202 (5.5), 234 (sh, 4.8), 272 (4.7), 312 (4.8), 478 (4.1).

 $Ru(5,5''-bis-(4-\{2-[2-(2-hex-5-ynyloxyethoxy)ethoxy]ethoxy\}-$ 2,6-dimethylphenyl)-[2,2';6',2'']-terpyridine)₂·2PF₆ (8). To an anhydrous solution of 5 (0.100 g, 0.075 mmol) in acetonitrile (15 mL) was added anhydrous cesium carbonate (0.243 g, 0.74 mmol) and 23b (0.229 g, 0.60 mmol). The contents were heated at reflux for 12 h. The reaction was cooled to room temperature, poured into aqueous KPF₆ (25 mL) and filtered over Celite. The red oily precipitate was washed with water and transferred to a clean flask with dichloromethane. The resulting red solution was dried over magnesium sulfate, filtered, and the solvent removed. Purification by column chromatography on silica gel eluted with CH₃CN-H₂O-aqueous KPF₆, 96 : 4 : 0.04 afforded a red glassy solid (0.09 g, 56%). Mp 54-58 °C. ¹H NMR (500 MHz, acetone- d_6 , δ): 8.97 (d, J = 8.0 Hz, 4H), 8.87 (d, J = 8.0 Hz, 4H), 8.39 (t, J = 8.0 Hz, 2H), 7.97 (dd, J = 2.0, 3.0 Hz, 2.0)8.0 Hz, 4H), 7.65 (d, J = 2.0 Hz, 4H), 6.63 (s, 8H), 4.09 (t, J = 4.5 Hz, 8H), 3.79 (t, J = 4.5 Hz, 8H), 3.65–3.58 (m, 24H), 3.53 (t, J = 4.5 Hz, 8H), 3.45 (t, J = 5.0 Hz, 8H), 2.31 (t, J = 2.5 Hz, 8H)4H), 2.18 (dt, J = 2.5, 5.0 Hz, 8H), 1.65–1.62 (m, 8H), 1.62 (s, 24H), 1.57–1.54 (m, 8H). ¹³C NMR (125 MHz, acetone- d_6 , δ): 159.03, 156.78, 155.51, 152.98, 140.65, 139.82, 137.03, 136,20, 127.51, 124.14, 123.71, 113.81, 84.01, 70.44, 70.32, 70.29, 70.10, 69.91, 69.32, 69.06, 67.33, 25.14, 19.74, 19.67, 17.56. MALDI HRMS m/z: found 1896.8874 (M⁺); calcd (C₁₁₀H₁₃₄N₆O₁₆Ru) 1896.8900. UV–Vis (CH₃CN), λ_{max}/nm (log ε): 201 (5.6), 273 (4.9), 312 (5.0), 478 (4.3).

Ru(5,5''-bis-{2,6-dimethyl-4-[2-(4'-prop-2-ynylbiphenyl-4-yloxy)ethoxy]phenyl}-[2,2';6',2'']-terpyridine)₂·2PF₆ (10). To an anhydrous solution of 5 (0.100 g, 0.075 mmol) in acetonitrile

(5 mL) was added anhydrous cesium carbonate (0.192 g, 0.59 mmol), 24b (0.123 g, 0.37 mmol), and triglyme (100 µL). The contents were heated at reflux for 36 h. The reaction was cooled to room temperature, poured into aqueous KPF₆ (25 mL) and filtered over Celite. The red oily precipitate was washed with water and transferred to a clean flask with dichloromethane. The resulting red solution was dried over magnesium sulfate, filtered, and the solvent removed. Purification by column chromatography on silica gel eluted with CH₃CN-H₂O-aqueous KPF₆ 96 : 4 : 0.04 afforded a red solid (0.12 g, 70%). Mp 154–155 °C. ¹H NMR (400 MHz, acetone-d₆, δ): 8.95 (d, J = 8.0 Hz, 4H), 8.70 (d, J = 8.0 Hz, 4H), 8.38 (t, J = 8.0 Hz, 2H), 7.98 (dd, J = 2.0, 8.0 Hz, 4H), 7.66 (d, J =2.0 Hz, 4H), 7.56 (d, J = 8.0 Hz, 16H), 7.07 (d, J = 8.0 Hz, 8H), 7.04 (d, J = 8.0 Hz, 8H), 6.69 (s, 8H), 4.83 (d, J = 2.4 Hz, 8H), 4.36 (bd, J = 4.5 Hz, 16H), 3.12 (t, J = 2.4 Hz, 4H), 1.63 (s, 24H). ¹³C NMR (125 MHz, acetone- d_6 , δ): 159.78, 158.99, 157.90, 157.70, 156.39, 153.86, 141.52, 140.74, 138.07, 137.17, 134.71, 134.20, 128.63, 128.46, 128.32, 125.06, 124.63, 116.11, 115.80, 114.72, 79.82, 77.05, 67.43, 67.39, 56.32, 20.62. MALDI HRMS m/z: found 2048.7409 (M⁺); calcd (C₁₃₀H₁₁₀N₆O₁₂Ru) 2048.7220. UV–Vis (CH₃CN), λ_{max}/nm (log ε): 201 (5.9), 266 (5.3), 312 (4.9), 478 (4.1).

Ru(8-(4-{2-[2-(2-hex-5-ynyloxyethoxy)ethoxy]ethoxy}-2,6dimethylphenyl)-2-[5-(4-{2-[2-(2-hex-5-ynyloxyethoxy)ethoxy]ethoxy}-2,6-dimethylphenyl)pyridin-2-yl]-[1,10]-phenanthroline)2. $2PF_6$ (13). To an anhydrous solution of 12 (0.70 g, 0.051 mmol)in acetonitrile (10 mL) was added anhydrous cesium carbonate (0.132 g, 0.40 mmol) and 23b (0.155 g, 0.40 mmol). The contents were heated at reflux for 12 h. The reaction was cooled to room temperature, poured into aqueous KPF_6 (15 mL) and filtered over Celite. The red oily precipitate was washed with water and transferred to a clean flask with dichloromethane. The resulting red solution was dried over magnesium sulfate, filtered, and the solvent removed. Purification by column chromatography on silica gel eluted with CH₃CN-H₂O-aqueous KPF₆ 96 : 4 : 0.04 afforded a red glassy solid (63 mg, 56%). Mp 58-62 °C. ¹H NMR (500 MHz, acetone- d_6 , δ): 9.22 (d, J = 8.0 Hz, 2H), 9.02 (d, J = 8.0 Hz, 2H), 8.95 (d, J = 8.0 Hz, 2H), 8.53 (d, J = 8.0 Hz, 2H, 8.49 (d, J = 2.0 Hz, 2H), 8.39 (d, J = 8.0 Hz, 2H), 8.01 (dd, J = 2.0, 8.0 Hz, 2H), 7.93 (d, J = 2.0 Hz, 2H), 7.72 (d, J = 2.0 Hz, 2H), 6.62 (s, 2H), 6.59 (s, 2H), 6.55 (s, 2H), 6.53 (s, 2H), 4.05₂ (t, J = 5.0 Hz, 4H), 4.05 (t, J = 5.0 Hz, 4H), 3.76 (t, J = 5.0 Hz, 4H), 3.75 (t, J = 5.0 Hz, 4H), 3.63–3.51 (m, 32H), 3.45 (t, J = 6.0 Hz, 8H), 2.31 (t, J = 2.0 Hz, 4H), 2.18 (dt, J = 2.0, 7.0 Hz, 8H), 1.64–1.62 (m, 8H), 1.63 (s, 12H), 1.57–1.54 (m, 8H), 1.40 (s, 6H), 1.37 (s, 6H). ¹³C NMR (100 MHz, acetone-d₆, δ): 159.42*, 157.76, 155.75, 155.67, 154.84, 149.19, 146.84, 141.10, 140.49, 139.79, 137.88, 137.64, 137.57, 137.46, 137.37, 134.79, 131.39, 130.64, 129.65, 128.59, 128.44, 127.97, 125.33, 122.84, 114.31*, 114.24*, 84.71*, 71.17*, 71.06*, 71.02*, 70.84*, 70.66*, 70.04*, 69.84*, 68.05*, 29.44*, 26.04*, 20.75, 20.66, 20.49, 20.36, 18.48*. MALDI HRMS m/z: found 1944.8859 (M⁺); calcd ($C_{114}H_{134}N_6O_{16}Ru$) 1944.8894. UV–Vis (CH₃CN), $\lambda_{\text{max}}/\text{nm}$ (log ε): 202 (5.4), 273 (5.2), 349 (4.7), 499 (4.0).

Ru(terpy-catenane)·2PF₆ (7a). To a solution of RuCl₂-(PCy₃)₂CH₂Ph (3 mg, 3.5 µmol) in degassed dichloromethane (50 mL) at room temperature was added a solution of **6** (0.15 g, 0.07 mmol) in dichloromethane (10 mL) *via* syringe pump over 12 h. After stirring for an additional 48 h, the red solution was reduced to dryness and purified by column chromatography on silica gel with CH₃CN–H₂O–aqueous KPF₆ 95 : 5 : 0.05 as the eluant to yield a red glassy solid (0.04 g, 27%) as a mixture of *cis/trans* isomers. This mixture was dissolved in dichloromethane–ethanol (1 : 1, 4 mL) and to it was added Pd/C (10 mol%). The suspension was stirred for 20 h under a hydrogen atmosphere. The slurry was filtered and reduced to dryness to yield a red glassy solid (0.04 g, 98%). Mp 118–120 °C. ¹H NMR (500 MHz, acetone-*d*₆, δ): 8.94 (d, *J* = 8.0 Hz, 4H), 8.86 (d, J = 8.0 Hz, 4H), 8.32 (t, J = 8.0 Hz, 2H), 8.02 (dd, J = 2.0, 8.0 Hz, 4H), 7.61 (d, J = 2.0 Hz, 4H), 6.64 (s, 8H), 4.13 (bt, J = 5.0 Hz, 8H), 3.77 (bt, J = 5.0 Hz, 8H), 3.64–3.56 (m, 32H), 3.49 (t, J = 7.0 Hz, 8H), 1.63 (bs, 8H), 1.60 (s, 24H), 1.44 (bs, 16H). ¹³C NMR (125 MHz, acetone- d_6 , δ): 160.02, 157.58, 156.43, 154.01, 141.69, 140.49, 137.87, 136.85, 128.30, 125.08, 124.58, 114.77, 71.93, 71.52, 71.29, 71.24, 70.88, 70.33, 68.48, 30.75, 30.56, 27.11, 20.59. MALDI HRMS *m*/*z*: found 1796.8537 (M⁺); calcd (C₁₀₂H₁₃₀N₆O₁₆Ru) 1796.8587. UV–Vis (CH₃CN), λ_{max}/nm (log ε): 202 (5.5), 235 (sh, 5.0), 272 (4.9), 312 (5.0), 479 (4.2).

Ru(terpy-macrocycle).2PF₆ (7b). Prepared simultaneously and hydrogenated in an identical manner to compound 7a (0.068 g, 47%) cis/trans isomers. Hydrogenation yielded a red glassy solid, 7b (0.065 g, 97%). Mp 132-134 °C. ¹H NMR (500 MHz, acetone- d_6 , δ): 8.96 (d, J = 8.0 Hz, 4H), 8.88 (d, J = 8.0 Hz, 4H), 8.40 (t, J = 8.0 Hz, 2H), 7.96 (dd, J = 2.0, 8.0 Hz, 4H), 7.67 (d, J = 2.0 Hz, 4H), 6.66 (s, 4H), 6.64 (s, 4H), 4.13 (t, J = 5.0 Hz, 8H), 3.77 (t, J = 5.0 Hz, 8H), 3.62 (t, J =5.0 Hz, 8H), 3.57-3.53 (m, 16H), 3.45 (t, J = 5.0 Hz, 8H), 3.33 (t, J = 7.0 Hz, 8H), 1.73 (s, 12H), 1.51 (s, 12H), 1.43 (q, J = 7.0 Hz, 8H), 1.19 (bs, 16H). ¹³C NMR (125 MHz, acetone- d_6, δ): 160.13, 157.73, 156.45, 153.91, 141.61, 140.83, 137.89, 137.70, 137.26, 128.41, 125.07, 124.67, 115.16, 115.06, 71.81, 71.61, 71.39, 71.28, 70.83, 70.35, 68.54, 30.65, 30.40, 26.99, 20.82, 20.62. MALDI HRMS m/z: found 1796.8624 (M⁺); calcd (C₁₀₂H₁₃₀N₆O₁₆Ru) 1796.8581. UV–Vis (CH₃CN), λ_{max}/nm (log ε): 203 (5.4), 236 (sh, 4.9), 273 (4.8), 311 (4.9), 479 (4.1).

Standard reaction conditions for copper-mediated coupling (compounds 9, 11, and 14a–b). To enough acetonitrile to generate a 1.0 mM solution of the corresponding pre-catenane Ru(II) complex (1.0 mol eq.) was added copper(II) acetate monohydrate (20 mol eq.), and the mixture heated at reflux for 12–32 h, or until no starting material was visible by TLC (SiO₂, CH₃CN–H₂O–aqueous KPF₆ 95 : 5 : 0.05). The reactions were cooled and reduced to dryness. The residues were suspended in dichloromethane, filtered and the solvent evaporated to yield the crude red compounds. The crude mixtures were purified by column chromatography on silica gel with 100% CH₃CN \rightarrow CH₃CN–H₂O–aqueous KPF₆ 95 : 5 : 0.05 as the eluant, except for compounds 11a–b which required centrifugal chromatography on silica gel with 100% CH₂Cl₂ \rightarrow 2% MeOH as the eluant.

Ru(terpy-acteylcatenane)-2PF₆ (9a). A red glassy solid (0.012 g, 25%). Mp 65–70 °C. ¹H NMR (500 MHz, acetoned₆, δ): 8.95 (d, J = 8.0 Hz, 4H), 8.87 (d, J = 8.0 Hz, 4H), 8.36 (t, J = 8.0 Hz, 2H), 8.01 (dd, J = 2.0, 8.0 Hz, 4H), 7.62 (d, J = 2.0 Hz, 4H), 6.34 (s, 8H), 4.12 (t, J = 5.0 Hz, 8H), 3.78 (t, J = 5.0 Hz, 8H), 3.64–3.60 (m, 24H), 3.55 (t, J = 5.0 Hz, 8H), 3.50 (t, J = 6.0 Hz, 8H), 2.36 (t, J = 6.0 Hz, 8H), 1.69–1.63 (m, 16H), 1.62 (s, 24H). ¹³C NMR (125 MHz, acetone-d₆, δ): 159.11, 156.71, 155.51, 153.07, 140.75, 139.72, 136.98, 136.21, 127.43, 124.16, 123.69, 113.85, 77.50, 70.60, 70.40, 70.31, 70.07, 69.99, 69.36, 67.56, 65.54, 25.06, 19.73, 19.69, 18.35. MALDI HRMS *m/z*: found 1892.8710 (M⁺); calcd (C₁₁₀H₁₃₀N₆O₁₆Ru) 1892.8581. UV–Vis (CH₃CN), λ_{max}/nm (log ε): 201 (5.2), 235 (sh, 4.7), 272 (4.6), 312 (4.7), 480 (3.9).

Ru(terpy-acetylmacrocycle)·2PF₆ (9b). A red glassy solid (0.017 g, 36%). Mp 79–84 °C. ¹H NMR (500 MHz, acetoned₆, δ): 8.98 (d, J = 8.0 Hz, 4H), 8.88 (d, J = 8.0 Hz, 4H), 8.42 (t, J = 8.0 Hz, 2H), 7.98 (dd, J = 2.0, 8.0 Hz, 4H), 7.65 (d, J = 2.0 Hz, 4H), 6.66 (s, 4H), 6.65 (s, 4H), 4.12 (t, J = 5.0 Hz, 8H), 3.79 (t, J = 5.0 Hz, 8H), 3.64 (t, J = 5.0 Hz, 8H), 3.58, (t, J = 5.0 Hz, 8H), 3.56 (t, J = 5.0 Hz, 8H), 3.48 (t, J = 5.0 Hz, 8H), 3.57 (t, J = 7.0 Hz, 8H), 2.15 (t, J = 7.0 Hz, 8H), 1.69 (s, 12H), 1.59 (s, 12H), 1.52 (q, J = 7.0 Hz, 8H), 1.44 (q, J = 7.0 Hz, 8H), 153.95, 141.60, 140.85, 137.92, 137.86, 137.31, 128.37, 125.12, 124.68, 115.02, 114.88, 78.20, 71.59, 71.44, 71.32, 70.89, 70.86, 70.29, 68.47, 66.28, 29.65, 25.90, 20.81, 20.64, 19.18. MALDI HRMS *m/z*: found 1892.8663 (M⁺); calcd (C₁₁₀H₁₃₀N₆O₁₆Ru) 1892.8581. UV–Vis (CH₃CN), λ_{max} /nm (log ε): 202 (5.2), 236 (4.8), 272 (4.9), 311 (4.8), 478 (4.0).

Ru(terpy-biphencatenane)·2PF₆ (11a). A red solid (0.021 g, 29%). Mp > 340 °C. ¹H NMR (500 MHz, acetone- d_6 , δ): 8.78 (d, J = 8.0 Hz, 4H), 8.76 (d, J = 8.5 Hz, 4H), 8.19 (t, J = 8.0 Hz, 2H), 7.93 (dd, J = 2.0, 8.5 Hz, 4H), 7.60 (d, J = 9.0 Hz, 8H), 7.58 (d, J = 2.0 Hz, 4H), 7.57 (d, J = 9.0 Hz, 8H), 7.12 (d, J = 9.0 Hz, 8H), 7.03 (d, J = 9.0 Hz, 8H), 6.64 (s, 8H), 5.01 (s, 8H), 4.41 (bd, J = 8.0 Hz, 16H), 1.55 (s, 24H). ¹³C NMR (125 MHz, acetone- d_6 , δ): 159.34, 158.58, 157.69, 157.05, 156.24, 153.84, 141.50, 140.50, 137.94, 135.05, 134.20, 128.54, 128.50, 128.47, 124.95, 124.55, 116.22, 116.17, 114.72, 106.00, 76.50, 71.12, 66.90, 66.55, 56.72, 20.53. MALDI HRMS m/z: found 2044.6826 (M⁺); calcd (C₁₃₀H₁₀₆N₆O₁₂Ru) 2044.6907. UV–Vis (CH₃CN), λ_{max} /nm (log ε): 202 (5.6), 268 (5.2), 314 (4.8), 478 (4.0).

Ru(terpy-biphenmacrocyle)·2PF₆ (11b). A red solid (0.021 g, 21%). Mp > 340 °C. ¹H NMR (500 MHz, acetone- d_6 , δ): 8.96 (d, J = 8.0 Hz, 4H), 8.85 (d, J = 8.0 Hz, 4H), 8.40 (t, J = 8.0 Hz, 2H), 7.93 (dd, J = 2.0, 8.5 Hz, 4H), 7.65 (d, J = 2.0 Hz, 4H), 7.52 (d, J = 8.5 Hz, 8H), 7.50 (d, J = 8.5 Hz, 8H), 7.04 (d, J = 8.5 Hz, 8H), 7.00 (d, J = 8.5 Hz, 8H), 4.98 (s, 8H), 4.41 (bd, 16H), 1.73, (s, 12H), 1.45 (s, 12H). ¹³C NMR (125 MHz, acetone- d_6 , δ): 159.70, 158.87, 157.72, 157.64, 156.48, 153.89, 141.46, 140.85, 138.01, 137.89, 137.19, 135.03, 134.33, 128.68, 128.44, 128.40, 125.04, 124.64, 116.31, 116.18, 115.69, 115.03, 76.54, 70.99, 67.54, 65.02, 56.58, 20.78, 20.62. MALDI HRMS *m/z*: found 2044.7012 (M⁺); calcd (C₁₃₀H₁₀₆N₆O₁₂Ru) 2044.6907. UV–Vis (CH₃CN), λ_{max}/nm (log ε): 202 (5.5), 266 (5.1), 311 (4.7), 479 (4.0).

Ru(pherpy-catenane) \cdot 2PF₆ (14a). A red glassy solid (0.012 g, 20%). Mp 122–128 °C. ¹H NMR (500 MHz, acetone-*d*₆, δ): 9.12 (d, J = 9.0 Hz, 2H), 9.00 (d, J = 8.0 Hz, 2H), 8.93 (d, J =9.0 Hz, 2H), 8.53 (d, J = 9.0 Hz, 2H), 8.53 (d, J = 2.0 Hz, 2H), 8.41 (d, J = 9.0 Hz, 2H), 8.04 (dd, J = 2.0, 8.0 Hz, 2H), 7.89 (d, J = 2.0 Hz, 2H), 7.69 (d, J = 2.0 Hz, 2H), 6.62 (s, 2H), 6.60 (s, 2H), 6.55 (s, 2H), 6.52 (s, 2H), 4.08 (bt, J = 6.0 Hz, 8H), 3.74 (bt, J = 6.0 Hz, 8H), 3.61–3.59 (m, 24H), 3.55 (bt, J = 6.0 Hz, 8H), 3.49 (bt, J = 6.0 Hz, 8H), 2.39 (bt, J =6.0 Hz, 8H), 1.70-1.664 (m, 16H), 1.62 (s, 6H), 1.61 (s, 6H), 1.39 (s, 6H), 1.36 (s, 6H). ¹³C NMR (125 MHz, acetone- d_6 , δ): 160.00, 159.98, 158.15, 156.36, 156.13, 155.46, 149.63, 147.31, 141.65, 140.74, 140.32, 138.11, 138.01, 137.95, 137.84, 137.76, 135.05, 131.82, 130.98, 130.14, 128.87, 128.71, 128.24, 125.67, 123.14, 114.77*, 114.71*, 78.53*, 71.50*, 71.31*, 71.22*, 70.99*, 70.91*, 70.26*, 68.49*, 66.56*, 29.72*, 26.01*, 20.74, 20.65, 20.48, 20.35, 19.32*. MALDI HRMS m/z: found 1940.8534 (M⁺); calcd (C₁₁₄H₁₃₀N₆O₁₆Ru) 1940.8581. UV–Vis (CH₃CN), $\lambda_{\rm max}/{\rm nm}$ (log ε): 202 (5.3), 236 (4.9), 297 (4.8), 343 (4.6), 501 (4.0).

5-5'' - Bis{4-[2-(4' - allyloxybiphenyl-4 - yloxy)ethoxy]-2,6dimethylphenyl}-[2,2':6'2"]-terpyridine (15). An anhydrous solution of **3b** (0.197 g, 0.42 mmol), **22b** (0.330 g, 1.14 mmol), and cesium carbonate (0.689 g, 2.11 mmol) in DMF (10 mL) was heated at 80 °C for 12 h with an oil bath. The crude reaction mixture was quenched with water (30 mL) and extracted with chloroform (30 mL). The organic layer was separated, washed with water $(3 \times 20 \text{ mL})$, dried over magnesium sulfate, filtered and reduced to dryness. Purification by column chromatography on alumina with dichloromethane-hexane 3:1 as the eluant afforded a white solid (0.357 g, 88%). Mp 188-189 °C. ¹H NMR $(500 \text{ MHz}, \text{CDCl}_3, \delta)$: 8.73 (d, J = 8.0 Hz, 2H), 8.53 (d, J =2.0 Hz, 2H), 8.52 (d, J = 8.0 Hz, 2H), 8.02 (t, J = 8.0 Hz, 1H), 7.69 (dd, J = 2.0, 8.0 Hz, 2H), 7.50 (d, J = 9.0 Hz, 4H), 7.48 (d, J = 9.0 Hz, 4H), 7.03 (d, J = 9.0 Hz, 4H), 6.98 (d, J = 9.0 Hz, 4H), 6.80 (s, 4H), 6.09 (m, 2H), 5.44 (dd, J = 2.0, 18.0 Hz, 2H), 5.31, (dd, J = 2.0, 12.0 Hz, 2H), 4.58 (d, J = 6.0 Hz, 4H), 4.39 (bs, 8H), 2.10 (s, 12H). ¹³C NMR (125 MHz, CDCl₃, δ): 158.08, 157.83, 157.80, 155.40, 154.67, 150.03, 138.35, 138.09, 138.01, 136.69, 133.91, 133.56, 133.34, 130.88, 127.79, 127.75, 125.55, 120.90, 117.74, 115.03, 115.00, 113.79, 68.88, 66.61, 66.42, 21.19. MALDI HRMS m/z: found 978.4525 (M⁺); calcd (C₆₅H₆₀N₃O₆) 978.4477.

8-{4-[2-(4'-Allyloxybiphenyl-4-yloxy)ethoxy]-2,6-dimethylphenyl}-2-(5-{4-[2-(4'-allyloxybiphenyl-4-yloxy)ethoxy]-2,6dimethylphenyl}pyridin-2-yl)-[1,10]-phenanthroline (18). An anhydrous solution of 4b (0.256 g, 0.52 mmol), 22b (0.447 g, 1.55 mmol), and cesium carbonate (0.840 g, 2.58 mmol) in DMF (13 mL) was heated at 80 °C for 12 h with an oil bath. The crude reaction mixture was quenched with water (30 mL) and extracted with chloroform (30 mL). The organic layer was separated, washed with water (3 \times 20 mL), dried over magnesium sulfate, filtered and reduced to dryness. Purification by column chromatography on alumina with dichloromethanehexane 3:1 as the eluant afforded a white solid (0.473 g, 92%). Mp 203–204 °C. ¹H NMR (400 MHz, CDCl₃, δ): 9.07 (d, J =8.0 Hz, 1H), 9.08 (d, J = 2.0 Hz, 1H), 8.86, (d, J = 8.0 Hz, 1H), 8.56 (d, J = 2.0 Hz, 1H), 8.42 (d, J = 8.0 Hz, 1H), 8.07 (d, J = 2.0 Hz, 1H), 7.88 (d, J = 8.0 Hz, 1H), 7.82 (d, J =8.0 Hz, 1H), 7.75 (dd, J = 2.0, 8.0 Hz, 1H), 7.49 (d, J = 8.0 Hz, 4H), 7.47 (d, J = 8.0 Hz, 4H), 7.02 (d, J = 8.0 Hz, 4H), 6.97 (d, J = 8.0 Hz, 4H), 6.82 (s, 2H), 6.79 (s, 2H), 6.06 (m, 2H),5.43 (dd, J = 2.0, 18.0 Hz, 2H), 5.28, (dd, J = 2.0, 12.0 Hz, 2H), 4.55 (d, J = 6.0 Hz, 4H), 4.38 (s, 4H), 4.37 (s, 4H), 2.10 (s, 6H), 2.07 (s, 6H). ¹³C NMR (125 MHz, CDCl₃, δ): 158.18*, 158.00*, 157.75*, 156.39, 154.67, 151.99, 149.83, 145.74, 144.93, 138.50, 138.06, 137.91, 137.06, 137.01, 136.53, 135.93, 133.81, 133.78, 133.47, 133.45, 133.28*, 130.84, 130.61, 128.81, 128.66, 127.71*, 127.65*, 126.78, 126.74, 122.54, 120.70, 117.62* 114.97*, 114.94*, 113.79, 113.73, 68.78*, 66.54*, 66.38, 66.35, 21.20, 21.12. MALDI HRMS m/z: found 1002.4458 (M⁺); calcd (C₆₇H₆₀N₃O₆) 1002.4477.

Fe(5-5"-bis{4-[2-(4'-Allyloxybiphenyl-4-yloxy)ethoxy]-2,6dimethylphenyl $\left\{-[2,2':6'2''] - \text{terpyridine}\right)_2 \cdot 2PF_6$ (16). To a suspension of **15** (0.174 g, 0.18 mmol) in acetone (20 mL), at 55 °C, was added a solution of $FeSO_4{\cdot}7H_2O$ (0.049 g, 0.18 mmol) in water (12 mL) dropwise to generate an intense purple solution, which was heated at reflux for 1 h. The crude reaction was quenched with aqueous KPF_6 to generate a purple precipitate, which was filtered and washed with water. The precipitate was dissolved in dichloromethane, dried over magnesium sulfate, filtered, and reduced to dryness to yield a purple solid (0.203 g, 99%). Mp 158–159 °C. ¹H NMR (500 MHz, acetone- d_6 , δ): 9.14 (d, J = 8.0 Hz, 4H), 8.87 (d, J = 8.0 Hz, 4H), 8.67 (t, J =8.0 Hz, 2H), 7.97 (dd, J = 2.0, 8.0 Hz, 4H), 7.56 (d, J = 8.0 Hz, 8H), 7.54 (d, J = 8.0 Hz, 8H), 7.35 (d, J = 2.0 Hz, 4H), 7.04 (d, J = 8.0 Hz, 8H), 7.03 (d, J = 8.0 Hz, 8H), 6.83 (s, 8H),6.10 (m, 4H), 5.43 (dd, J = 2.0, 17.0 Hz, 4H), 5.27 (dd, J =2.0, 11.0 Hz, 4H), 4.62 (bd, J = 5.0 Hz, 8H), 4.37 (bd, J =8.0 Hz, 16H), 1.56 (s, 24H). ¹³C NMR (125 MHz, CD₂Cl₂, δ): 160.42, 159.43, 158.40, 158.29, 156.20, 153.91, 1491.78, 139.90, 137.52, 134.16, 134.14, 134.13, 134.01, 133.72, 128.10, 128.03, 127.56, 124.08, 117.67, 115.50, 115.38, 114.75, 69.28, 67.14, 67.08, 20.74. MALDI HRMS m/z: found 2010.8226 (M⁺); calcd (C₁₃₀H₁₁₈N₆O₁₂Fe) 2010.8152. UV-Vis (CH₃CN), λ_{max}/nm (log ε): 202 (5.6), 269 (5.1), 333 (4.7), 556 (3.9).

Fe(8-{4-[2-(4'-allyloxybiphenyl-4-yloxy)ethoxy]-2,6-dimethylphenyl}-2-(5-{4-[2-(4'-allyloxybiphenyl-4-yloxy)ethoxy]-2,6dimethyl-phenyl}pyridin-2-yl)-[1,10]-phenanthroline)₂·2PF₆ (19). To a suspension of 18 (0.218 g, 0.22 mmol) in acetone (20 mL), at 55 °C, was added a solution of FeSO₄·7H₂O (0.060 g, 0.22 mmol) in water (15 mL) dropwise to generate an intense purple solution, which was heated at reflux for 1 h. The crude reaction was quenched with aqueous KPF₆ to generate a purple precipitate, which was filtered and washed with water. The precipitate was dissolved in dichloromethane, dried over magnesium sulfate, filtered, and reduced to dryness to yield a purple solid (0.233 g, 91%). Mp 160-163 °C. ¹H NMR $(500 \text{ MHz}, \text{ acetone-} d_6, \delta)$: 9.45 (d, J = 8.8 Hz, 2H), 9.23 (d, J =8.8 Hz, 2H), 9.03 (d, J = 8.4 Hz, 2H), 8.64 (d, J = 8.8 Hz, 2H), 8.44 (d, J = 2.0 Hz, 2H), 8.41 (d, J = 8.8 Hz, 2H), 7.97 (dd, J = 2.0, 8.4 Hz, 2H), 7.56 (d, J = 10.0 Hz, 8H), 7.55 (d, J = 2.0 Hz, 2H), 7.54 (d, J = 10.0 Hz, 8H), 7.45 (d, J = 2.0 Hz, 2H), 7.03 (d, J = 10.0 Hz, 8H), 7.01 (d, J = 10.0 Hz, 8H), 6.66 (s, 2H),6.64 (s, 2H), 6.58 (s, 2H), 6.56 (s, 2H), 6.09 (m, 4H), 5.45 (d, J = 17.2 Hz, 4H), 5.27 (d, J = 10.8 Hz, 4H), 4.62 (m, 8H), 4.35 (bs, 8H), 4.32 (bs, 8H), 1.56 (s, 6H), 1.54 (s, 6H), 1.26 (s, 12H). ¹³C NMR (125 MHz, CD_2Cl_2 , δ): 161.92*, 159.70*, 158.89*, 158.84, 158.06, 158.00, 156.41, 151.79, 148.68, 141.76, 141.29, 140.41, 138.80, 138.04, 137.91, 137.87, 137.73, 137.12, 134.80*, 134.32*, 134.06*, 130.80, 130.67, 130.03, 129.02, 128.56, 128.36*, 128.33, 128.31*, 125.22, 123.68, 117.41*, 115.93*, 115.80*, 114.70*, 114.63*, 69.32*, 67.43*, 67.36*, 20.66, 20.59, 20.31, 20.19. MALDI HRMS m/z: found 2058.8076 (M⁺); calcd (C₁₃₄H₁₁₈N₆O₁₂Fe) 2058.8152. UV–Vis (CH₃CN), λ_{max}/nm $(\log \varepsilon)$: 203(5.7), 267 (5.2), 357 (4.7), 579 (4.0).

Fe(terpycat)·2PF₆ (17). To a solution of RuCl₂(PCy₃)₂CH₂Ph (8 mg, 0.20 mmol) in dichloromethane (20 mL) was added a solution of 16 (0.115 g, 0.05 mol) in dichloromethane (20 mL) via syringe pump over 12 h. The resulting purple solution was stirred for an additional 60 h. The crude reaction mixture was reduced to dryness and purified by column chromatography on silica gel with CH₃CN-H₂O-aqueous KPF₆ 96 : 4 : 0.04 as the eluant. Subsequent centrifugal chromatography with dichloromethane-methanol 100 : 3 as the eluant afforded a purple solid (0.033 g, 30%) as a mixture of cis/trans isomers. This mixture (0.022 g, 0.010 mmol) was dissolved in dichloromethane-ethanol 1 : 1 (6 mL), and to it was added Pt/alumina (5 mol%). The resulting solution was stirred under a hydrogen atmosphere for 36 h. The crude mixture was filtered, and the solution reduced to dryness to afford a purple solid (0.020 g, 91%). Mp 218-223 °C. 1H NMR (500 MHz, acetone- d_6 , δ): 8.91 (d, J = 8.0 Hz, 4H), 8.73 (d, J = 8.0 Hz, 4H), 8.42 (t, J = 8.0 Hz, 2H), 7.87 (dd, J = 2.0, 8.0 Hz, 4H), 7.56 (d, J = 8.8 Hz, 8H), 7.55 (d, J = 8.8 Hz, 8H), 7.24 (d, J = 2.0 Hz, 4H), 7.06 (d, J = 8.8 Hz, 8H), 7.03 (d, J = 8.8 Hz, 8H), 6.61 (s, 8H), 4.44 (d, J = 2.8 Hz, 8H), 4.37 (d, J = 2.8 Hz, 8H), 4.21 (bs, 8H), 3.06 (bs, 8H), 1.46 (s, 24H). ¹³C NMR (125 MHz, acetone- d_6 , δ): 161.11, 159.26, 159.22, 158.28, 157.54, 154.59, 141.47, 141.35, 139.33, 137.87, 134.40, 133.83, 128.62, 128.34, 128.31, 124.58, 124.30, 116.24, 115.87, 114.67, 67.99, 66.67, 66.34, 26.26, 20.45. MALDI HRMS m/z: found 1958.7705 (M⁺); calcd ($C_{126}H_{114}N_6O_{12}Fe$) 1958.7839. UV–Vis (CH₃CN), $\lambda_{\text{max}}/\text{nm}$ (log ε): 201 (5.5), 272 (5.0), 334 (4.5), 554 (3.8).

Terpy catenane (1). To a solution of **17** (0.020 g, 0.009 mmol) in acetonitrile-water 1 : 1 (12 mL) was added 10% sodium hydroxide (1 mL) followed by a slow addition of hydrogen peroxide (35% in water) until no purple color remained. The resulting solution was extracted with chloroform (20 mL). The organic layer was removed, washed with water (3 \times 15 mL), dried over magnesium sulfate, filtered and reduced to dryness. Purification by column chromatography on alumina with chloroform-hexane 3 : 1 as the eluant afforded a white solid (0.010 g, 59%). ¹H NMR (500 MHz, CDCl₃, δ): 8.60 (d, J = 8.0 Hz, 4H), 8.50 (d, J = 2.0 Hz, 4H), 8.39 (d, J =8.0 Hz, 4H), 7.93, (t, J = 8.0 Hz, 2H), 7.32 (dd, J = 2.0, 8.0 Hz, 4H), 7.28 (d, J = 9.0 Hz, 8H), 7.27 (d, J = 9.0 Hz, 8H), 6.82 (d, J = 9.0 Hz, 8H), 6.80 (d, J = 9.0 Hz, 8H), 6.74 (s, 8H),4.43 (t, J = 4.0 Hz, 8H), 4.27 (t, J = 4.0 Hz, 8H), 3.90 (t, J = 6.0 Hz, 8H), 1.95 (s, 24H), 1.87 (t, J = 6.0 Hz, 8H). ¹³C NMR (125 MHz, CDCl₃, δ): 157.94, 157.23, 157.21, 155.07, 154.67, 149.46, 138.57, 138.14, 137.83, 136.67, 133.81, 133.10, 131.01, 127.59, 127.57, 121.27, 120.37, 115.32, 114.73, 114.57, 66.82, 66.23, 65.63, 25.23, 21.12. MALDI HRMS m/z: found 1903.8420 (M⁺); calcd (C₁₂₆H₁₁₄N₆O₁₂) 1903.8568.

Fe(pherpycat)·2PF₆ (20). To a solution of 19 (0.33 g, 0.14 mmol) in dichloromethane (200 mL) was added a solution of RuCl₂(PCy₃)₂CH₂Ph (12 mg, 0.01 mmol) in dichloromethane (20 mL) via cannula. The purple solution was stirred for 72 h. The crude reaction mixture was reduced to dryness and purified by column chromatography on silica gel with CH₃CN-H₂Oaqueous KPF_6 97 : 3 : 0.03 as the eluant. Subsequent centrifugal chromatography with dichloromethane-methanol 100:3 as the eluant afforded a purple solid (0.063 g, 20%) as a mixture of cis/trans isomers. This mixture (0.023 g, 0.010 mmol) was dissolved in a solution of dichloromethane-ethanol 1:1(6mL) and to it was added Pt/alumina (5 mol%). The resulting solution was stirred under a hydrogen atmosphere for 12 h. The crude mixture was filtered and the solution reduced to dryness to afford a purple solid (0.022 g, 96%). Mp 224-230 °C. ¹H NMR (500 MHz, $CDCl_3, \delta$): 9.20 (d, J = 9.0 Hz, 2H), 8.98 (d, J = 9.0 Hz, 2H), 8.88 (d, J = 8.5 Hz, 2H), 8.45 (d, J = 9.5 Hz, 2H), 8.35 (d, J = 2 Hz, 2H), 8.29 (d, J = 9.5 Hz, 2H), 7.88 (dd, J = 2.0, 8.5 Hz, 2H), 7.60 (d, J = 8.5 Hz, 8H), 7.58 (d, J = 8.5 Hz, 8H), 7.44 (d, J = 2.0 Hz, 2H), 7.36 (d, J = 2.0 Hz, 2H), 7.10 (d, J = 8.5 Hz, 8H), 7.01 (d, J = 8.5 Hz, 8H), 6.59 (s, 2H), 6.57 (s, 2H), 6.51 (s, 2H), 6.49 (s, 2H), 4.4 (bt, J = 4.0 Hz, 8H), 4.44 (bt, J = 4.0 Hz, 8H), 4.23 (bt, J = 4.0 Hz, 8H), 2.09 (bt, J = 4.0 Hz, 8H), 1.45 (s, 6H), 1.44 (s, 6H), 1.15 (s, 12H). ¹³C NMR (125 MHz, CDCl₃, δ): 161.83*, 159.32*, 159.14, 158.26*, 158.04, 157.94, 156.36, 151.67, 148.68, 141.59, 141.25, 140.33, 138.67, 137.98, 137.86, 137.81, 137.68, 136.96, 134.38*, 13384*, 130.64, 129.81, 128.94, 128.50, 128.36*, 128.30, 128.28*, 126.36, 125.09, 123.60, 116.22*, 115.91* 114.61*, 114.58*, 67.95*, 66.63*, 66.24*, 26.24*, 20.57, 20.50, 20.23, 20.12. MALDI HRMS m/z: found 2006.7866 (M+); calcd $(C_{130}H_{114}N_6O_{12}Fe)$ 2006.7839. UV–Vis (CH₃CN), λ_{max}/nm (log ε): 202 (5.6), 260 (5.2), 356 (4.5), 579 (3.9).

Pherpy catenane (2). To a solution of $20 (0.022 \text{ g}, 10 \mu \text{mol})$ in acetonitrile-water 1:1 (6 mL) was added 10% sodium hydroxide (1 mL) followed by a slow addition of hydrogen peroxide (35% in water) until no purple color remained. The resulting solution was extracted with dichloromethane (15 mL). The organic layer was removed, washed with water $(3 \times 15 \text{ mL})$, dried over magnesium sulfate, filtered and reduced to dryness. Purification by column chromatography on silica with 2-3% MeOH in chloroform as the eluant afforded a white solid (5 mg, 27%). ¹H NMR (500 MHz, $CDCl_3, \delta$: 8.98 (d, J = 8.0 Hz, 2H), 8.83 (d, J = 2.0 Hz, 2H), 8.77 (d, J = 8.5 Hz, 2H), 8.54 (d, J = 2.0 Hz, 2H), 8.40 (d, J = 8.0 Hz, 2H), 8.07 (d, J = 2.0 Hz, 2H), 7.86 (d, J = 9.0 Hz, 2H), 7.80 (d, J = 9.0 Hz, 2H), 7.41 (dd, J = 2.0, 8.5 Hz, 2H), 7.30–7.25 (m, 8H), 8.86 (d, J = 8.5 Hz, 4H), 6.79 (d, J = 8.5 Hz, 4H), 6.77 (s, 4H), 6.73 (s, 4H), 4.44 (bt, J = 5.0 Hz, 8H), 4.31 (bs, 8H), 3.88 (bt, J = 5.0 Hz, 8H), 1.97 (s, 6H), 1.97 (s, 6H), 1.96 (s, 6H), 1.94 (s, 6H), 1.85 (bs, 8H). MALDI HRMS m/z: found 1951.8663 (MH⁺); calcd (C₁₃₀H₁₁₅N₆O₁₂) 1951.8568.

References and notes

- 1 (a) G. Schill, Catenanes, Rotaxanes, and Knots. 1971, New York, Academic Press Inc.; (b) J. S. Siegel, Science, 2004, 304, 1256–59.
- 2 (a) K. S. Chichak, S. J. Cantrill, A. R. Pease, S.-H. Chiu, G. W. V. Cave, J. L. Atwood and J. F. Stoddart, *Science*, 2004, **304**, 1308–1312; (b) J. P. Sauvage and C. Dietrich-Buchecker, *Molecular Catenanes, Rotaxanes and Knots: A Journey Through the World of Molecular Topology*, Wiley-VCH, Weinheim, 1999.
- 3 (a) A. H. Flood, R. J. A. Ramirez, W.-Q. Deng, R. P. Muller, W. A. Goddard, III and J. F. Stoddart, Aust. J. Chem., 2004, 57(4), 301–322; (b) B. Korybut-Daszkeiwicz, A. Wieckowska, R. Ilewicz, S. Domagala and K. Wozniak, Angew. Chem., Int. Ed., 2004, 43(13), 1668–1672; (c) P. Mobian, J.-M. Kern and J.-P. Sauvage, Angew. Chem., Int. Ed., 2004, 43(18), 2392–2395; (d) M. Cavallini, F. Biscarini, S. Leon, F. Zerbetto, G. Bottari and D. A. Leigh, Science, 2003, 299(5606), 531; (e) T. Da Ros, D. M. Guldi, A. F. Morales, D. A. Leigh, M. Prato and R. Turco, Org. Lett., 2003, 5(5),

689–691; (f) V. Balzani, A. Credi, F. M. Raymo and J. F. Stoddart, Angew. Chem., Int. Ed., 2000, **39**(19), 3348–3391; (g) V. Balzani, A. Credi and M. Venturi, in Stimulating Concepts in Chemistry, M. Shibasaki, J. F. Stoddart and F. Vögtle, eds., Wiley-VCH, Weinheim, 2001, p. 255; (h) V. Balzani, M. Gomez-Lopez and J. F. Stoddart, Acc. Chem. Res., 1998, **31**(7), 405–414; (i) J. P. Sauvage, Acc. Chem. Res., 1998, **31**(10), 611–619.

- 4 (a) T. J. Hubin and D. H. Busch, *Coord. Chem. Rev.*, 2000, 200(202), 5–52; (b) T. J. Hubin, A. G. Kolchinski, A. L. Vance and D. H. Busch, 'Template Control of Supramolecular Architecture', in *Advances in Supramolecular Chemistry*, G. W. Gokel, ed., JAI Press, Connecticut, USA, 1999, vol. 5, pp. 237–357.
- 5 (a) G. F. Swiegers and T. J. Malefetse, Chem. Rev., 2000, 100, 3483–3537; (b) J.-P. Sauvage, Transition Metals in Supramolecular Chemistry, John Wiley & Sons, New York, 1999.
- 6 C. Kaes, A. Katz and M. W. Hosseini, *Chem. Rev.*, 2000, **100**, 3553–3590.
- 7 W. W. Brandt, Chem. Rev., 1954, 54, 959-1017.
- 8 A. M. W. C. Thompson, Coord. Chem. Rev., 1997, 160, 1-52.
- 9 M. Benaglia, F. Ponzini, C. R. Woods and J. S. Siegel, Org. Lett., 2001, 3(7), 967–969.
- 10 (a) S. Toyota, C. R. Woods, M. Benaglia, R. Haldimann, K. Warnmark, K. Hardcastle and J. S. Siegel, *Angew. Chem., Int. Ed.*, 2001, **40**(4), 751–754; (b) C. R. Woods, M. Benaglia, S. Toyota, K. Hardcastle and J. S. Siegel, *Angew. Chem., Int. Ed.*, 2001, **40**(4), 749–751.
- (a) C. O. Dietrich-Buchecker, J. P. Sauvage and J. P. Kintzinger, *Tetrahedron Lett.*, 1983, 24(46), 5095–5098; (b) M. Weck, B. Mohr, J. P. Sauvage and R. H. Grubbs, *J. Org. Chem.*, 1999, 64(15), 5463– 5471; (c) C. Dietrich-Buchecker, G. Rapenne and J. P. Sauvage, *Coord. Chem. Rev.*, 1999, 186(V186), 167–176.
- 12 C. O. Dietrich-Buchecker, P. A. Marnot and J. P. Sauvage, *Tetrahe*dron Lett., 1982, 23(50), 5291–5294.
- 13 (a) J.-C. Chambron, J.-P. Collin, V. Heitz, D. Jouvenot, J.-M. Kern, P. Mobian, D. Pomeranc and J.-P. Sauvage, *Eur. J. Org. Chem.*, 2004, **8**, 1627–1638; (b) P. Mobian, J.-M. Kern and J.-P. Sauvage, *Inorg. Chem.*, 2003, **42**(26), 8633–8637; (c) P. Mobian, J. M. Kern and J. P. Sauvage, *J. Am. Chem. Soc.*, 2003, **125**(8), 2016–2017; (d) F. Arico, P. Mobian, J. M. Kern and J. P. Sauvage, *Org. Lett.*, 2003, **5**(11), 1887–1890; (c) D. A. Leigh, P. J. Lusby, S. J. Teat, A. J. Wilson and J. K. Y. Wong, *Angew. Chem., Int. Ed.*, 2001, **40**(8), 1538–1543.
- 14 C. Piguet, G. Bernardinelli, A. F. Williams and B. Bocquet, Angew. Chem., Int. Ed. Engl., 1995, 34, 582–584.
- 15 J. P. Sauvage and M. Ward, Inorg. Chem., 1991, 30(20), 3869-3874.
- 16 (a) D. J. Cardenas, P. Gavina and J.-P. Sauvage, J. Am. Chem. Soc., 1997, **119**(11), 2656–2664; (b) D. J. Cardenas, J. P. Collin, P. Gavina, J. P. Sauvage, A. De Cian, J. Fischer, N. Armaroli, L. Flamigni, V. Vicinelli and V. Balzani, J. Am. Chem. Soc., 1999, **121**(23), 5481– 5488; (c) A. Livoreil, C. O. Dietrich-Buchecker and J. P. Sauvage, J. Am. Chem. Soc., 1994, **116**(20), 9399–9400.
- 17 For a recent report on a metal-free catenane templated by Zn(II) with terpyridine and phenanthroline, see: C. Hamann, J. M. Kern and J. P. Sauvage, *Inorg. Chem.*, 2003, **42**, 1877–1883.
- 18 N. Belfrekh, C. Dietrich-Buckecker and J.-P. Sauvage, *Inorg. Chem.*, 2000, **39**, 5169–5172.
- 19 (a) Doubly interlocked catenanes from symmetric components are also chiral by virtue of a helical twist. For examples of these analogs, see: A. C. Try, M. M. Harding, D. G. Hamilton and J. K. M. Sanders, *Chem. Commun.*, 1998, **6**, 723-724; (b) C. Dietrich-Buchecker and J. P. Sauvage, *Chem. Commun.*, 1999, **7**, 615–616; (c) J. F. Nierengarten, C. O. Dietrich-Buchecker and J. P. Sauvage, *New J. Chem.*, 1996, **20**(6), 685–693; (d) J. F. Nierengarten, C. O. Dietrich-Buchecker and J. P. Sauvage, *J. Am. Chem. Soc.*, 1994, **116**(1), 375–376; (e) F. Ibukuro, M. Fujita, K. Yamaguchi and J.-P. Sauvage, *J. Am. Chem. Soc.*, 1999, **121**(47), 11014–11015.
- 20 (a) J. C. Chambron, D. K. Mitchell and J. P. Sauvage, J. Am. Chem. Soc., 1992, 114(12), 4625–4631; (b) D. K. Mitchell and J. P. Sauvage, Angew. Chem., Int. Ed. Engl., 1988, 27(7), 930–931; (c) C. P. McArdle, S. Van, M. C. Jennings and R. J. Puddephatt, J. Am. Chem. Soc., 2001; (d) A. Mohry, F. Vögtle, M. Nieger and H. Hupfer, Chirality, 2000, 12(2), 76–83; (e) C. Yamamoto, Y. Okamoto, T. Schmidt, R. Jäger and F. Vögtle, J. Am. Chem. Soc., 1997, 119(43), 10547–10548; (f) S. Ottens-Hildebrandt, T. Schmidt, J. Harren and F. Vögtle, Liebigs Ann., 1995, 10, 1855–60; (g) P. R. Ashton, I. Iriepa, M. V. Reddington, N. Spencer, A. M. Z. Slawin, J. F. Stoddart and D. J. Williams, Tetrahedron Lett., 1994, 35(27), 4835–4838.
- 21 (a) J. J. Moore, J. J. Nash, P. E. Fanwick and D. R. McMillin, *Inorg. Chem.*, 2002, 41(24), 6387–6396; (b) F. Barigelletti, B. Ventura, J. P. Collin, R. Kayhanian, P. Gavina and J. P. Sauvage, *Eur. J. Inorg. Chem.*, 2000, 1, 113–119; (c) J. P. Collin, P. Gavina, J. P. Sauvage, A. DeCian and J. Fischer, *Aust. J. Chem.*, 1997, 50(10), 951–957;

- 22 (a) U. Lehmann, O. Henze and A. D. Schlüter, *Chem. Eur. J.*, 1999, 5(3), 854–859; (b) U. S. Schubert and C. Eschbaumer, *Org. Lett.*, 1999, 1(7), 1027–1029; (c) U. S. Schubert, C. Eschbaumer and G. Hochwimmer, *Synthesis*, 1999, 5, 779–782.
- 23 J. C. Loren and J. S. Siegel, Angew. Chem., Int. Ed., 2001, 40(4), 754-757.
- 24 The prefix [n] is often associated with term catenane to indicate the number of interlocking molecular cycles per molecule, a convention that is useful when comparing multi-component catenanes. This paper designates [2]catenane simply as catenane.
- 25 A. Fürstner, Angew. Chem., Int. Ed., 2000, 39, 3012-3043.
- 26 P. Siemsen, R. C. Livingston and F. Diederich, Angew. Chem., Int. Ed., 2000, 39, 2632–2657.
- 27 (a) A. Merz and H. Bachmann, J. Am. Chem. Soc., 1995, 117, 901– 908; (b) M. Delgado and J. D. Martin, J. Org. Chem., 1999, 64, 4798–4816.
- 28 I. P. Evans, A. Spencer and G. Wilkinson, J. Chem. Soc., Dalton Trans., 1973, 204–209.
- 29 (a) B. Mohr, M. Weck, J. P. Sauvage and R. H. Grubbs, Angew. Chem., Int. Ed., 1997, 36(12), 1308–1310; (b) P. Schwab, R. H. Grubbs and J. Ziller, J. Am. Chem. Soc., 1996, 118, 100–110.
- 30 Successful alkylation only occurred in the presence of tri(ethylene) glycol dimethyl ether (triglyme) and cesium carbonate. Little to no reactivity was observed in the absence of triglyme or when potassium carbonate was used, suggesting that a cesium-triglyme or cesium-polyethylene glycol coordinating effect facilitates the alkylation of the poorly soluble poly-phenoxide ion of **5**. This behavior was not observed when **5** was alkylated with polyethylene glycol linkers **21b** and **23b**. Here it is presumed that the linkers play the role of the triglyme.
- 31 Crystal structure analysis of **11a**: crystal from dichloromethane-benzene; empirical formula including solvent molecules $C_{172,23}H_{106}Cl_{3,11}F_{12}N_6O_{13}P_2Ru$, $M_r = 3000.6$; crystal system triclinic, space group $P\overline{1}$, a = 16.1362(14), b = 17.4816(16), c = 30.711(3) Å, a = 77.635(2), $\beta = 75.632(2)$, $\gamma = 65.267(2)^\circ$, V = 7562.3(12) Å³, Z = 2, $D_x = 1.318$ g cm⁻³, T = -173 °C; crystal dimensions: 0.19 × 0.17 × 0.16 mm, Bruker SMART area-detector diffractometer, MoK α radiation $\lambda = 0.71073$ Å, $\mu = 0.260$ mm⁻¹, $\theta_{max} = 22.57^\circ$, 41159 measured reflections, 19541 independent reflections, $R_{int} =$

0.044, 12817 reflections with $I > 2\sigma(I)$, refinement on F^2 with SHELXL-97, 1116 parameters, $R(F)[I > 2\sigma(I)$ reflections] = 0.098, $wR(F^2)$ [all reflections] = 0.307, $S(F^2) = 1.088$, $\Delta \rho_{max} = 1.31$ e Å⁻³. CCDC reference number 270917. See http://dx.doi.org/10.1039/ b506101f for crystallographic data in CIF or other electronic format.

- 32 (a) Crystal structure analysis of the Fe(II) complex with 3a: crystal obtained from dichloromethane-benzene; empirical formula including solvent molecules $C_{66}H_{64}F_{12}FeN_6O_5P_2$, $M_r = 1367.0$, orthorhombic, space group *Pbcn*, a = 23.3906(5), b = 11.4829(2), c = 23.2945(6) Å, V = 6256.7(2)Å³, Z = 4, $D_x = 1.451$ g cm⁻³, T = -113 °C, crystal di-mensions: $0.20 \times 0.10 \times 0.07$ mm, Nonius KappaCCD area-detector diffractometer, MoKa radiation, $\lambda = 0.71073$ Å, $\mu = 0.384$ mm⁻¹, $\theta_{\rm max} = 25^{\circ}, 75497$ measured reflections, 5489 independent reflections, 3041 reflections with $I > 2\sigma(I)$, absorption correction based on analysis of equivalent reflections (SORTAV), refinement on F^2 with SHELXL-97,³³ 481 parameters, $R(F) [I > 2\sigma(I) \text{ reflections}] = 0.075$, $wR(F^2)$ [all reflections] = 0.234, $S(F^2) = 1.021$, $\Delta \rho_{\text{max}} = 1.25$ e Å⁻³. The hexafluorophosphate anion is highly disordered; the asymmetric unit includes one 50%-occupied site for a water molecule. CCDC reference number 274362. See http://dx.doi.org/10.1039/b506101f for crystallographic data in CIF or other electronic format; (b) Crystal structure analysis of the Ru(II) complex with 3a: crystal obtained from dichloromethane-benzene; empirical formula including solvent molecules $C_{66}H_{64}F_{12}N_6O_5P_2Ru$, $M_r = 1412.2$, orthorhombic, space group *Pbcn*, a = 23.1733(3), b = 11.9603(1), c = 22.8083(3) Å, V = 6321.5(1) Å³, Z = 4, $D_x = 1.484$ g cm⁻³, T = -113 °C, crystal dimensions: $0.20 \times 0.18 \times 0.07$ mm, Nonius KappaCCD area-detector diffractometer, MoK α radiation, $\lambda = 0.71073$ Å, $\mu = 0.390 \text{ mm}^{-1}, \ \theta_{\text{max}} = 25^{\circ}, \ 101070 \text{ measured reflections}, \ 5577$ independent reflections, 3965 reflections with $I > 2\sigma(I)$, absorption correction based on analysis of equivalent reflections (SORTAV), refinement on F^2 with SHELXL-97,³³ 481 parameters, R(F) [I > $2\sigma(I)$ reflections] = 0.061, $wR(F^2)$ [all reflections] = 0.189, $S(F^2)$ = 1.044, $\Delta \rho_{\text{max}} = 1.31$ e Å⁻³. The hexafluorophosphate anion is highly disordered; the asymmetric unit includes one 50%-occupied site for a water molecule. CCDC reference number 274363. See http://dx.doi.org/10.1039/b506101f for crystallographic data in CIF or other electronic format.
- 33 G. M. Sheldrick, SHELXL-97, Program for refinement of crystal structures, University of Göttingen, Germany, 1997.