Inorg. Chem. 2005, 44, 3103–3111



Square Planar vs Tetrahedral Geometry in Four Coordinate Iron(II) Complexes

Eric J. Hawrelak,^{†,‡} Wesley H. Bernskoetter,[†] Emil Lobkovsky,[†] Gordon T. Yee,[§] Eckhard Bill,^{II} and Paul J. Chirik^{*,†}

Department of Chemistry and Chemical Biology, Baker Laboratory, Cornell University, Ithaca, New York 14853, Department of Chemistry, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061-0212, and Max-Planck-Institute for Bioinorganic Chemistry, Stiftstrasse 34-36, D-45470 Mülheim an der Ruhr, Germany

Received December 20, 2004

The geometric preferences of a family of four coordinate, iron(II) d⁶ complexes of the general form L₂FeX₂ have been systematically evaluated. Treatment of Fe₂(Mes)₄ (Mes = 2,4,6-Me₃C₆H₂) with monodentate phosphine and phosphite ligands furnished square planar *trans*-P₂Fe(Mes)₂ derivatives. Identification of the geometry has been accomplished by a combination of solution and solid-state magnetometry and, in two cases (P = PMe₃, PEt₂Ph), X-ray diffraction. In contrast, both tetrahedral and square planar coordination has been observed upon complexation of chelating phosphine ligands. A combination of crystallographic and magnetic susceptibility data for (depe)Fe-(Mes)₂ (depe = 1,2-bis(diethylphosphino)ethane) established a tetrahedral molecular geometry whereas SQUID magnetometry and Mössbauer spectroscopy on samples of (dppe)Fe(Mes)₂ (dppe = 1,2-bis(diphenylphosphino)ethane) indicated a planar molecule. When dissolved in chlorinated solvents, the latter compound promotes chlorine atom abstraction, forming tetrahedral (dppe)Fe(Mes)CI and (dppe)FeCl₂. Ligand substitution reactions have been studied for both structural types and are rapid on the NMR time scale at ambient temperature.

Introduction

The geometry and electronic structure of a transition metal complex are interrelated properties and dictate both the physical characteristics and chemical reactivity of a given molecule. Four coordinate compounds of the first transition series are particularly intriguing, given the choice between a low-/intermediate-spin square planar ground state and a high-spin tetrahedral configuration (Scheme 1).^{1,2} The tetrahedral arrangement is sterically preferred and typically observed with large ligands and small metal ions.³ For high-spin complexes with dⁿ electronic configurations with $n \ge 2$, occupation of antibonding molecular orbitals is unavoid-

- [§] Virginia Polytechnic Institute and State University.
- ^{II} Max-Planck-Institute for Bioinorganic Chemistry.

- (2) The d-orbital splitting diagrams shown in Scheme 1 are for idealized, high-symmetry cases.
 (3) Greenwood, N. N.; Earnshaw, A. *Chemistry of the Elements*, 2nd ed.;
- (3) Greenwood, N. N.; Earnshaw, A. Chemistry of the Elements, 2nd ed.; Butterworth Heinemann: Oxford, U.K., 1997; pp 1156–1166.

10.1021/ic048202+ CCC: \$30.25 © 2005 American Chemical Society Published on Web 03/29/2005





able, providing little ligand field stabilization energy (LFSE). In contrast, square planar coordination, while sterically disfavored, offers substantial electronic stabilization that is a natural consequence of a high-lying, empty, antibonding $d(x^2-y^2)$ orbital (Scheme 1).

For d⁸ metal ions, the factors that govern the choice between square planar and tetrahedral geometry are fairly well understood.³ For larger second and third row metals such as Pd(II), Pt(II), and Au(III), the LFSE dominates and square planar geometries are almost exclusively observed.³ The lighter Ni(II) congener offers both limiting structural types. When weak field ligands such as halides or arylated phosphines are present, as is the case with NiCl₄^{2–} or (Ph₃P)₂-NiCl₂, tetrahedral coordination is observed^{4.5} whereas compounds with strong field ligands or alkylated phosphines such

Inorganic Chemistry, Vol. 44, No. 9, 2005 3103

 $[\]ast$ Author to whom correspondence should be addressed. E-mail: pc92@ cornell.edu.

[†] Cornell University.

[‡] Current address: Department of Chemistry, 237A Hartline Science Center, Bloomsburg University, Bloomsburg, PA 17815.

⁽¹⁾ Favas, M. C.; Kepert, D. L. Prog. Inorg. Chem. 1980, 27, 325.

as Ni(CN)₄²⁻ and (Cy₃P)₂NiCl₂ are square planar.⁶ Interestingly, complexes supported by mixed alkyl/aryl phosphines such as $[(PhCH_2)Ph_2P]_2NiBr_2$ contain both tetrahedral and square planar geometries in the same crystal lattice.⁷

Unlike the d⁸ cases, the geometric preferences of lower d^n configurations have not been as thoroughly studied. Our laboratory has been exploring the chemistry of organometallic iron complexes as potential surrogates for precious metals in catalytic bond forming reactions such as olefin hydrogenation and hydrosilation.⁸ As part of these investigations, we have prepared and characterized a family of four coordinate, in some cases enantiomerically pure, iron(II) dialkyl complexes.9 While formally 14 electron species, these compounds are unreactive toward olefins, possibly a consequence of their tetrahedral geometry and high-spin, S = 2ground state, which provides no vacant low-lying molecular orbitals for substrate coordination. On the basis of this hypothesis, we became interested in exploring alternative geometries and spin states with the goal of generating catalytically active iron complexes.

Square planar coordination for iron(II) L_2FeX_2 compounds, while amply precedented, is generally limited to complexes containing porphyrin, salicylaldimato, and phthalocyanine ligands, where the macrocyclic chelate confines the geometric preference of the metal center.^{10,11} Iron(II) complexes containing only mono- or bidentate ligands where the molecule can adopt either geometry have not been systematically investigated. While both structural types have been reported, only tetrahedral L_2FeX_2 complexes have been crystallographically characterized.^{12–15} The geometry of the square planar complexes, such as (PhEt₂P)₂Fe(C₆Cl₅)₂,¹⁶ has

- (4) Gill, N. S.; Nyholm, R. S. J. Chem. Soc. 1959, 3997.
- (5) Venanzi, L. M. J. Chem. Soc. 1958, 719.
- (6) Cotton, F. A.; Wilkinson, G. Advanced Inorganic Chemistry, 5th ed.; Wiley-Interscience: New York, 1988; pp 742–755.
- (7) Kilbourn, B. T.; Powell, H. M.; Darbyshire, J. A. C. Proc. Chem. Soc. 1963, 207.
- (8) Bart, S. C.; Lobkovsky, E.; Chirik, P. J. J. Am. Chem. Soc. 2004, 126, 13794.
- (9) Bart, S. C.; Hawrelak, E. J.; Schmisseur, A. K.; Lobkovsky, E.; Chirik, P. J. Organometallics 2004, 23, 237.
- (10) Examples of square planar, L₂FeX₂ complexes with macrocyclic ligands have been crystallographically characterized. For representative examples, see: (a) Kirner, J. F.; Dow, W.; Scheidt, W. R. Inorg. Chem. **1976**, 15, 1685. (b) Silver, M. E.; Ibers, J. A. J. Am. Chem. Soc. **1983**, 105, 4108. (c) Strauss, S. H.; Silver, M. E.; Long, K. M.; Thompson, R. G.; Hudgens, R. A.; Spartalian, K.; Ibers, J. A. J. Am. Chem. Soc. **1985**, 107, 4207. (d) Ghiladi, R. A.; Kretzer, R. M.; Guzei, I.; Rheingold, A. L.; Neuhold, Y.-M.; Hatwell, K. R.; Zuberbuhler, A. D.; Karlin, K. D. Inorg. Chem. **2001**, 40, 5754. (e) Jacoby, D.; Floriani, C.; Chiesi-Villa, A.; Rizzoli, C. Chem. Commun. **1991**, 220.
- (11) For representative examples, see: (a) Evanglesti, M.; Bartolome, J.; de Jongh, L. J.; Filoti, G. *Phys. Rev. B* 2002, 66, 144410. (b) Ohya, T.; Takeda, J.; Kobayashi, N.; Sato, M. *Inorg. Chem.* 1990, 29, 3734. (c) Ohya, T.; Kobayashi, N.; Sato, M. *Inorg. Chem.* 1987, 26, 2506. (d) Labarta, A.; Molins, E.; Vinas, X.; Tejada, J.; Caubet, A.; Alvarez, S. *J. Chem. Phys.* 1984, 80, 444. (e) Molins, E.; Labarta, A.; Tejada, J.; Caubet, A.; Alvarez, S. *Transition Met. Chem.* 1983, 83, 377. (f) Dale, B. W.; Williams, R. J.; Edwards, P. R.; Johnson, C. E. *J. Chem. Phys.* 1968, 49, 3445.
- (12) Hermes, A. R.; Girolami, G. S. Organometallics 1987, 6, 763.
- (13) Hill, D. H.; Sen, A. J. Am. Chem. Soc. 1988, 110, 1651.
- (14) Klose, A.; Solari, E.; Floriani, C.; Chiesi-Villa, A.; Rizzoli, C.; Re, N. J. Am. Chem. Soc. 1994, 116, 9123.
- (15) Müller, H.; Seidel, W.; Görls, H. Z. Anorg. Allg. Chem. 1996, 622, 756.
- (16) Chatt, J.; Shaw, B. L. J. Chem. Soc. 1961, 285.



Figure 1. Molecular structure of *trans*- $(PhEt_2P)_2Fe(C_6Cl_5)_2$ at 30% probability ellipsoids. Hydrogen atoms are omitted for clarity.

Table 1. Selected Bond Distances (Å) and Angles (deg) for trans-(PhEt₂P)₂Fe(C₆Cl₅)₂ and **6**

trans-(PhEt ₂ P) ₂ Fe(C ₆ Cl ₅) ₂		6	
Fe(1)-C(1) Fe(1)-P(1)	1.999(3) 2.2990(8)	Fe(1)-C(1) Fe(1)-P(1)	2.031(2) 2.2597(6)
$\begin{array}{l} C(1) - Fe(1) - C(1A) \\ C(1) - Fe(1) - P(1) \\ C(1A) - Fe(1) - P(1) \\ C(1A) - Fe(1) - P(1A) \end{array}$	180.0 88.31(9) 91.69(9) 88.31(9)	$\begin{array}{l} C(1)-Fe(1)-C(1A) \\ C(1)-Fe(1)-P(1) \\ C(1A)-Fe(1)-P(1) \\ C(1A)-Fe(1)-P(1A) \end{array}$	180.0 89.72(7) 90.29(7) 90.28(7)

been inferred on the basis of magnetic susceptibility data, where compounds with S = 1 ground states are consistent with intermediate-spin, d⁶ planar molecules.¹⁷ Here we describe the synthesis and characterization of a family of P₂Fe(Mes)₂ (P = phosphine, phosphite; Mes = 2,4,6-Me₃C₆H₂) complexes and explore the factors that govern their geometric preferences. In addition, crystallographic characterization of (PhEt₂P)₂Fe(C₆Cl₅)₂ and two square planar P₂-Fe(Mes)₂ compounds is also presented.

Results and Discussion

Crystallographic and NMR Spectroscopic Characterization of *trans*-(PhEt₂P)₂Fe(C₆Cl₅)₂. Because no example of a square planar iron(II) compound with monodentate ligands had been structurally characterized, our study commenced with crystallographic verification of the structure of the previously reported trans-(PhEt₂P)₂Fe(C₆Cl₅)₂.⁹ Yellow plates suitable for X-ray diffraction were obtained by layering a concentrated dichloromethane solution with pentane. The solid-state structure (Figure 1) definitively establishes the square planar geometry of the molecule, in agreement with the previous assignment based on magnetometry.9 Selected bond distances and angles are presented in Table 1. The crystallography also reveals trans-phosphine and pentachlorophenyl ligands that are related by a center of inversion. To avoid steric interactions with the phosphines, the aryl ligands are orthogonal to the square plane, placing the orthochlorine substituents above and below the iron center. The structural data are consistent with Chatt and Shaw's asser-

⁽¹⁷⁾ Seidel, V. W.; Lattermann, K.-J. Z. Anorg. Allg. Chem. 1982, 488, 69.

Chart 1



tion¹⁶ that strong field, *ortho*-substituted aryl groups are essential for square planar coordination. In addition, the aryl groups on the phosphine ligands are essentially parallel to the pentachlorophenyl ligands, presumably to avoid steric interactions with the ring substituents.

Analysis of *trans*-(PhEt₂P)₂Fe(C₆Cl₅)₂ by ¹H NMR spectroscopy in benzene- d_6 confirms the presence of only one compound in solution. As expected for an S = 1 molecule, the phosphine resonances are shifted dramatically from their diamagnetic reference values. For example, the ethyl substituents on the phosphine ligands appear at -18.6 (CH₃) and -0.15 (CH₂) ppm, while three peaks are observed at -7.8, -53.2, and 12.1 (*para*) ppm for the corresponding phenyl protons. Attempts to observe either ³¹P or ¹³C NMR spectra have been unsuccessful.

Evaluation of Geometric Preferences of L₂Fe(Mes)₂ Compounds. With both NMR spectroscopic and solid-state characterization of a square planar iron(II) complex in hand, the geometric preferences of a family of four coordinate Fe-(II), d⁶ compounds were systematically evaluated. For rapid screening, a series of neutral donors was added to Fe₂(Mes)₄ (eq 1). This iron precursor was chosen due to its synthetic availability, versatility as a synthon for a range of iron complexes, and reported ability to provide both square planar and tetrahedral complexes. Using this approach, Floriani and co-workers have prepared $(py)_2Fe(Mes)_2$ and $(phen)Fe(Mes)_2$ (py = pyridine, phen = 9,10 phenanthroline).^{14,18} X-ray crystallography has established that both molecules are tetrahedral in the solid state. Likewise, Siedel and co-workers have synthesized and structurally characterized tetrahedral $(dme)Fe(Mes)_2$ (dme = 1,2-dimethoxyethane).¹⁵



Nitrogen donors were the first class of ligand explored. Treatment of Fe₂(Mes)₄ with *N*,*N*-dimethylaminopyridine (DMAP) furnished a yellow powder identified as (DMAP)₂Fe-(Mes)₂ (1) (Chart 1). The solution magnetic moment (benzene- d_6 , 22 °C) of 1 was measured by the Evans method²⁵ and produced a magnetic moment of 5.1 μ_B ,

- (18) Magill, C. P.; Floriani, C.; Chiesi-Villa, A.; Rizzoli, C. *Inorg. Chem.* **1994**, *33*, 1928.
- (19) Carlin, R. L. *Magnetochemistry*; Springer-Verlag: Berlin, 1986; p 24. (20) Sellers, S. P.; Korte, B. J.; Fitzgerald, J. P.; Reiff, W. M.; Yee, G. T.
- *J. Am. Chem.* 502, **1998**, *120*, 4662, 100
- (21) McGarvey, B. R. Inorg. Chem. 1988, 27, 4691.
- (22) Lorber, C.; Choukroun, R.; Costes, J.-P.; Donnadieu, B. C. R. Chim. 2002, 5, 251.



Figure 2. SQUID magnetization data for 3.

consistent with a high-spin, S = 2 tetrahedral molecule. This result is consistent with Floriani's observation of a tetrahedral geometry for (py)₂Fe(Mes)₂, which has also been characterized by X-ray diffraction.¹⁴ Addition of a chelating diamine such as *N*,*N*,*N'*,*N'*-tetramethylethylenediamine (tmeda) to Fe₂-(Mes)₄ yielded (tmeda)Fe(Mes)₂ (**2**) as a tan solid in modest yield (Chart 1). Measurement of the benzene-*d*₆ solution magnetic moment at 22 °C produced a value of 5.1 $\mu_{\rm B}$, consistent with an S = 2 tetrahedral molecule. Addition of more hindered amines such as NEt₃ or *N*,*N*,*N'*,*N'*-tetraethylethylenediamine to Fe₂(Mes)₄ produced no reaction.

In addition to nitrogen-based ligands, bis(mesityl)iron(II) complexes supported by phosphines and phosphites were studied. The commercial availability of these ligands along with their diverse steric and electronic properties makes them attractive for a systematic study of the coordination preferences of four coordinate iron(II) complexes.

Addition of the smallest phosphine in the series, PMe₃, to Fe₂(Mes)₄ in diethyl ether at -78 °C afforded (Me₃P)₂Fe- $(Mes)_2$ (3) as a yellow crystalline solid in high yield. The solution magnetic moment of 3 was measured by the Evans method in benzene- d_6 and produced a value of 2.9 $\mu_{\rm B}$, consistent with the spin-only value for two unpaired electrons. The S = 1 ground state suggests an intermediate-spin, d⁶ planar molecule. Corroboration of this value was obtained from solid state magnetic susceptibility measurements on a powdered sample of **3** (Figure 2). The γT versus T data may be fitted to a zero-field-splitting model for $S = 1^{19}$ with a temperature-independent paramagnetism term that yields g $= 2.26 \pm 0.01$, $D/k = 51 \pm 1$ K, and TIP $= 5 \times 10^{-4}$ (emu K)/mol. The high value of the TIP may reflect a trace ferromagnetic impurity. Deviation of the g-value from the free electron value is consistent with incomplete quenching of the orbital angular momentum as has been seen previously for square planar Fe(II).²⁰

As with *trans*-(PhEt₂P)₂Fe(C₆Cl₅)₂, ¹H NMR spectroscopy has proven useful for the identification of **3**. A representative spectrum recorded at 22 °C in benzene- d_6 is presented in Figure 3. A relatively sharp ($\Delta v_{1/2} = 36$ Hz) peak corresponding to 18 hydrogens is observed at -35.8 ppm and is assigned as the phosphine methyl groups.²¹ Distinct resonances for the *ortho* and *para* methyl groups of the mesityl ligands are observed at 20.5 and -4.8 ppm, respectively,



Figure 3. ¹H NMR spectrum of 3 in benzene- d_6 .



Figure 4. Molecular structure of **3** at 30% probability ellipsoids. Hydrogen atoms are omitted for clarity.

Table 2. Selected Bond Distances (Å) and Angles (deg) for 3

Fe(1)-C(7)	2.032(3)	Fe(1)-P(1)	2.2552(10)
Fe(1)-C(16)	2.017(3)	Fe(1)-P(2)	2.2475(10)
C(16)-Fe(1)-C(7)	176.61(13)	C(7)-Fe(1)-P(1)	89.62(9)
C(16)-Fe(1)-P(1)	89.31(9)	C(7)-Fe(1)-P(2)	90.91(9)
C(16)-Fe(1)-P(2)	90.91(9)	P(1)-Fe(1)-P(2)	177.74(4)

while a single resonance is observed at 25.5 ppm for the *meta* hydrogen on the aryl rings.

Single crystals suitable for X-ray diffraction were obtained by cooling a concentrated diethyl ether solution of **3** to -35°C. The molecular structure is shown in Figure 4, and selected bond distances and angles are presented in Table 2. As with the molecular structure of *trans*-(PhEt₂P)₂Fe(C₆Cl₅)₂, **3** is square planar with trans phosphine and mesityl ligands. The mesityl ligands are oriented perpendicular to the plane



of the iron with the *ortho* methyl groups avoiding steric interactions with the phosphine ligands and protecting the axial sites on the iron center, again supporting the Chatt and Shaw hypothesis.¹⁶

Using the synthetic approach outlined in eq 1, several other bis(phosphine)iron(II) bis(mesityl) complexes were prepared in moderate to good yield (Figure 5). Each complex was characterized by a combination of ¹H NMR spectroscopy, elemental analysis, and solution magnetometry. For all of the complexes containing monodentate phosphine ligands, the magnetic susceptibility data are consistent with intermediate-spin, S = 1 ground states, indicative of a square planar geometry. The geometry of 6 was corroborated by X-ray diffraction, and the solid-state structure is presented in Figure 6, while selected bond distances and angles are reported in Table 1. 6 is structurally similar to both trans-(PhEt₂P)₂Fe- $(C_6Cl_5)_2$ and 3 with a square planar iron center and *trans*phosphine ligands. The orientation of both the phosphines and the aryl ligands is similar to that observed in trans- $(PhEt_2P)_2Fe(C_6Cl_5)_2$. The iron-carbon bonds in 6 are elongated by 0.03 Å as compared to those of *trans*-(PhEt₂P)₂-



Figure 6. Molecular structure of 6 at 30% probability ellipsoids. Hydrogen atoms are omitted for clarity.

 $Fe(C_6Cl_5)_2$, while the iron-phosphorus bonds are contracted by approximately the same distance.

Treatment of Fe₂(Mes)₄ with the trialkyl phosphite, P(OEt)₃, furnished *trans*-[(EtO)₃P]₂Fe(Mes)₂ (**8**), which has a solution magnetic moment (benzene- d_6 , 22 °C) of 2.7 μ_B , consistent with a planar, intermediate-spin S = 1 molecule. Attempts to prepare similar bis(phosphite) complexes by addition of more hindered and electron-poor precursors such as P(OPh)₃, P(OSiMe₃)₃, or P(O-2,6-tBu₂C₆H₃)₃ produced no reaction.

The effect of chelating phosphines on the geometric preference of bis(phosphine)iron(II) bis(mesityl) complexes was also explored. Stirring either dppf (dppf = 1,2-bis-(diphenylphosphino)ferrocene) or (\pm) -BINAP (BINAP = 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl) with Fe₂(Mes)₄ in diethyl ether produced no reaction even after extended reaction times or gentle heating. However, addition of dmpe (1,2-bis(dimethylphosphino)ethane) to an ethereal solution of Fe₂(Mes)₄ furnished a bright yellow powder identified as (dmpe)Fe(Mes)₂ (9) on the basis of magnetic data and elemental analysis.

Further characterization of **9** by solution ¹H NMR spectroscopy was hampered by the low solubility of the compound in common organic solvents such as toluene, benzene, THF, diethyl ether, acetone, acetonitrile, and dimethyl sulfoxide. Dissolution of **9** in chlorinated solvents such as dichloromethane- d_2 resulted in decomposition (vide infra). Despite these limitations, a magnetic moment of $4.2 \mu_B$ was measured in the solid state using the Gouy method. While low, this value is most consistent with tetrahedral coordination. The origin of the low moment is not understood at this time and may be a consequence of an impurity or population of a low-spin, planar isomer. It should be noted, however, that the magnetic measurements were conducted on analytically pure material.

To circumvent the solubility problems associated with 9, a chelating phosphine with more lipophilic alkyl substituents was explored. Addition of depe (1,2-bis(diethylphosphino)ethane) to Fe₂(Mes)₄ furnished an orange powder identified as (depe)Fe(Mes)₂ (**10**) in 79% yield (Figure 7). Solution



Figure 7. Iron(II) bis(mesityl) complexes with chelating phosphine ligands.



Figure 8. Molecular structure of **10** at 30% probability ellipsoids. Hydrogen atoms are omitted for clarity.

Table 3. Selected Bond Distances (Å) and Angles (deg) for 10

Fe(1)-C(11)	2.0702(14)	Fe(1)-P(1)	2.4692(4)
Fe(1)-C(20)	2.0745(3)	Fe(1)-P(2)	2.4859(4)
C(11)-Fe(1)-C(20)	120.41(5)	C(20)-Fe(1)-P(1)	100.90(4)
C(11)-Fe(1)-P(1)	126.18(4)	C(20)-Fe(1)-P(2)	123.20(4)
C(11)-Fe(1)-P(2)	100.97(4)	P(1)-Fe(1)-P(2)	80.35(1)

magnetometry in benzene- d_6 at 22 °C provided a magnetic moment of 4.4 μ_B , slightly lower than the spin-only value expected for four unpaired electrons but well within the range of high-spin tetrahedral ferrous complexes.⁹

The tetrahedral geometry of 10 was corroborated by X-ray diffraction. The solid-state structure is presented in Figure 8, and selected bond distances and angles are provided in Table 3. As expected for a tetrahedral iron center, the mesityl ligands are oriented above and below the iron-phosphine chelate plane and are slightly twisted to avoid steric interactions between the ortho methyl groups. The ethyl substituents on the phosphorus atoms are directed in a "front-back" arrangement where the two carbon chain directed toward the iron is parallel to the plane of the adjacent aryl ring. Comparison of the iron-phosphorus bond distances in 10 to those found in *trans*-(PhEt₂P)₂Fe(C_6Cl_5)₂, 3, and 6 establishes a lengthening by approximately 0.2 Å from square planar to tetrahedral coordination, consistent with population of antibonding molecular orbitals in the tetrahedral case. In contrast, little change is observed in the iron-carbon bond lengths in the two limiting geometries.

The tetrahedral geometry of 10 is somewhat surprising in light of the observation of square planar complexes for the monodentate phosphines and phosphite in 3-8. At first glance, it may appear that the tetrahedral geometry is favored



Figure 9. SQUID magnetization data for 10 (red) and 11 (blue).

due to the inability of the chelate to adopt a trans disposition of the phosphines and mesityl ligands. However, Siedel and Lattermann¹⁷ have reported (dppe)Fe(Mes)₂ (**11**) (dppe = 1,2-bis(diphenylphosphino)ethane) as a planar molecule on the basis of magnetic susceptibility data. Observation of a square planar complex with *arylated* chelating phophines and tetrahedral molecules with *alkylated* phosphines also contrasts the coordination preferences observed in classic Ni-(II) complexes.³⁻⁶

On the basis of these puzzling observations, 11 was prepared and studied in more detail. As was reported previously,¹⁷ addition of dppe to an ethereal solution of Fe₂- $(Mes)_4$ afforded an orange powder, identified as **11**, that is sparingly soluble in ethereal and hydrocarbon solvents. The low solubility of the compound in common NMR solvents precluded measurement of the solution magnetic moment. Solid-state magnetic data were collected from 2 to 300 K using SQUID magnetometry (Figure 9). For comparison, similar data was recorded on powdered samples of 10, a crystallographically verified tetrahedral molecule. For 11, the magnetic moment from 50 to 300 K ranges from 3.3 to 3.9 $\mu_{\rm B}$ and is slightly higher than the values measured over the same temperature range for 3. Importantly, the values of the magnetic susceptibility for 11 are substantially lower than those for 10, supporting Siedel's original conclusion of a planar molecule with an intermediate-spin, S = 1 ground state.

Mössbauer spectroscopy was also used to investigate the geometry of **11**. For comparison, spectra were recorded on the known tetrahedral compounds, **10** and (–)-(sparteine)-FeCl₂²² as well as square planar **6**. Selected spectra are presented in Figure 10, and the experimentally determined isomer shifts and quadrupole splittings are reported in Table 4. As expected for a series of iron(II) compounds, the isomer shifts are essentially invariant for the four compounds. In contrast, **6** and **11** have large quadrupole splitting parameters consistent with planar, S = 1 molecules in which covalency contributions to the electric field gradient are significant. The unique appearance of the zero-field spectra showing unusually large quadrupole splittings of more than 4 mm/s and small isomer shifts of 0.3–0.5 mm/s is fully characteristic of ferrous ions in the intermediate S = 1 spin state. Thus,



Figure 10. Zero-field Mössbauer spectra for 6, 10, and 11 recorded at 80 K.

Table 4. Mössbauer Parameters for a Series of Four Coordinate

 Iron(II) Compounds

compound	δ (mm/s)	quad splitting (mm/s)
6	0.31	4.63
10	0.39	1.71
11	0.33	4.53
((-)-sparteine)FeCl ₂	0.91	2.43

both magnetic susceptibility studies and Mössbauer spectroscopy are consistent with a square planar geometry for **11**.

Attempts to verify the geometry of **11** by X-ray diffraction have been frustrated by the inability to obtain single crystals. The complex is essentially insoluble in both hydrocarbon and ethereal solvents but dissolves freely in dichloromethane. Recrystallization of the complex from a concentrated CH₂-Cl₂ solution of **11** layered with diethyl ether at -35 °C afforded large yellow blocks suitable for X-ray diffraction. The solid-state structure, shown in Figure 11, is not the desired complex but rather (dppe)Fe(Mes)Cl (**12**). Monitoring the reaction between **11** and dichloromethane- d_2 by ¹H NMR spectroscopy established clean conversion to **12** over the course of minutes at ambient temperature. Concomitant with



Figure 11. Molecular structure of **12** at 30% probability ellipsoids. Hydrogen atoms are omitted for clarity.

Table 5. Selected Bond Distances (Å) and Angles (deg) for 12

Fe(1)-C(1)	2.044(5)	Fe(1)-P(1)	2.4598(15)
Fe(1)-Cl(1)	2.2486(14)	Fe(1)-P(2)	2.5169(15)
C(1)-Fe(1)-Cl(1)	117.17(14)	Cl(1)-Fe(1)-P(1)	98.60(6)
C(1)-Fe(1)-P(1)	123.21(14)	Cl(1)-Fe(1)-P(2)	127.06(16)
C(1) - Fe(1) - P(2)	127.06(16)	P(1) - Fe(1) - P(2)	82.01(5)

this process is a second chlorine atom abstraction by 12 to form (dppe)FeCl₂ (eq 2). This competing atom transfer process makes isolation of pure 12, free of 11 or (dppe)-FeCl₂, challenging. In both chlorine atom abstraction reactions, the fate of the mesityl ligand is unknown.



The solid-state structure (Figure 11) and metrical parameters (Table 5) clearly establish a tetrahedral geometry for **12**. The remaining mesityl group is oriented essentially perpendicular to the iron-chlorine bond vector while the phenyl substituents on the phosphorus atoms are arranged in a propeller-like fashion and are related by an idealized C_2 axis of symmetry. Importantly, the iron-phosphorus bond lengths of 2.4598(15) and 2.5169(15) Å found in **12** are elongated from the distances of ~2.3 Å present in the corresponding square planar complexes. In comparison, the iron-carbon bond is barely elongated in **12** as compared to the square planar complexes.

Because substitution of one mesityl ligand by a chloride induces a geometric change from square planar to tetrahedral, the coordination preferences of iron(II) bis(halide) complexes were examined. Heating 2 equiv of PEt₂Ph, a phosphine known to support square planar complexes, with either anhydrous FeCl₂ or FeBr₂ furnished (PhEt₂P)₂FeX₂ (X = Cl (**13**), Br (**14**)) (eq 3). Measurement of the solution magnetic moments (Evans method) yielded μ_{eff} values of 4.8 (**13**) and 5.0 $\mu_{\rm B}$ (**14**), consistent with a *S* = 2 ground states and a tetrahedral geometries. Observation of tetrahedral complexes with **12–14** highlights the importance of *ortho*-substituted aryl groups for isolation of square planar complexes.



Several other bis(phosphine)ferrous complexes were targeted to further test this hypothesis. Attempts to isolate Fe₂-Ph₄ or trap the compound in situ in the presence of both monodentate and chelating phosphines resulted in decomposition. Likewise, attempts to alkylate (dppe)FeCl₂ or **13** with MeLi or KCH₂Ph did not produce a tractable product. It should be noted that Hermes and Girolami have synthesized and in many cases crystallographically characterized (dippe)FeR₂ (dippe = bis(diisopropylphosphino)ethane) complexes, all of which are tetrahedral.¹² **Ligand Exchange Reactions.** The isolation and crystallographic characterization of a family of square planar iron-(II) complexes offered the opportunity to study ligand exchange processes at an intermediate-spin, S = 1 metal center and potentially compare the outcome of these processes to more familiar, S = 0, d⁸ complexes. Addition of 2 equiv of PMe₃ to **4** resulted in rapid formation of **3** with liberation of 2 equiv of free PEt₃. A similarly rapid reaction was observed for the addition of PMe₃ to **6** (eq 4). Addition of PEt₃ or PEt₂Ph to **3** produced no reaction, indicating that coordination of PMe₃ is in fact thermodynamically favored. Attempts to measure rate constants for the exchange reactions have not been successful due to the rapid rate at which these reactions reach equilibrium.



Exchange reactions were also carried out with **5** and **6**. Addition of PMe₂Ph to **6** rapidly furnished **5** and free PEt₂-Ph whereas, in the converse experiment, addition of PEt₂Ph to **5** produced no ligand exchange (eq 5). Given the electronic similarity of these two ligands in combination with the results of the PMe₃ exchange studies, the thermodynamic preference for phosphine coordination appears to be dominated by steric effects, where smaller ligands bind more tightly to the [Fe-(Mes)₂] fragment than their more hindered counterparts.



Exchange reactions between square planar and tetrahedral complexes were also investigated. Addition of 2 equiv of PMe₃ to $(py)_2Fe(Mes)_2$ resulted in immediate and quantitative ligand exchange at 22 °C in benzene- d_6 . Addition of a slight excess of pyridine to **3** produced no reaction. These experiments clearly demonstrate the thermodynamic preference for phosphine over amine coordination. As with the square planar complexes, substitution of pyridine in $(py)_2Fe(Mes)_2$ with PMe₃ or pyridine- d_5 is too fast at ambient temperature to be measured by conventional NMR techniques.



Concluding Remarks

The coordination preferences of a family of four coordinate, $d^6 L_2 FeX_2$ complexes have been systematically studied as a function of both neutral and anionic ligands. For iron-(II) bis(mesityl) compounds, high-spin, S = 2 tetrahedral

complexes are observed exclusively when supported by monodentate or chelating amine ligands. These results, in conjunction with those previously reported for oxygen chelates,¹⁵ suggest that, for first row donors, steric factors overcome competing electronic stabilization offered by the square planar alternative. For monodentate phosphines and phosphites, the opposite situation is observed, where planar, intermediate-spin, S = 1 complexes are favored. Ligand exchange studies establish a preference for phosphine over amine coordination and hence a square planar over a tetrahedral geometry. Thus, stronger phosphine ligands overcome the dominant steric effect observed with the nitrogen and oxygen donors, providing greater ligand field stabilization energy and square planar complexes. Within the series of phosphines, preferential coordination of smaller phosphines was observed.

For chelating phosphines, the conclusions are not as straightforward. Alkyl-substituted chelates such as dmpe and depe furnished tetrahedral molecules whereas the arylated phosphine, dppe, afforded a square planar complex, contrasting observations in classic Ni(II), d⁸ chemistry. This shift in geometric preference is not immediately obvious and may be a result of subtle steric interactions. Significantly, replacement of the *ortho*-substituted aryl ligands with weaker field halides induces a change in geometric preference, producing tetrahedral molecules regardless of the field strength of the supporting neutral ligand. These observations are in agreement with Chatt and Shaw's assertion¹⁶ that *ortho*-substituted aryl groups are necessary for intermediate-spin, S = 1 molecules.

Experimental Section

General Considerations. All air- and moisture-sensitive manipulations were carried out using standard vacuum line, Schlenk, and cannula techniques or in an Mbraun inert-atmosphere drybox containing an atmosphere of purified nitrogen. Solvents for airand moisture-sensitive manipulations were initially dried and deoxygenated using literature procedures.²³ The Mbraun drybox was equipped with a cold well designed for freezing samples in liquid nitrogen. Argon and hydrogen gas were purchased from Airgas Inc. and passed through a column containing manganese oxide supported on vermiculite and 4 Å molecular sieves before admission to the high vacuum line. Benzene- d_6 was purchased from Cambridge Isotope Laboratories and distilled from sodium metal under an atmosphere of argon and stored over 4 Å molecular sieves. Chloroform-d and dichloromethane- d_2 were purchased from Cambridge Isotope Laboratories and dried over CaH₂. Phosphines and phosphites were purchased from Aldrich, Strem, and Acros and were used as received. $Fe_2(Mes)_4^{24}$ was prepared according to the literature procedure.

¹H NMR spectra were recorded on a Varian Mercury 300 and Inova 400 and 500 spectrometers operating at 299.763, 399.780, and 500.62 MHz, respectively. All chemical shifts are reported relative to SiMe₄ using ¹H (residual) chemical shifts of the solvent as a secondary standard. For paramagnetic compounds, ¹H NMR data are reported with the chemical shift followed by the peak width at half-height in hertz followed by integration value and, where possible, peak assignment.

Unless stated otherwise, solution magnetic moments were measured at 22 °C by the method originally described by Evans²⁵ with stock and experimental solutions containing a known amount of a ferrocene standard. Solid-state magnetic moments were recorded using a Quantum Design SQUID magnetometer. Magnetization versus temperature data were recorded in a 5000 G applied field from 5 to 300 K. The sample holder was constructed from an NMR tube as previously described.²⁶ Diamagnetic contributions were estimated using Pascal's constants. The sample used was recrystallized several times and purity determined by elemental analysis.

Mössbauer data were recorded on an alternating constantacceleration spectrometer. The minimum experimental line width was 0.24 mm s⁻¹ (full width at half-height). A constant sample temperature was maintained with an Oxford Instruments Variox or an Oxford Instruments Mössbauer-Spectromag 2000 cyrostat. Reported isomer shifts (δ) are referenced versus iron metal at 293 K.

Single crystals suitable for X-ray diffraction were coated with polyisobutylene oil in a drybox and were quickly transferred to the goniometer head of a Siemens SMART CCD Area detector system equipped with a molybdenum X-ray tube ($\lambda = 0.71073$ Å). Preliminary data revealed the crystal system. A hemisphere routine was used for data collection and determination of lattice constants. The space group was identified and the data were processed using the Bruker SAINT program, and data were corrected for absorption using SADABS. The structures were solved using direct methods (SHELXS) completed by subsequent Fourier synthesis and refined by full-matrix least-squares procedures. Elemental analyses were performed by Robertson Microlit Laboratories, Inc., in Madison, NJ.

Preparation of (DMAP)Fe(Mes)₂ (1). A scintillation vial was charged with 0.102 g (0.173 mmol) of Fe₂(Mes)₄ and approximately 5 mL of diethyl ether. With stirring, 0.085 g (0.696 mmol) of DMAP in 3 mL of diethyl ether was added and the resulting reaction mixture stirred for 3 h. The resulting solid was collected by filtration, washed with pentane, and dried in vacuo yielding 0.157 g (84%) of 1 as a yellow powder. Anal. Calcd for C₃₀H₄₂FeN₄: C, 70.03; H, 8.23; N, 10.89. Found: C, 70.26; H, 8.14; N, 10.58. Magnetic susceptibility (benzene-*d*₆): $\mu_{eff} = 5.1 \ \mu_{B}$. ¹H NMR (benzene-*d*₆): $\delta = 4.59$ (59.3), 33.4 (241), 45.2 (814), 92.1 (546), 117 (452).

Preparation of (TMEDA)Fe(Mes)₂ (2). This molecule was prepared in a manner similar to that for 1 with 0.200 g (0.340 mmol) of Fe₂(Mes)₄ and 0.079 g (0.680 mmol) of TMEDA yielding 0.117 g (42%) of tan solid identified as 2. Anal. Calcd for C₂₄H₃₈FeN₂: C, 70.24; H, 9.33; N, 6.83. Found: C, 69.84; H, 9.24; N, 6.54. Magnetic susceptibility (benzene-*d*₆): $\mu_{\text{eff}} = 5.1 \ \mu_{\text{B}}$. ¹H NMR (benzene-*d*₆): $\delta = 2.13$ (68.3), 35.24 (1140), 85.17 (84.9), 113 (203).

Preparation of *trans*-(Me₃P)₂Fe(Mes)₂ (3). A 50 mL roundbottomed flask was charged with 0.200 g (0.340 mmol) of Fe₂-(Mes)₄ and approximately 10 mL of diethyl ether. With stirring, 0.103 g (1.40 mmol) of PMe₃ was added by a 31.6 mL calibrated gas bulb. The deep red solution was stirred for 2 h forming a yellow solution over time. The solvent was removed in vacuo, and the resulting yellow solid was washed with pentane and dried to afford

⁽²³⁾ Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J. *Organometallics* **1996**, *15*, 1518.

⁽²⁴⁾ Klose, A.; Solari, E.; Ferguson, R.; Floriani, C.; Chiesi-Villa, A.; Rizzoli, C. Organometallics 1993, 12, 2414.

⁽²⁵⁾ Sur, S. K. J. Magn. Reson. 1989, 82, 169.

⁽²⁶⁾ de Bruin, B.; Bill, E.; Bothe, E.; Weyhermüller, T.; Wieghardt, K. *Inorg. Chem.* **2000**, *39*, 2936.

Four Coordinate Iron(II) Complexes

0.254 g (84%) of **3**. Anal. Calcd for C₂₄H₄₀FeP₂: C, 64.58; H, 9.03. Found: C, 64.74; H, 9.05. Magnetic susceptibility (benzene- d_6): $\mu_{\rm eff} = 2.9 \ \mu_{\rm B}$. ¹H NMR (benzene- d_6): $\delta = -35.8$ (36, 18H, P(*CH*₃)₃), -4.8 (20, 6H, *CH*₃-*p*-C₆H₂), 20.5 (79, 12H, *CH*₃-*o*-C₆H₂), 25.5 (25, 4H, C₆H₂).

Preparation of *trans*-(**E**t₃**P**)₂**Fe**(**Mes**)₂ (4). This molecule was prepared in a manner similar to that for **1** with 0.250 g (0.430 mmol) of Fe₂(Mes)₄ and 0.201 g (1.7 mmol) of PEt₃ yielding 0.318 g (70%) of **4**. Anal. Calcd for C₃₀H₅₂FeP₂: C, 67.92; H, 9.88. Found: C, 68.82; H, 9.66. Magnetic susceptibility (benzene-*d*₆): $\mu_{eff} = 2.8$ μ_{B} . ¹H NMR (benzene-*d*₆): δ , -35.2 (845, 8H, CH₃CH₂), -19.8 (727.6, 12H, *CH*₃CH₂), -2.2 (1841, 6H, *CH*₃-*p*-C₆H₂), 10.4 (936.6, 12H, *CH*₃-*o*-C₆H₂), 27.6, (979.2, 4H, *m*-C₆H₂).

Preparation of *trans*-(**PhMe**₂**P**)₂**Fe**(**Mes**)₂ (5). A scintillation vial was charged with 0.200 g (0.340 mmol) of Fe₂(Mes)₄ and approximately 5 mL of diethyl ether. With stirring, 0.188 g (1.40 mmol) of PPhMe₂ in approximately 3 mL of diethyl ether was added at ambient temperature forming a deep red reaction mixture. An orange precipitate formed over the course of 3 h. The solid was collected by filtration and washed with pentane to afford 0.183 g (78%) of **5**. Anal. Calcd for C₃₄H₄₄FeP₂: C, 71.58; H, 7.77. Found: C, 71.53; H, 8.04. Magnetic susceptibility (benzene-*d*₆): $\mu_{\rm eff} = 2.7 \,\mu_{\rm B}$. ¹H NMR (benzene-*d*₆) δ = -33.5 (30.3, 12H, PC₆H₅-(*CH*₃)₂), -5.33 (5.3, 6H, *CH*₃-p-C₆H₂), 0.45 (14.3, 4H), 0.70 (23.4, 4H), 21.3 (74.6, 12H, *CH*₃-o-C₆H₂), 24.7 (11.1, 4H, C₆H₂).

Preparation of *trans*-(**PhEt₂P**)₂**Fe**(**Mes**)₂ (6). This molecule was prepared in a manner similar to that for **5** with 0.500 g (0.850 mmol) of Fe₂(Mes)₄ and 0.565 g (3.4 mmol) of PEt₂Ph in 10 mL of diethyl ether yielding 0.862 g (81%) of an orange powder identified as **6**. Anal. Calcd. for C₃₈H₅₂FeP₂: C, 72.84; H, 8.36. Found: C, 72.43; H, 7.97. Magnetic susceptibility (benzene-*d*₆): $\mu_{eff} = 2.7 \ \mu_{B}$. ¹H NMR (benzene-*d*₆): $\delta = -44.0 (232, 4H), -20.53 (151, 12H,$ *CH*₃-CH₂), -4.0 (368, 6H, C₆H₂-p-*CH*₃), -3.3 (495, 4H), 0.86 (197, 8H, CH₃*CH*₂), 9.1 (171, 2H, p-*H*-C₆H₅), 14.8 (262, 12H, C₆H₂-o-*CH*₃), 25.4 (185, 4H).

Preparation of *trans*-(**Ph**₂**MeP**)₂**Fe**(**Mes**)₂ (7). This molecule was prepared in a manner similar to that for **5** with 0.100 g (0.170 mmol) of Fe₂(Mes)₄ and 0.136 g (0.680 mmol) of PPh₂Me yielding 0.183 g (78%) of a red solid identified as **7**. Anal. Calcd for C₄₄H₄₈-FeP₂: C, 76.08; H, 6.96. Found: C, 75.69; H, 6.25. Magnetic susceptibility (benzene-*d*₆): $\mu_{eff} = 2.8 \,\mu_{B}$. ¹H NMR (benzene-*d*₆): $\delta = -7.0 \,(179, 8H), 2.2 \,(159, 12H, CH₃-o-C₆H₂), 10.4 (169, 6H), 17.9 (184, 4H), 21.3 (167, 4H), 23.4 (72.5, 6H).$

Preparation of [(EtO)₃P]₂Fe(Mes)₂ (8). This molecule was prepared in a manner similar to that for **5** with 0.235 g (0.390 mmol) of Fe₂(Mes)₄ and 0.332 g (2.0 mmol) of P(OEt)₃ yielding 0.287 g (57%) of a red solid identified as **8**. Magnetic susceptibility (benzene-*d*₆): $\mu_{\text{eff}} = 2.7 \ \mu_{\text{B}}$. ¹H NMR (benzene-*d*₆): $\delta = -8.2$ (43.6, 6H, *CH*₃-*p*-C₆H₂), -4.1 (213.2, 30H, P(O*CH*₂*CH*₃)), 19.0 (51.0, 4H, C₆*H*₂), 32.4 (137, 12H, *CH*₃-*o*-C₆H₂).

Preparation of (dmpe)Fe(Mes)₂ (9). This molecule was prepared in a manner similar to that for **5** with 0.047 g (0.080 mmol) of Fe₂(Mes)₄ and 0.024 g (0.160 mmol) of dmpe yielding 0.058 g (82%) of yellow solid identified as **9**. Anal. Calcd for C₂₄H₃₈FeP₂: C, 64.87; H, 8.62. Found: C, 64.89; H, 8.63. Magnetic susceptibility (Guoy balance): $\mu_{\text{eff}} = 4.2 \ \mu_{\text{B}}$.

Preparation of (depe)Fe(Mes)₂ (10). This molecule was prepared in a manner similar to that for **5** with 0.250 g (0.425 mmol) of Fe₂(Mes)₄ and 0.175 mg of depe yielding 0.332 g (79%) of an

orange solid identified as **10**. Anal. Calcd for $C_{28}H_{46}FeP_2$: C, 67.20; H, 9.26. Found: C, 67.01; H, 8.75. Magnetic susceptibility (benzene-*d*₆): $\mu_{eff} = 4.4 \, \mu_{B}$. ¹H NMR (benzene-*d*₆): $\delta = 9.1 \, (302, 12H, C_6H_2-o-CH_3)$, 18.4 (836, 6H, $C_6H_2-p-CH_3$), 90.3, -94.1 (2060, 20 H, CH_2CH_3), 127.9 (520, 4H, PCH_2CH_2P).

Preparation of (dppe)Fe(Mes)₂ (11). This molecule was prepared in a manner similar to that for **3** with 0.337 g (0.570 mmol) of Fe₂(Mes)₄ and 0.457 g (1.2 mmol) of dppe yielding 0.594 g (75%) of an orange solid identified as **11**. Anal. Calcd for C₄₄H₄₆-FeP₂: C, 76.30; H, 6.69. Found: C, 76.27; H, 6.31. Magnetic susceptibility (benzene-*d*₆): $\mu_{eff} = 3.7 \,\mu_{B}$. ¹H NMR (benzene-*d*₆) $\delta = -3.7 (120, 6H, CH₃-p-C_6H₂), 16.5 (153, 12H, CH₃-o-C_6H₂),$ 92.1 (2904), 97.6 (2880), 103 (163, 8H), 132 (386, 4H).

Preparation of (PhEt₂P)₂FeCl₂ (13). A 50 mL round-bottomed flask was charged with 0.363 g (2.90 mmol) of FeCl₂, 1.0 g (6.0 mmol) of PhEt₂P, and approximately 30 mL of toluene. A reflux condenser and a 180° needle valve were attached, and the reaction mixture was heated to reflux for 2 h. The reaction mixture was transferred into the drybox and filtered through Celite. The toluene was removed in vacuo, and the off-white residue was washed several times with pentane and dried to afford 1.23 g (93%) of **13**. Magnetic susceptibility (benzene-*d*₆): $\mu_{eff} = 4.8 \ \mu_{B}$. ¹H NMR (benzene-*d*₆): $\delta = -2.6 (57.4, 2H, p-C_6H_5), 2.57 (435.5, 4H), 14.1 (61.5, 4H), 77.9 (1543, 20 H, C₂H₅).$

Preparation of (PhEt₂P)₂FeBr₂ (14). This molecule was prepared in a manner similar to that for **13** with 0.156 g (0.720 mmol) of FeCl₂, 0.250 g (1.5 mmol) of PhEt₂P, and approximately 20 mL of toluene and afforded 0.302 mg (77%) of **14**. Magnetic susceptibility (benzene- d_6): $\mu_{eff} = 5.0 \,\mu_B$. ¹H NMR (benzene- d_6): $\delta = -3.4$ (52.3, 2H, p-C₆ H_5), 4.6 (356.2, 4H), 14.0 (52.6, 4H), 77.1 (648.6, 20 H, C₂ H_5).

Procedure for a Typical Exchange Reaction. A J. Young NMR tube was charged with 0.020 g (0.035 mmol) of **5** and approximately 1.0 mL of benzene- d_6 . Using a calibrated gas bulb, 2 equiv (0.070 mmol) of PMe₃ was added at 77 K. The solution was thawed and shaken and the ¹H NMR spectrum recorded.

¹H NMR Characterization of (py)₂Fe(Mes)₂. ¹H NMR (benzened₆): $\delta = -5.14$ (77.2, CH, py), 23.40 (10.8, CH, *p*-py), 23.4 (127.1, CH, py), 41.6 (752.3, CH₃, *o*-C₆H₂), 112.5 (168.2, CH₃, *p*-C₆H₂), 138 (270.6, CH, *m*-C₆H₂).

¹H NMR Characterization of (PhEt₂P)₂Fe(C₆Cl₅)₂. ¹H NMR (benzene- d_6): $\delta = -53.0$ (29.5, Ph), -18.5 (19.1, CH_3), -7.9 (9.2, Ph), 11.8 (16.6, *p*-Ph).

Acknowledgment. We thank the Cornell Department of Chemistry and Chemical Biology and the National Science Foundation (CAREER Award to P.J.C.) for financial support. P.J.C. also acknowledges the Research Corp. (Cottrell Scholarship) and the Packard Foundation for a fellowship in science and engineering. G.T.Y. acknowledges the National Science Foundation (CHE-023488) and the Virginia Tech ASPIRES program for each providing partial support for the purchase of the SQUID magnetometer.

Supporting Information Available: Crystallographic data for *trans*-(PhEt₂P)₂Fe(C₆Cl₅)₂, **3**, **6**, **9**, and **11** (CIF, PDF). This material is available free of charge via the Internet at http://pubs.acs.org. IC048202+