

Neutron diffraction study of the highly distorted octahedral complex $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OPh})_3]_2$

Nam Nhat Ho^a, Robert Bau^{a,*}, Sax A. Mason^b

^a Department of Chemistry, University of Southern California, Los Angeles, CA 90089, USA

^b Institut Laue-Langevin, Avenue des Martyrs, F-38042, Grenoble, France

Received 6 February 2003; received in revised form 17 April 2003; accepted 17 April 2003

Abstract

A neutron diffraction analysis was carried out at 20 K on a single crystal of $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OPh})_3]_2$ in order to accurately locate the hydride ligands. The two hydrides are bonded terminally to the Fe atom, and their small size causes the title complex to assume a highly distorted octahedral geometry. Distances and angles involving the hydride ligands are as follows: $\text{Fe}-\text{H}(1) = 1.521(2) \text{ \AA}$, $\text{Fe}-\text{H}(2) = 1.529(2) \text{ \AA}$, $\text{H}(1)\cdots\text{H}(2) = 2.011(3) \text{ \AA}$, and $\text{H}(1)-\text{Fe}-\text{H}(2) = 82.49(13)^\circ$. Final agreement factors: $R(F) = 2.8\%$ and $R(wF^2) = 5.4\%$.

© 2003 Elsevier Science B.V. All rights reserved.

Keywords: Neutron diffraction; Iron hydride; *cis*-Dihydride complexes

1. Introduction

Complexes of the type FeH_2L_4 have had a long history. Although $\text{FeH}_2(\text{CO})_4$ has been known since 1931 when it was first synthesized by Hieber and Leutert [1], there were some initial doubts about the locations of the hydride ligands. Some early investigators proposed a tetrahedral structure for $\text{FeH}_2(\text{CO})_4$, postulating that the hydrogen atoms might be bonded to the oxygen atoms of the carbonyl ligands [2]. Eventually, an electron diffraction study [3] laid all speculations to rest by demonstrating that $\text{FeH}_2(\text{CO})_4$ has a highly distorted *cis*-octahedral geometry, with a stereochemically-active hydride ligand and normal Fe–H bonds of length $1.56(2) \text{ \AA}$. However, in that study the precision of the Fe–H distance measurement was rather low because of the inherent limitations of the electron diffraction technique. In this paper, we describe the results of a single-crystal neutron diffraction study on the related compound $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OPh})_3]_2$, in which the Fe–H distances are measured with much higher precision.

2. Experimental

2.1. Preparation of $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OPh})_3]_2$

The title compound was prepared according to the method published by Brunet et al. [4]. In a typical experiment, an 80 ml solution of $\text{K}[\text{FeH}(\text{CO})]_4$ (11 mmol) in THF:H₂O (30:50 ml) is prepared under argon; to this solution, triphenyl phosphite (6.83 g, 22 mmol) is added. An evolution of CO gas is observed, after which (24 h later) evaporation of THF under reduced pressure leads to the formation of an aqueous phase and the concurrent formation of a white precipitate. After water is removed, the microcrystalline product is washed with water previously distilled under argon. Attempts to grow large crystals using a variety of different conditions and solvents were tried, and we found that the best quality crystals could be grown by allowing hexane to diffuse slowly into a THF solution of the title compound at -20°C .

2.2. Neutron diffraction analysis

A large, transparent, colorless crystal with dimension of $4.3 \times 2.4 \times 1.0 \text{ mm}$ was used in the neutron analysis. The sample was mounted in air onto a 1-mm diameter

* Corresponding author. Tel.: +1-213-740-2692; fax: +1-213-740-0930.

E-mail address: bau@usc.edu (R. Bau).

vanadium pin using epoxy resin. Data were collected on instrument D19 at the Institut Laue Langevin equipped with a Displex cryostat and a curved position-sensitive area detector [5], to which was added a new ‘square’ (192 × 192 mm) position-sensitive microstrip detector [6]. The crystal was cooled at a rate of 2 K min^{−1} to 20 K while the intensity and peak profile of a strong reflection was monitored, and no change in the mosaicity of that reflection was observed. The space group was confirmed to be $P\bar{1}$ at 20 K. During data collection, three standard reflections were monitored regularly and showed no significant variation in intensity. A total of 15 316 reflections were collected and merged to yield 6402 independent reflections with positive F^2 values [7]. After an analytical absorption correction with the program D19ABS, based on the ILL version of the CCSL program system [8], phasing of the neutron data was carried out using the atomic coordinates of the non-hydrogen atoms obtained from a previous X-ray analysis [9,10]. Least-squares refinement of all atomic coordinates, and anisotropic temperature factors, resulted in final agreement factors of $R(F) = 2.8\%$ and $R(wF^2) = 5.4\%$ for all reflections. Further details of the data collection and refinement are shown in Table 1.

Table 1
Crystal data and structure refinement for $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OPh})_3]_2$

Empirical formula	$\text{C}_{38}\text{H}_{32}\text{FeO}_8\text{P}_2$
Formula weight	732.00
Temperature (K)	20(2)
Wavelength (Å)	1.3150
Crystal system	Triclinic
Space group	$P\bar{1}$
Unit cell dimensions	
<i>a</i> (Å)	9.5682(4)
<i>b</i> (Å)	10.3878(4)
<i>c</i> (Å)	18.2514(7)
α (°)	102.323(2)
β (°)	94.527(2)
γ (°)	107.112(2)
<i>V</i> (Å ³)	1674.16(9)
<i>Z</i>	2
<i>D</i> _{calc} (Mg m ^{−3})	1.452
Absorption coefficient (cm ^{−1})	1.720
Crystal size (mm)	4.3 × 2.4 × 1.0
θ range for data collection (°)	2.14–61.68
Index ranges	$-12 \leq h \leq 10$, $-7 \leq k \leq 13$, $-23 \leq l \leq 22$
Reflections collected	15316
Independent reflections	6402 [$R_{\text{int}} = 0.0217$]
Refinement method	Full-matrix least-squares on F^2
Data/restraints/parameters	6402/0/733
Goodness-of-fit on F^2	1.409
Final <i>R</i> indices [$I > 2\sigma(I)$]	$R_1 = 0.0264$, $wR_2 = 0.0540$
<i>R</i> indices (all data)	$R_1 = 0.0276$, $wR_2 = 0.0543$
Extinction coefficient	0.00145(6)
Largest difference peak and hole (fm Å ^{−3})	0.372 and −0.379




Fig. 1. Molecular structure of $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OPh})_3]_2$ at 20 K, with hydrogen atoms in the phenyl groups removed for clarity. Ellipsoids are shown at the 50% probability level.

In Fig. 1, the structure of $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OPh})_3]_2$ is shown with all aryl hydrogen atoms omitted for clarity. Selected distances and angles in the molecule are listed in Table 2. Significantly, this is the most precise structural characterization of a H_2FeL_4 -type complex to date, with extremely low esd's for the Fe–H distances (1.521(2) and 1.529(2) Å) and the H··H distance (2.011(3) Å). Part of the motivation for this ultra-high-resolution study was that we wanted to confirm the capabilities of a new square detector recently installed at the D19 beam line [6]. This microstrip detector proved

Table 2
Selected bond distances (Å) and angles (°) in $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OPh})_3]_2$

Bond lengths (Å)	
Fe(1)–P(2)	2.099(2)
Fe(1)–P(1)	2.119(2)
Fe(1)–C(1)	1.787(1)
Fe(1)–C(2)	1.798(1)
Fe(1)–H(1)	1.520(2)
Fe(1)–H(2)	1.529(2)
H(1)··H(2)	2.011(3)
Bond angles (°)	
C(1)–Fe(1)–C(2)	100.90(6)
H(1)–Fe(1)–H(2)	82.49(13)
P(2)–Fe(1)–P(1)	154.41(5)
C(2)–Fe(1)–P(2)	103.37(6)
C(2)–Fe(1)–P(1)	96.08(6)
C(1)–Fe(1)–P(1)	94.81(7)
C(1)–Fe(1)–P(2)	97.62(7)
C(1)–Fe(1)–H(1)	89.29(10)
C(2)–Fe(1)–H(1)	169.81(10)
C(1)–Fe(1)–H(2)	171.77(10)
C(1)–Fe(1)–H(1)	87.32(10)
P(2)–Fe(1)–H(1)	74.54(10)
P(1)–Fe(1)–H(1)	83.37(10)
P(2)–Fe(1)–H(2)	80.84(11)
P(1)–Fe(1)–H(2)	83.67(10)

to be very reliable and stable, with resolution (3×3 mm) well-matched to neutron experiments.

3. Discussion

In this report, we describe an accurate analysis of $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OC}_6\text{H}_5)_3]_2$ that confirms a distorted octahedral geometry with the carbonyl ligands in mutually *cis* positions and the phosphite ligands in mutually *trans* positions. In an earlier X-ray analysis of this complex, the following distances and angles involving the hydride ligands were reported: $\text{Fe}-\text{H} = 1.44(2), 1.42(2)$ Å; $\text{H} \cdots \text{H} = 1.81(1)$ Å; $\text{H}-\text{Fe}-\text{H} = 79.1(13)^\circ$ [10]. Our more precise neutron diffraction study gives the following molecular parameters: $\text{Fe}-\text{H} = 1.520(2), 1.529(2)$ Å; $\text{H} \cdots \text{H} = 2.011(3)$ Å; $\text{H}-\text{Fe}-\text{H} = 82.49(13)^\circ$. The differences confirm the usual finding [11] that $\text{M}-\text{H}$ distances derived from X-ray data are generally shorter than their true values, because during least-squares refinement the H position in an X-ray analysis is often drawn in towards the electron density of the heavy atom.

Various bond distances and angles in selected *cis*-dihydride iron complexes are summarized in Table 3. Perhaps the most striking feature in this table is the highly distorted value of the *trans*- LFeL angle, which ranges from $136.70(6)^\circ$ in $\text{FeH}_2[\text{PPh}(\text{OEt})_2]_4$ [12] to $154.41(5)^\circ$ in the title complex, far less than the 180° expected for an ideal octahedral structure. Curiously, the extent of this distortion appears to be independent of the steric bulk of the L ligand, which varies from small (CO) to large ($\text{P}(\text{OPh})_3$). In other words, the FeL_4 portion of the FeH_2L_4 skeleton in virtually all cases seems to be midway between octahedral and tetrahedral. This observation is rather puzzling because other six-coordinate hydrido complexes that have been analyzed by neutron diffraction, such as $\text{MnH}(\text{CO})_5$ [13] and *fac*-

$\text{IrH}_3(\text{PMePh}_2)_3$ [14], do not show anywhere near that amount of distortion from octahedral symmetry.


The H atoms in $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OPh})_3]_2$ make a slightly acute $\text{H}-\text{Fe}-\text{H}$ angle of $82.5(1)^\circ$, and hence can be considered very slightly ‘compressed’ as compared to that in an ideal octahedral complex. Nevertheless, the corresponding $\text{H} \cdots \text{H}$ distance of $2.011(3)$ Å is definitely in the non-bonding range, very far from the $1.5\text{--}1.6$ Å distance that marks the beginning of incipient $\text{H}-\text{H}$ bond formation [11b,15,16]. Originally, we were speculating that the large steric bulk of the triphenylphosphite ligands might force the hydride ligands to assume an unusually close $\text{H} \cdots \text{H}$ distance, such as the $1.357(7)$ Å found in $\text{ReH}_7[\text{P}(\text{tolyl})_3]_2$ [15] and $1.49(4)$ Å in $[\text{OsH}_5(\text{PMePh}_2)_3]^+$ [16]. However, that turned out not to be the case.

Finally, as far as the $\text{Fe}-\text{H}$ distance itself is concerned, the values of $1.521(2)$ and $1.529(2)$ Å we find in $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OPh})_3]_2$ agree very well with other terminal $\text{Fe}-\text{H}$ bond lengths measured via neutron diffraction: $1.514(6)$ and $1.538(7)$ Å in $\text{FeH}_2(\eta^2\text{-H}_2)(\text{PPh}_2\text{Et})_3$ [17] and $1.535(12)$ Å in $[\text{Fe}(\text{H})(\eta^2\text{-H}_2)(\text{dppe})]^+[\text{BPh}_4]^-$ [18]. From all of these results, we can estimate an average value of 1.53 Å for a covalent $\text{Fe}-\text{H}$ single bond, a number which fits very well along the following trend for first-row transition metals: $\text{Mn}-\text{H} = 1.59$ Å, $\text{Fe}-\text{H} = 1.53$ Å, $\text{Co}-\text{H} = 1.52$ Å, and $\text{Ni}-\text{H} = 1.45$ Å [11a]. We also note that this distance is slightly but significantly shorter than $\text{Fe}-\text{H}$ distances involving non-classical dihydrogen ligands, $\eta^2\text{-H}_2$, which have been measured to be in the $1.58\text{--}1.61$ Å range [17,18].

4. Supplementary material

Crystallographic data for the structural analysis have been deposited with the Cambridge Crystallographic

Table 3
Bond distances and angles in selected iron *cis*-dihydride complexes



	L	L*	$\text{Fe}-\text{H}$ (Å)	$\text{H}-\text{Fe}-\text{H}$ ($^\circ$)	$\text{H} \cdots \text{H}$ (Å)	$\text{L}-\text{Fe}-\text{L}$ ($^\circ$)	$\text{L}^*-\text{Fe}-\text{L}^*$ ($^\circ$)	Method ^a	References
$\text{FeH}_2(\text{CO})_4$	CO	CO	1.56(2)	100(10)	2.39(2)	148.5(15)	96.0(6)	E	[3]
$\text{FeH}_2(\text{CO})_2[\text{P}(\text{OPh})_3]_2$	$\text{P}(\text{OR})_3$	CO	1.525(2) (avg.)	82.5(1)	2.011(3)	154.41(5)	100.90(6)	N	This work
$\text{FeH}_2[\text{PPh}(\text{OEt})_2]_4$	$\text{PR}(\text{OR}')_2$	$\text{PR}(\text{OR}')_2$	1.51(4) (avg.)	89(2)	2.12(5)	136.70(6)	102.28(6)	X	[12]
$\text{FeH}_2(\eta^2\text{-H}_2)(\text{PPh}_2\text{Et})_3$	PR_3	PR_3, H_2	1.526(7) (avg.)	88.2(3)	2.124(7)	149.8(2)		N	[17]
$\text{FeH}_2(\text{N}_2)(\text{PPh}_2\text{Et})_3$	PR_3	PR_3, N_2	1.43(4) (avg.)	79(2)	1.82(4)	148.9(1)	97.2(2)	X	[18]
$\text{FeH}_2(\text{CO})_3(\text{AsPh}_3)$	CO	AsPh_3, CO	1.35 (avg.)	84	1.81	151.6(3)	100.3(3)	X	[19]

^a Experimental technique: E, electron diffraction; N, neutron diffraction; X, X-ray diffraction.

Data Center, CCDC no. 208347 for compound $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OC}_6\text{H}_5)_3]_2$. Copies of this information may be obtained free of charge from The Director, CCDC, 12 Union Road, Cambridge, CB2 1EZ UK (Fax: +44-1223-336033; e-mail: deposit@ccdc.cam.ac.uk or [www: http://www.ccdc.cam.ac.uk](http://www.ccdc.cam.ac.uk)).

Acknowledgements

This research was supported by the National Science Foundation Grant CHE-98-16294 and NATO Grant CRG-921317.

References

- [1] W. Hieber, F. Leutert, *Naturwissenschaften* 19 (1931) 360.
- [2] R.V.G. Ewens, M.W. Lister, *Trans. Faraday Soc.* 35 (1939) 681.
- [3] E.A. McNeill, F.R. Scholer, *J. Am. Chem. Soc.* 99 (1977) 6243.
- [4] J.-J. Brunet, F.B. Kindela, D. Neibecker, *Inorg. Syn.* 29 (1992) 156.
- [5] M. Thomas, R.F.D. Stansfield, M. Berneron, A. Filhol, G. Greenwood, J. Jacobe, D. Feltin, S.A. Mason, in: P. Convert, J.B. Forsyth (Eds.), *Position Sensitive Detection of Thermal Neutrons*, Academic Press, London, 1983, p. 344.
- [6] F.G. Clergeau, P. Convert, D. Feltin, H.E. Fischer, B. Guerard, T. Hansen, G. Manzin, A. Oed, P. Palleau, *IEEE Trans. Nucl. Sci.* 48 (2001) 1075.
- [7] Neutron data reduction was carried out using the program RETREAT C. Wilkinson, H.W. Khamis, R.F.D. Stansfield, G.J. McIntyre, *Appl. Crystallogr.* 21 (1988) 471.
- [8] J.C. Matthewman, P. Thompson, P.J. Brown, *J. Appl. Cryst.* 15 (1982) 167.
- [9] R. Bau, N.N. Ho, unpublished results.
- [10] V. Arion, J.-J. Brunet, D. Neibecker, *Inorg. Chem.* 40 (2001) 2628.
- [11] (a) R. Bau, R.G. Teller, *Struct. Bond.* 44 (1981) 1;
(b) R. Bau, M.H. Drabnis, *Inorg. Chim. Acta* 259 (1997) 27.
- [12] L.J. Guggenberger, D.D. Titus, M.T. Flood, R.E. Marsh, A.A. Orio, H.B. Gray, *J. Am. Chem. Soc.* 94 (1972) 1135.
- [13] S.J. LaPlaca, W.C. Hamilton, J.A. Ibers, A. Davison, *Inorg. Chem.* 8 (1969) 1928.
- [14] R. Bau, C.J. Schwerdtfeger, L. Garlaschelli, T.F. Koetzle, *J. Chem. Soc. Dalton* (1993) 3359.
- [15] L. Brammer, J.A.K. Howard, O. Johnson, T.F. Koetzle, J.L. Spencer, M. Andre, *Chem. Commun.* 4 (1991) 241.
- [16] T.J. Johnson, A. Albinati, T.F. Koetzle, J. Ricci, O. Eisenstein, J.C. Huffman, K.G. Caulton, *Inorg. Chem.* 33 (1994) 4966.
- [17] L.S. Van Der Sluys, J. Eckert, O. Eisenstein, J.H. Hall, J.C. Huffman, S.A. Jackson, T.F. Koetzle, G.J. Kubas, P.J. Vergamini, K.G. Caulton, *J. Am. Chem. Soc.* 112 (1990) 4831.
- [18] J.S. Ricci, T.F. Koetzle, M.T. Bautista, T.M. Hofstede, R.H. Morris, J.F. Sawyer, *J. Am. Chem. Soc.* 11 (1989) 8823.
- [19] H. Berke, G. Huttner, L. Zsolnai, *Chem. Ber.* 114 (1981) 3549.