

Iron(IV) Porphyrins from Iron(III) Porphyrin Cation Radicals

John T. Groves,^{*a} Robert Quinn,^a Thomas J. McMurry,^a George Lang,^{*b} and Brian Bosob

^a Department of Chemistry, The University of Michigan, Ann Arbor, Michigan 48109, U.S.A.

^b Department of Physics, Pennsylvania State University, University Park, Pennsylvania 16802, U.S.A.

The preparation and spectroscopic data of FeTMP(ClO₄)₂ [(1), TMP = 5,10,15,20-tetramesitylporphyrinato] are reported; the reaction of (1) with NaOMe–MeOH gave a new red species, (2), which is indicated to exist (a) as a symmetrically ligated complex with a vacant d_{x²–y²} iron orbital (¹H n.m.r.) and (b) in the iron(IV) state (Mössbauer spectroscopy).

The importance of oxidized iron–porphyrin complexes in heme-containing enzymes is now fully appreciated.^{1–3} However, the electronic structures and mechanistic interrelationships of such species have remained obscure owing to the lack of simple chemical examples. Recently, at least three types of iron(III) porphyrin cation radical complexes have been recognized.^{4–8} Further, an oxo-iron(IV)⁹ and an oxo-iron(IV)¹⁰ porphyrin cation radical^{10,11} have been characterized. Only the oxo-,¹² nitrido-, and carbido-¹³ ligands are known to stabilize the iron(IV) state with respect to the iron(III) porphyrin cation radical.¹⁴ We report here the first direct conversion of an iron(III) porphyrin cation radical into an iron(IV) porphyrin by ligand metathesis.

The treatment of perchlorato-5,10,15,20-tetramesitylporphyrinatoiron(III) [FeTMP(ClO₄)₂] with an excess of iron(III) perchlorate in methylene chloride at room temperature resulted in the formation of a bright green solution. Removal of the solvent and recrystallization of the resulting solid from toluene–hexane afforded crystalline material which gave a satisfactory elemental analysis for FeTMP(ClO₄)₂, (1). Solutions of (1) in toluene or methylene chloride were stable for short periods at room temperature; however, immediate reduction of (1) occurred in methanol or acetone at room temperature. The spectroscopic properties of (1) were similar to other iron(III) porphyrin cation radical complexes.^{5–7} The visible spectrum of (1) displayed a broad absorbance at 800 nm and other absorbances as shown in Figure 1(a). The infrared spectrum of (1) showed an intense band at 1270 cm^{–1}¹⁵ and there were no apparent e.s.r. signals for (1) at 110 K. The ¹H n.m.r. spectrum of (1) at 25 °C in CD₂Cl₂ was similar to that of Fe^{III}TPP(ClO₄)₂^{5,7} (TPP = tetraphenylporphyrinato) (pyrrole H, δ 18.7; *meta* H, 55; *o*-methyl, 20; *p*-methyl, 12.5). The assignment of the pyrrole and *meta* protons was confirmed by comparison of the spectra of the corresponding deuteriated porphyrins. The titration of (1) with one equivalent of tetrabutylammonium iodide regenerated FeTMP(ClO₄). This result and the spectroscopic data are consistent with a one-electron oxidation of the porphyrin ring.

The addition of two equivalents of sodium methoxide in methanol to CD₂Cl₂ solutions of (1) at –50 °C caused the immediate production of a bright red species (2) with a visible spectrum dramatically different from that of (1) [Figure 1(b)].

The oxidation of FeTMP(OH)¹⁰ with iodosylmesitylene (Mes-IO) in methylene chloride–methanol also produced (2) but the addition of methanol to (1) did not. The 360 MHz ¹H n.m.r. spectrum of (2) in CD₂Cl₂ at –78 °C shows a pyrrole resonance at very high field (δ –37.5) and other resonances near the expected diamagnetic positions (*meta* H, 7.72, s; *p*-methyl, 2.86, s; *o*-methyl, 2.4, br.). A broad resonance in the spectrum of (2) prepared in methanol at δ 3.6 which was absent in samples prepared from CD₃OD was assigned to co-ordinated methoxide. The general features of this spectrum are similar to those reported recently for low spin, six-co-ordinate manganese(III) porphyrin complexes.¹⁶

The e.s.r. spectrum of (2) shows only weak signals due to high-spin iron(III). The Mössbauer spectrum of ⁵⁷Fe-(2) (δ_{Fe} –0.025 mm/s, Δ*E*_Q 2.1 mm/s) is unlike those reported for





Figure 1. (a) Visible absorption spectrum for (1) in CH₂Cl₂ at 25 °C; (b) visible absorption spectrum of (2), 1.28 × 10^{–5} M in CH₂Cl₂–methanol at –40 °C.




Figure 2. Comparative Mössbauer data for iron porphyrins; ■ high-spin Fe^{III}, □ low-spin Fe^{III}, ● intermediate spin (admixed) Fe^{III}, ○ high spin Fe^{II}, △ low-spin Fe^{II}, ▲ oxo- and imido-Fe^{IV}, * (2). Cf. refs. 11, 17, and 18. Inset shows the zero field Mössbauer spectrum of ⁵⁷Fe-(2) at 4.2 K.

iron(III) porphyrins and similar to the relatively few known iron(IV) species (Figure 2).^{11,17,18} The sharp quadrupole doublet for (2) (Figure 2 inset) was unaffected by changes in temperature or a small transverse magnetic field indicating a non-Kramers spin system.

Taken together these results indicate that the co-ordination of methoxide, a strong π -donating ligand, has shifted the site of oxidation from the porphyrin ring in (1) to the iron in (2).

Support of this research by the National Institutes of Health (to J. T. G. and to G. L.) is gratefully acknowledged. Fellowship support for R. Q. was provided by the Dow Chemical Company.

Received, 12th June 1984; Com. 816

References

- W. D. Hewson and L. P. Hager, in 'The Porphyrins,' ed. D. Dolphin, Academic Press, New York, 1979, Vol. VIII, pp. 295–332.
- H. B. Dunford and J. S. Stillman, *Coord. Chem. Rev.*, 1976, **19**, 187.
- R. H. Felton, G. S. Owen, D. Dolphin, and J. Fajer, *J. Am. Chem. Soc.*, 1971, **93**, 6332.
- P. Gans, J.-C. Marchon, C. A. Reed, and J.-R. Regnard, *Nouv. J. Chim.*, 1981, 203.
- G. Buisson, A. Deronzier, E. Duee, P. Gans, J.-C. Marchon, and J.-R. Regnard, *J. Am. Chem. Soc.*, 1984, **104**, 6793.
- M. A. Phillippi and H. A. Goff, *J. Am. Chem. Soc.*, 1982, **104**, 6026.
- W. F. Scholz, C. A. Reed, Y. J. Lee, W. R. Scheidt, and G. Lang, *J. Am. Chem. Soc.*, 1982, **104**, 6791.
- H. M. Goff and M. A. Phillippi, *J. Am. Chem. Soc.*, 1983, **105**, 7567.
- D. H. Chin, G. N. La Mar, and A. L. Balch, *J. Am. Chem. Soc.*, 1980, **102**, 4344.
- J. T. Groves, R. C. Haushalter, M. Nakamura, T. E. Nemo, and B. J. Evans, *J. Am. Chem. Soc.*, 1981, **103**, 2884.
- J. E. Penner-Hahn, T. J. McMurphy, M. Renner, L. Latos-Grazynsky, K. S. Eble, I. M. Davis, A. L. Balch, J. T. Groves, J. H. Dawson, and K. O. Hodgson, *J. Biol. Chem.*, 1983, **258**, 12761.
- W. Hiller, J. Strähle, A. Datz, M. Hanack, W. E. Hatfield, L. W. ter Haar, and P. Gülich, *J. Am. Chem. Soc.*, 1984, **106**, 329.
- K. M. Kadish, R. K. Rhodes, L. A. Bottomley, and H. M. Goff, *Inorg. Chem.*, 1981, **20**, 3195; D. R. English, D. N. Hendrickson, and K. S. Suslick, *ibid.*, 1982, **22**, 367.
- F. Arena, P. Gans, and J.-C. Marchon, *J. Chem. Soc., Chem. Commun.*, 1984, 196.
- E. T. Shimomura, M. A. Phillippi, H. M. Goff, W. F. Scholz, and C. A. Reed, *J. Am. Chem. Soc.*, 1981, **103**, 6778.
- A. P. Hanson and H. M. Goff, *Inorg. Chem.*, 1984, in the press. We thank Professor Goff for this information prior to publication.
- B. Boso, G. Lang, T. J. McMurphy, and J. T. Groves, *J. Chem. Phys.*, 1983, 1122 and references therein.
- G. Simonneaux, W. F. Scholz, C. A. Reed, and G. Lang, *Biochim. Biophys. Acta*, 1982, **716**, 1.