This article was downloaded by: [Carnegie Mellon University] On: 20 January 2015, At: 03:11 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides and Nucleotides

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lncn19</u>

Synthesis and Characterization of Short Oligonucleotide Segments Containing Nonnatural Internucleoside Amine- and Amide Linkages

Gorm Vang Petersen ^a & Jesper Wengel ^a

^a Department of Chemistry , University of Odense , DK-5230, Odense M, Denmark

Published online: 16 Feb 2007.

To cite this article: Gorm Vang Petersen & Jesper Wengel (1995) Synthesis and Characterization of Short Oligonucleotide Segments Containing Nonnatural Internucleoside Amine- and Amide Linkages, Nucleosides and Nucleotides, 14:3-5, 925-928

To link to this article: http://dx.doi.org/10.1080/15257779508012503

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

SYNTHESIS AND CHARACTERIZATION OF SHORT OLIGONUCLEOTIDE SEGMENTS CONTAINING NONNATURAL INTERNUCLEOSIDE AMINE- AND AMIDE LINKAGES

Gorm Vang Petersen and Jesper Wengel*

Department of Chemistry, University of Odense, DK-5230 Odense M, Denmark

Abstract: The synthesis of thymidine dimers in which the phosphodiester linkage has been replaced by either piperazine or N,N'-dimethylethylenediamine are described. The dimers containing piperazine were incorporated into oligodeoxynucleotides on which thermal and enzymatic stability experiments were performed.

Modified oligonucleotides have achieved much attention because of their potential use as therapeutics in the treatment of AIDS and certain forms of cancer. The oligonucleotides can be modified in the nucleobase, the carbohydrate moiety or the internucleoside linkage. The oligonucleotides containing modified linkages can be divided into two classes, one in which the phosphor atom is retained and one in which the natural phosphodiester linkage is replaced by achiral, neutral moieties.¹ Here we wish to report the synthesis of the four novel diamine linked thymidine dimers **6**, **8**, **10** and **11** and the incorporation of **7** and **9** into oligonucleotide sequences. The thermal and enzymatic stability of these oligodeoxynucleotides were evaluated.

The key synthons 1 and 2 were synthesized by refluxing 1-[2-deoxy-5-O-(4,4'dimethoxytrityl)-3-O-methanesulfonyl- β -D-*threo*-pentofuranosyl]thymine prepared from 5'-O-(4,4'-dimethoxytrityl)thymidine using the protocol outlined by Fox and Miller² with either piperazine or N,N'-dimethylethylenediamine. Thus affording 1 in 30% and 2 in 19% yield after chromatographic purification. The 5'-aldehyde 3³ and 5'-carboxylic acid 4⁴ were prepared from thymidine as previously reported. The dimers 5 and 10 were synthesized by reductive amination of 3 with either 1 or 2 using titanium tetraisopropoxide in toluene and sodium cyanoborohydride giving 5 in 33% and 10 in 25% yield. 6 was obtained by treating 5 with tetrabutylammoniumfluoride (TBAF) in THF in 46% yield. The amide linkage in 8 and 11 was formed by dicyclohexylcarbodiimide (DCC)/Nhydroxysuccinimide (NHS) activation of 4 followed by addition of either 1 or 2 furnishing 8 and 11 in 60% and 73% yield, respectively. The two phosphoramidites 7 and 9 were obtained by reacting 6 and 8 with 2-cyanoethyl-N,N-diisopropylphosphoramidochloridite in the presence of N,N-diisopropylethylamine hence giving 7 and 9 in yields of 84% and 82% after precipitation from hexanes.



 $\begin{aligned} Reagents: (i) \ 1) \ Ti(OiPr)_4 (toluene/3A \ molecular \ sieves 2) \ NaBH_3CN, (ii) \ NHS/DCC/DMF, (iii) \ TBAF/THF \\ (iv) \ NCCH_2CH_2OP(Cl)NiPr_2/EtNiPr_2/CH_2Cl_2. \ DMT=4,4'-dimethoxytrityl, \ T=thymin-1-yl. \end{aligned}$

The phosphoramidites 7 and 9 were incorporated into different oligonucleotide sequences by standard phosphoramidite methodology on an automated DNA-synthesizer⁵ in an average yield of 50%.

The thermal stability of the duplexes formed by the modified oligodeoxynucleotides and their DNA complements were determined by melting experiments as previously described.⁶ It can be seen that incorporation of 7 once or twice in the middle of an oligodeoxynucleotide results in a considerable lowering of the melting temperature T_m by approximately 11 °C per modification. Contrary to this, incorporation of the amide 9 once or twice in the middle only causes a slight decrease in T_m (approximately 2 °C per modification).

	n nhannka	T_m (°C) / h_T	ΔT_{m} (°C)		h _T	
	Sequence $(5' \rightarrow 3')$	w.t.	8	11	8	11
A	CACCAACT*TCTTCCACA	64/1.34	-10.8	-1.6	1.25	1.28
B	CACCAACT*TCT*TCCACA	64/1.34	-11.1	-2.0	1.23	1.26
С	TTAACTTCTTCACAT*TC	55/1.34	-2.8	-2.4	1.33	1.34
D	T*TAACTTCTTCACAT*TC	55/1.34	-2.0	-1.6	1.31	1.30

Table 1. Oligodeoxynucleotides Synthesized and Melting Experiments

 T^*T = modified thymidine dimer, w.t. = wild type DNA, ΔT_m = decrease in T_m per modification,

 h_{T} = thermal hyperchromicity

As expected, incorporation of the modified dimers 7 and 9 at the ends do not lead to a significant change in T_m values. The hyperchromicities of all the modified oligodeoxy-nucleotides deviate to some extent from the wild type oligodeoxynucleotides indicating a distortion of the secondary structure of the DNA duplexes formed compared to the wild type duplexes, the distortion being less for the end-modified oligomers as expected.

The fact that 3'-phoshpodiesterases play a predominant role in the *in vivo* degradation of natural oligodeoxynucleotides⁷ prompted us to study the enzymatic stability of the modified oligodeoxynucleotides towards the 3'-exonuclease snake venom phosphodiesterase (SV PDE). Incorporation of 7 and 9 gave rise to an increase in stability towards 3'-exonucleases by a factor of 5 to 6, while incorporation in the middle lead to a rapid degradation of the 17-mers to a stabilized 9- or 12-mer (depending on the number of modifications in the oligodeoxynucleotide).

In conclusion, synthesis of novel diamine linked thymidine dimers have been accomplished. Two of these have been incorporated into different oligodeoxynucleotides. Oligomers containing the amide 9 are promising candidates as antisense oligonucleotides in that the duplex stability only slightly is affected and the stability towards nucleolytic degradation is increased considerably. On the contrary oligomers containing amine 7 seems to be of no use in antisense oligonucleotides due to the pronounced destabilization observed for duplexes formed with complementary DNA strands.

REFERENCES

- 1. Varma, R. S. Synlett. 1993, 621.
- 2. Fox, J. J.; Miller, N. C. J. Org. Chem. 1963, 28, 936.
- Weis, A. L.; Hausheer, F. H.; Chaturvedula, P. V. C.; Delecki, D. J.; Cavanaugh, P. F., Jr.; Moskwa, P. S.; Oakes, F. T. Oligonucleotide analogs for inhibiting gene expression, International Patent WO 92/02534 1992, CA 117: 105700.

- 4.. Moss, G. P.; Reese, C. B.; Schofield, K.; Shapiro, R.; Todd, Lord J. Chem. Soc. **1963**, 1149.
- 5. The regular protocol of the DNA-synthesizer was used except that the coupling time for the modifications was increased from 2 to 12 minutes.
- Svendsen, M. L.; Wengel, J.; Dahl, O.; Kirpekar, F.; Roepstorff, P. Tetrahedron 1993, 49, 11341.
- 7.. Tidd, D. M.; Warenius, H. M. Br. J. Cancer, 1989, 60, 343.