


compounds.^[2,3,10–12] In fact, this mechanism was notably successful in explaining the redistribution reactions of various organosilicon, -germanium, and -phosphorus systems.^[13–15] We have previously given direct experimental evidence for 1,2-silyl-migration from Si to the metal M (**A**→**B** or **D**→**B'**) via isolating complexes of the type **B'** or **C**, which are formed in reactions of complexes of type **A** or **D**.^[2,3,16] Furthermore, we have observed fluxional behavior in the 1,3-migration of methyl groups on an externally donor-stabilized silyl(silylene)iron complex [Cp(CO)Fe(=SiMe₂←HMPA)SiMe₃] (HMPA = hexamethyl phosphoramide) by variable-temperature NMR spectroscopy.^[17] In this process, we assumed that the coordinated HMPA dissociates at elevated temperatures to generate a donor-free silyl(silylene) complex. We now give direct evidence for 1,3-alkyl migration (**B**→**B'** and vice versa) and 1,2-silyl migration from M to Si (**B**→**A** or **B'**→**D**) by employing newly synthesized, donor-free silyl(silylene)iron complexes.

Photolysis of [Cp'Fe(CO)₂Me] (**1a**: Cp' = η⁵-C₅Me₅ (Cp*); **1b**: Cp' = η⁵-C₅H₅ (Cp)) in the presence of HSiMe₂-SiMes₂Me (**2**; Mes = mesityl (2,4,6-trimethylphenyl)) produced the first donor-free silyl(silylene)iron complexes [Cp'Fe(CO)(=SiMes₂)SiMe₃] (**3a**: Cp' = Cp*, 60%; **3b**: Cp' = Cp, 38% yield, calculated by NMR spectroscopy [Eq. (1)]). Complex **3a** could be isolated as orange crystals in 40% yield, whereas isolation of **3b** was unsuccessful



because of its extreme instability. We have previously synthesized the tungsten analogue of **3a** by a similar method, but the chemistry has not been thoroughly investigated.^[16]

The molecular structure of **3a** is shown in Figure 1.^[18] The two mesityl groups are on the silylene ligand, while all of the three methyl groups are on the silyl ligand. The iron–silylene bond (Fe–Si(1) 2.154(1) Å) is about 9% shorter than the iron–silyl bond (Fe–Si(2) 2.343(2) Å) and is the shortest reported bond of this type.^[19] The silylene silicon atom is tricoordinate and its geometry is almost planar (sum of the three bond angles around Si(1) = 359.3 (2)°). No intermolecular bonding interaction was found. The ²⁹Si NMR spectra of **3a** and **3b** show signals for the dimesitylsilylene ligand at extremely low field (365.8 ppm for **3a** and 372.0 ppm for **3b**), which is characteristic of the donor-free dialkyl- or diaryl-silylene complexes.^[5,16] Also present are the resonances for the trimethylsilyl ligand (28.4 ppm for **3a** and 31.0 ppm for **3b**). These data unambiguously demonstrate the donor-free silyl(silylene)iron structures. In each of the ¹H NMR spectra of **3a** and **3b**, all the four *o*-Me groups, four *m*-H atoms, and two *p*-Me groups in two mesityl groups are inequivalent at room temperature. Apparently, the extremely congested





Figure 1. ORTEP drawing of **3a** showing thermal ellipsoids at the 50% probability level. Selected bond lengths [Å] and angles [°]: Fe(1)–Si(1) 2.154(1), Fe(1)–Si(2) 2.343(1), Fe(1)–C(11) 1.724(4); Si(1)–Fe(1)–Si(2) 93.15, Fe(1)–Si(1)–C(12) 127.8(1), Fe(1)–Si(1)–C(21) 127.2(1), C(12)–Si(1)–C(21) 104.3(2).

structures of **3a** and **3b** lead to hindered rotation around both the Fe=Si and the Si–C(mesityl) bonds.

Sharma and Pannell previously reported that the photolysis of linear oligosilanyl–[Fe(CO)₂Cp] complexes containing more than three silicon atoms produces highly branched, tris(silyl)silyl iron complexes in high yields, for example, [(Me₃Si(Me₂Si)₃]Fe(CO)₂Cp] is converted to [(Me₃Si)₃Si–Fe(CO)₂Cp] on irradiation.^[12] In this reaction, the 1,2-silyl migration from the Fe center to the silylene silicon atom on the silyl(silylene) iron intermediates (corresponding to **B**→**A** or **B'**→**D**; Scheme 1) could play an important role.

To confirm this hypothesis, thermolysis of **3a** in the presence of several two-electron-donor ligands was carried out. As a result, when **3a** was heated to 80 °C for 6 h in the presence of *t*BuNC, a disilanyl complex [Cp*Fe(CO)(CN-



*t*Bu)SiMesMeSiMesMe₂] (**4**) was isolated as a main product in 25% yield [Eq. (2)]. The ²⁹Si NMR signals of **4** appear in the normal range of disilanyl iron complexes (9.5 ppm for Fe–Si and –11.2 ppm for terminal Si atoms). The molecular structure of **4** is shown in Figure 2.^[18] A *t*BuNC molecule is terminally coordinated to the iron center, and each of the α- and β-Si atoms of the disilanyl ligand is coordinated to a mesityl group. The Fe–Si(1) and Si(1)–Si(2) bond lengths are 2.4107(7) and 2.4004(9) Å, respectively, which are normal values for single bonds.

A mechanism that can rationalize the reactions in Equations (1) and (2) is illustrated in Scheme 2. From **1a**, successive CO dissociation, oxidative addition of **1**, methane reductive elimination, 1,2-silyl migration, and 1,3-methyl migration occur to afford **3a**. Three isomeric donor-free silyl(silylene) complexes (**3a**, **3a'**, and **3a''**) are in rapid





Figure 2. ORTEP drawing of **4** showing thermal ellipsoids at the 50% probability level. Selected bond lengths [Å] and angles [°]: Fe(1)–Si(1) 2.4107(7), Fe(1)–C(11) 1.732(2), Fe(1)–C(33) 1.808(2), Si(1)–Si(2) 2.4004(9), N(1)–C(33) 1.174(3); Fe(1)–Si(1)–Si(2) 118.74(3), Si(1)–Fe(1)–C(11) 84.23(9), Si(1)–Fe(1)–C(33) 94.92(7), C(11)–Fe(1)–C(33) 92.5(1), Fe(1)–C(33)–N(1) 173.4(2), C(33)–N(1)–C(34) 162.0(3).



Scheme 2. A mechanism for the formation of $[\text{Cp}^*\text{Fe}(\text{CO})(=\text{SiMe}_2)\text{SiMe}_3]$ (**3a**) and $[\text{Cp}^*\text{Fe}(\text{CO})(\text{CN}t\text{Bu})\text{SiMe}_2\text{SiMe}_2\text{SiMe}_2]$ (**4**).

equilibrium at room temperature, where **3a** is the major and only observable isomer. When this equilibrium mixture is heated in the presence of *t*BuNC, 1,2-migration of the silyl ligand onto the silylene ligand followed by coordination of *t*BuNC to the unsaturated iron center occurs to produce **4**. It should be noted that both **3a** and **4** take the structures that obviously minimize the steric repulsion between the bulky groups, namely, the two mesityl groups and a pentamethylcyclopentadienyl group. In other words, **3a** and **4** are the thermodynamically controlled products. Both the formation

of **3a** and its conversion to **4** involves 1,2-silyl migration and 1,3-alkyl and/or aryl migration processes. These are considered to be concerted processes with low energy barriers.^[20] Importantly, through the latter process, usually robust Si–C bonds readily cleave under extremely mild conditions: The typical bond dissociation energy of the Si–C single bond is 301 kJ mol^{-1} , which is comparable to that of the C–C single bond (346 kJ mol^{-1}).^[21]

In this paper, we have provided the most straightforward evidence for extremely facile 1,2- and 1,3-group migrations in silyl(silylene) complex systems. These observations clearly show how organosilicon species bound to a transition-metal center can change their structures in an amazingly dynamic fashion through extremely facile Si–C and Si–Si bond fission and formation processes. A more detailed elucidation of the dynamic behavior is underway.

Experimental Section

3a: A pentane solution (3 mL) of $[\text{Cp}^*\text{Fe}(\text{CO})_2\text{Me}]$ (**1a**; 1.02 g, 3.89 mmol) and $\text{HSiMe}_2\text{SiMe}_2\text{Me}$ (**2**; 1.01 g, 2.96 mmol) in a pyrex sample tube with a teflon vacuum valve was irradiated for 80 min with a 450 W medium-pressure Hg lamp immersed in a water bath (4°C). The reaction mixture was degassed every 20 min by a conventional freeze-pump-thaw cycle on a vacuum line. The reaction mixture was filtered through a glass filter and volatiles were removed from the filtrate under reduced pressure. The residue was recrystallized from pentane at -30°C to afford orange crystals of $[\text{Cp}^*\text{Fe}(\text{CO})(=\text{SiMe}_2)\text{SiMe}_3]$ (**3a**) in 40% yield (0.660 g, 1.18 mmol). ^1H NMR (300 MHz, $[\text{D}_6]$ benzene): $\delta = 0.59$ (s, 9H, SiMe_3), 1.56 (s, 15H, C_5Me_5), 2.05 (s, 3H, *o*-Me), 2.10 (s, 3H, *o*-Me), 2.12 (s, 3H, *o*-Me), 2.15 (s, 3H, *o*-Me), 2.74 (s, 3H, *p*-Me), 3.05 (s, 3H, *p*-Me), 6.51 (s, 1H, *m*-H), 6.56 (s, 1H, *m*-H), 6.79 (s, 1H, *m*-H), 6.86 ppm (s, 1H, *m*-H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75.5 MHz, $[\text{D}_6]$ benzene): $\delta = 9.5$ (SiMe_3), 10.1 (C_5Me_5), 21.1 (*p*-Me), 23.7 (*o*-Me), 24.0 (*o*-Me), 24.7 (*o*-Me), 24.9 (*o*-Me), 93.7 (C_5Me_5), 128.7 ($\text{C}_6\text{H}_2\text{Me}_3$), 129.2 ($\text{C}_6\text{H}_2\text{Me}_3$), 138.7 ($\text{C}_6\text{H}_2\text{Me}_3$), 138.9 ($\text{C}_6\text{H}_2\text{Me}_3$), 139.2 ($\text{C}_6\text{H}_2\text{Me}_3$), 139.3 ($\text{C}_6\text{H}_2\text{Me}_3$), 142.5 ($\text{C}_6\text{H}_2\text{Me}_3$), 142.8 ($\text{C}_6\text{H}_2\text{Me}_3$), 145.3 ($\text{C}_6\text{H}_2\text{Me}_3$), 145.6 ($\text{C}_6\text{H}_2\text{Me}_3$), 220.2 ppm (CO); $^{29}\text{Si}\{^1\text{H}\}$ NMR (59.6 MHz, $[\text{D}_6]$ benzene): $\delta = 28.4$ (SiMe_3), 365.8 ppm (SiMe_2); IR ($[\text{D}_6]$ benzene solution): $\tilde{\nu} = 1905 \text{ cm}^{-1}$ (s, ν_{CO}); MS (EI, 70 eV) 558 (M^+ , 8), 543 ($M^+ - \text{CH}_3$, 30), 515 ($M^+ - \text{CH}_3 - \text{CO}$, 12), 73 (SiMe_3 , 100); elemental analysis calcd (%) for $\text{C}_{32}\text{H}_{46}\text{FeOSi}_2$: C 68.79, H 8.30; found: C 69.07, H 8.41.

4: A toluene solution (5 mL) of **3a** (0.103 g, 0.184 mmol) and *t*BuNC (0.0730 g, 0.878 mmol) in a pyrex tube with a teflon vacuum valve was heated to 80°C for 6 h. After removal of volatiles, the yellow residue was recrystallized from toluene/hexane to afford yellow crystals of $[\text{Cp}^*\text{Fe}(\text{CO})(\text{CN}t\text{Bu})\text{SiMe}_2\text{SiMe}_2\text{SiMe}_2]$ (**4**) in 25% yield (0.030 g, 0.047 mmol). ^1H NMR (300 MHz, $[\text{D}_6]$ benzene): $\delta = 0.69$ (s, 3H, $\text{SiMe}_2\text{SiMe}_2$), 0.97 (s, 3H, $\text{SiMe}_2\text{SiMe}_2$), 1.09 (s, 1H, $\text{SiMe}_2\text{SiMe}_2$), 1.46 (s, 15H, C_5Me_5), 2.15 (s, 3H, *p*-Me), 2.23 (s, 3H, *p*-Me), 2.45 (s, 6H, *o*-Me), 2.52 (s, 3H, *o*-Me), 2.64 (s, 3H, *o*-Me), 6.78 (s, 2H, *m*-H), 6.84 (s, 1H, *m*-H), 6.88 ppm (s, 1H, *m*-H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75.5 MHz, $[\text{D}_6]$ benzene): $\delta = 6.4$ (SiMe_2), 6.7 (SiMe_2), 9.9 (C_5Me_5), 11.7 (SiMe_2), 21.1 ($\text{C}_6\text{H}_2\text{Me}_3$), 25.8 ($\text{C}_6\text{H}_2\text{Me}_3$), 26.8 ($\text{C}_6\text{H}_2\text{Me}_3$), 28.0 ($\text{C}_6\text{H}_2\text{Me}_3$), 31.2 (CMe_3), 56.3 (CMe_3), 92.7 (C_5Me_5), 128.9 ($\text{C}_6\text{H}_2\text{Me}_3$), 129.1 ($\text{C}_6\text{H}_2\text{Me}_3$), 129.3 ($\text{C}_6\text{H}_2\text{Me}_3$), 130.1 ($\text{C}_6\text{H}_2\text{Me}_3$), 136.3 ($\text{C}_6\text{H}_2\text{Me}_3$), 136.9 ($\text{C}_6\text{H}_2\text{Me}_3$), 137.2 ($\text{C}_6\text{H}_2\text{Me}_3$), 140.3 ($\text{C}_6\text{H}_2\text{Me}_3$), 144.5 ($\text{C}_6\text{H}_2\text{Me}_3$), 145.3 ($\text{C}_6\text{H}_2\text{Me}_3$), 176.6 (FeCN), 222.3 ppm (CO); $^{29}\text{Si}\{^1\text{H}\}$ NMR (59.6 MHz, $[\text{D}_6]$ benzene): $\delta = -11.2$ ($\text{SiMe}_2\text{SiMe}_2$), 9.5 ppm ($\text{SiMe}_2\text{SiMe}_2$); IR ($[\text{D}_6]$ benzene solution): $\tilde{\nu} = 1907 \text{ cm}^{-1}$ (s) (ν_{CO}); MS (EI, 70 eV) 641 (M^+ , 0.3), 626 ($M^+ - \text{Me}$, 0.6), 556 ($M^+ - \text{CO} - t\text{Bu}$, 4), 515 ($M^+ - \text{CO} - \text{CN}t\text{Bu} - \text{Me}$, 7), 464

($M^+ - CO - 2Me - Mes$, 100); elemental analysis calcd (%) for $C_{37}H_{55}FeONSi_2$: C 69.24, H 8.64, N 2.18; found: C 69.31, H 8.66, N 2.28.

Received: July 30, 2003 [Z52519]

Keywords: group migration · iron · ligands · rearrangement · transition metals

- [1] T. M. Trnka, R. H. Grubbs, *Acc. Chem. Res.* **2001**, *34*, 18.
- [2] H. Ogino, *Chem. Rec.* **2002**, *2*, 291.
- [3] M. Okazaki, H. Tobita, H. Ogino, *Dalton Trans.* **2003**, 493.
- [4] S. R. Klei, T. D. Tilley, R. G. Bergman, *Organometallics* **2002**, *21*, 3376, 4648.
- [5] J. D. Feldman, J. C. Peters, T. D. Tilley, *Organometallics* **2002**, *21*, 4065.
- [6] H. Sakaba, M. Tsukamoto, T. Hirata, C. Kabuto, H. Horino, *J. Am. Chem. Soc.* **2000**, *122*, 11 511.
- [7] T. A. Schmedake, M. Haaf, B. J. Paradise, D. Powell, R. West, *Organometallics* **2000**, *19*, 3263.
- [8] S. H. A. Petri, D. Eikenberg, B. Neumann, H.-G. Stammler, P. Jutzi, *Organometallics* **1999**, *18*, 2615.
- [9] B. Gehrhuis, P. B. Hitchcock, M. F. Lappert, H. Maciejewski, *Organometallics* **1998**, *17*, 5599.
- [10] K. H. Pannell, J. Cervantes, C. Hernandez, J. Cassias, S. P. Vincenti, *Organometallics* **1986**, *5*, 1056.
- [11] H. Tobita, K. Ueno, H. Ogino, *Chem. Lett.* **1986**, 1777.
- [12] H. K. Sharma, K. H. Pannell, *Chem. Rev.* **1995**, *95*, 1351.
- [13] K. Tamao, G.-R. Sun, A. Kawachi, *J. Am. Chem. Soc.* **1995**, *117*, 8043.
- [14] S. M. Katz, J. A. Reichl, D. H. Berry, *J. Am. Chem. Soc.* **1998**, *120*, 9844.
- [15] P. Braunstein, M. Knorr, C. Stern, *Coord. Chem. Rev.* **1998**, *178–180*, 903.
- [16] K. Ueno, S. Asami, N. Watanabe, H. Ogino, *Organometallics* **2002**, *21*, 1326.
- [17] K. Ueno, K. Nakano, H. Ogino, *Chem. Lett.* **1996**, 459.
- [18] **3a**: monoclinic; $P2_1/c$; $a = 10.4593(2)$, $b = 18.3566(6)$, $c = 16.0338(4)$ Å, $\beta = 96.5724(7)^\circ$; $V = 3058.2(1)$ Å³; $Z = 4$; $C_{32}H_{46}FeOSi_2$; $T = 150$ K, 26 965 reflections, 6692 independent ($R_{int} = 0.043$); $R1 = 0.046$ ($I > 3\sigma(I)$), $Rw = 0.106$; $\mu = 5.93$ cm⁻¹; Full-matrix least-squares on F^2 . **4**: monoclinic; $P2_1$; $a = 8.8296(2)$, $b = 20.6795(5)$, $c = 9.9852(2)$ Å, $\beta = 99.970(1)^\circ$; $V = 1795.68(7)$ Å³; $Z = 2$; $C_{37}H_{55}FeNOSi_2$; $T = 150$ K, 17 660 reflections, 4233 independent ($R_{int} = 0.036$); $R1 = 0.027$ ($I > 2\sigma(I)$), $Rw = 0.063$; $\mu = 5.14$ cm⁻¹; Full-matrix least-squares on F^2 . CCDC-215618 (**3a**) and CCDC-215619 (**4**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/conts/retrieving.html (or from the Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge CB21EZ, UK; fax: (+44)1223-336-033; or deposit@ccdc.cam.ac.uk).
- [19] Based on a search of the Cambridge Structural Database, CSD version 5.24 (November **2002**).
- [20] K. Morokuma, private communication; Z. Liu, PhD Thesis, Emory University, **2000**, chap. 3. MO calculation (B3LYPLANL2DZ) for the 1,3-migration of a Me group from a silyl ligand to a silylene ligand in $[CpFe(CO)(=SiMe_2)SiMe_3]$ showed that this is a concerted process with no intermediate and the activation energy of this process is 12.0 kcal mol⁻¹.
- [21] J. Emsley, *The Elements*, 3rd ed., Oxford University Press, Oxford, **1998**.