

Novel Steroidal Inhibitors of Human Cytochrome P450_{17 α} (17 α -Hydroxylase-C_{17,20}-lyase): Potential Agents for the Treatment of Prostatic Cancer

Gerard A. Potter,[†] S. Elaine Barrie, Michael Jarman,* and Martin G. Rowlands

Cancer Research Campaign Centre for Cancer Therapeutics at the Institute of Cancer Research,
Cancer Research Campaign Laboratory, 15 Cotswold Road, Sutton, Surrey SM2 5NG, U.K.

Received January 19, 1995[§]

Steroidal compounds having a 17-(3-pyridyl) substituent together with a 16,17-double bond have been synthesized, using a palladium-catalyzed cross-coupling reaction of a 17-enol triflate with diethyl(3-pyridyl)borane, which are potent inhibitors of human testicular 17 α -hydroxylase-C_{17,20}-lyase. The requirement for these structural features is stringent: compounds having 2-pyridyl (**9**), 4-pyridyl (**10**), or 2-pyridylmethyl (**11**) substituents instead of the 3-pyridyl substituent were either poor inhibitors or noninhibitory. Reduction of the 16,17-double bond to give 17 β -pyridyl derivatives diminished potency with 3-pyridyl substitution (**3** \rightarrow **27**; IC₅₀ for lyase, 2.9 \rightarrow 23 nM) but increased it with a 4-pyridyl substituent present (**10** \rightarrow **28**; IC₅₀ 1 μ M \rightarrow 53 nM). In contrast, a variety of substitution patterns in rings A–C of the steroid skeleton afforded inhibitors having potencies similar to those most closely related structurally to the natural substrates pregnenolone and progesterone, respectively 17-(3-pyridyl)androsta-5,16-dien-3 β -ol (**3**, K_{i,app} < 1 nM; IC₅₀ for lyase, 2.9 nM) and 17-(3-pyridyl)androsta-4,16-dien-3-one (**15**; IC₅₀ for lyase, 2.1 nM). Thus compounds having variously aromatic ring A (**18**), saturated rings A/B (**21**, **22**), and oxygenated ring C (**26**) exhibited IC₅₀ values for lyase (1.8–3.0 nM) falling within a 2-fold range. The most potent compounds are candidates for development as drugs for the treatment of hormone-dependent prostatic carcinoma.

Carcinoma of the prostate is now the most prevalent cancer in men in the USA. In 1993, 165 000 new cases were expected to be diagnosed, of which 35 000 will die of metastatic prostatic cancer.¹ The most widely accepted drug treatment is the use of GnRH agonists, which act by interfering with the production of testosterone by the testes and represent a medical alternative to orchiectomy.² However neither GnRH agonists nor orchiectomy deplete the synthesis of androgens through the adrenal route, and levels of testosterone and dihydrotestosterone in the prostate are respectively still 25% and 10% of pretreatment levels even after 3 months treatment with a GnRH agonist.³ The importance of androgen synthesis by the adrenal route in maintaining tumor growth is suggested by the improved therapeutic benefit, both in terms of increase in progression-free survival time and survival advantage, seen in patients treated with the combination of GnRH agonist or orchiectomy with an antiandrogen, compared with those given GnRH agonist or orchiectomy alone.^{4,5} It is proposed that the role of the antiandrogen is to counteract the stimulant action of residual androgens, synthesized through the adrenal route, on androgen receptors in the prostate cancer cells.

In principle, the effects of the combined therapy could be realized by a single drug which inhibits the enzyme steroidal 17 α -hydroxylase-C_{17,20}-lyase. This enzyme is responsible for androgenic hormone biosynthesis which produces dehydroepiandrosterone and androstenedione, immediate precursors of testosterone, from their respective precursors pregnenolone and progesterone, in both testes and adrenals. The imidazole antifungal agent ketoconazole inhibits this enzyme when given in high

doses to male patients and produces the symptoms of androgen suppression. This drug has been used to treat prostate cancer,⁶ and although success has been reported in some studies,^{7,8} it proved less promising in others.^{9,10} The undesirable side effects, coupled with the inconvenience of the three times daily schedule which is dictated by its short half-life, limit its potential clinical usefulness. Nevertheless the clinical results obtained, coupled with a very recent report that careful scheduling of ketoconazole can produce prolonged responses in previously hormone-refractory prostate cancer,¹¹ lend credence to the selection of this enzyme target and impetus to the design and development of a more enzyme-selective, less toxic, and less metabolically labile inhibitor.

We report here on the synthesis and inhibitory activity toward the individual 17 α -hydroxylase and C_{17,20}-lyase components of the target enzyme, obtained from human testis, of a variety of steroidal compounds having as their common structural feature a 17-(3-pyridyl) substituent together with a 16,17-double bond in the steroidal skeleton. We have previously explored nonsteroidal inhibitors containing a pyridyl residue, starting from the serendipitous discovery that certain esters of 4-pyridylacetic acid were effective inhibitors of the hydroxylase-lyase enzyme from rat testis,¹² findings which have in part been rationalized by crystallographic and molecular modeling studies.¹³ More recently, esters of 3-pyridylacetic acid have been evaluated, using enzyme from human testis.¹⁴

The design concept used here was to consider how a pyridyl substituent could be incorporated into the actual steroid skeleton such that the pyridyl nitrogen lone pair would coordinate to the iron atom of the heme cofactor in the active site of the enzyme. The initial step of the *de novo* mechanism-based design approach was to postulate a complete catalytic cycle for the enzyme

* To whom enquiries should be addressed.

[†] Present address: Chiroscience Ltd., Milton Rd., Cambridge CB4 4WE, U.K.

[§] Abstract published in *Advance ACS Abstracts*, May 15, 1995.





Figure 1. Postulated complete catalytic cycle for the 17 α -hydroxylase-C_{17,20}-lyase enzyme. For clarity, only the steroid D-ring and the cofactor iron atom are shown.

(Figure 1) and then to consider the juxtaposition between the steroid D-ring and the heme cofactor from the putative transition state geometry. For this purpose, three-dimensional molecular models were constructed of the putative transition states using the Cochrane orbit molecular modeling system. From this analysis, it was postulated that a steroid incorporating a 16,17-double bond with the 17-position substituted by a 2-pyridyl group may inhibit the hydroxylase step and a 3-pyridyl derivative may inhibit the lyase step, while a 4-pyridyl analog should not inhibit either step. However, the enzyme may not tolerate an aromatic ring attached to the 17-position, and all three compounds may be inactive, even if the coordination geometry is correct.

The steroidal skeleton chosen for the first compound which was synthesized on the basis of this concept, namely the novel steroid **3**, was that of pregnenolone, which appears to be the preferred substrate for the hydroxylase activity of the human enzyme in the testis.¹⁵ Alternative orientations of the pyridyl ring relative to the steroidal framework were explored by synthesizing the 2- (**9**) and 4- (**10**) pyridyl analogs, as was the effect of a spacer group between a 2-pyridyl residue and a C-17 (compound **11**). The second 17-(3-pyridyl) derivative synthesized was **15**, analogously related to progesterone, the alternative substrate for the hydroxylase activity of the target enzyme. Further molecules synthesized retained the ring D substitution pattern of **3** and **15** while further exemplifying the effect on enzyme inhibition of structural variations in rings

Scheme 1^a




A, B, and C. Finally, the effect of reducing the 16,17-double bond in **3** and **10** was explored.

Results

Chemistry. A general method for introducing the required 17-pyridyl 16,17-ene functionality into ring D was by palladium-catalyzed cross-coupling of steroidal 17-enol triflates with suitable pyridyl-containing nucleophilic coupling partners. For the synthesis of **3** (Scheme 1), dehydroepiandrosterone 3-acetate was converted into its 17-enol triflate **1** by base-catalyzed reaction with triflic anhydride in the presence of the hindered base 2,6-di-*tert*-butyl-4-methylpyridine. This reaction also produced the 3,5-diene **4** in 10% yield. The 3-pyridyl group was then introduced into the 17-position by reacting **1** with diethyl(3-pyridyl)borane in THF,


Chart 1



using bis(triphenylphosphine)palladium(II) chloride as catalyst (0.01 equiv) and aqueous Na₂CO₃ as nucleophilic activator. The reaction proceeded remarkably efficiently, without the potential side reactions of triflate hydrolysis or ethyl coupling, to give the acetate **2** in 84% isolated yield. From **4**, the 3-pyridyl derivative **5** was similarly obtained. The acetyl group of **2**, which was stable to the mildly basic conditions of the coupling reaction, was easily removed with aqueous methanolic NaOH to afford the target 3-pyridyl steroid **3**.


Although these coupling reactions were developed independently, the palladium-catalyzed cross-coupling of organoboron reagents with an enol triflate has been reported recently by Suzuki and co-workers.¹⁶ Their reactions employed arylboronic acids and 9-alkyl-9-BBN reagents and the mild base K₃PO₄ as the nucleophilic activator under strictly anhydrous conditions. Our use of diethyl(3-pyridyl)borane was prompted by its commercial availability (it is also easily synthesized¹⁷) and its previous use in palladium-catalyzed cross-coupling reactions with aryl iodides.¹⁸ Some features of our reaction compared with that of Suzuki are noteworthy. We found that the catalyst Pd(PPh₃)₂Cl₂ was superior to Pd(PPh₃)₄ and consistently gave better yields of coupled product. The catalyst could also be used at much lower levels, and even at 0.001 equiv, good yields were obtained with prolonged reaction times. Importantly our reaction did not require anhydrous conditions, and indeed an aqueous THF solvent system was employed. Our method of introducing the 17-pyridyl 16,17-ene functionality was more efficient and higher yielding than a previous route,^{19,20} reaction of 3-pyridyllithium with a 17-keto steroid and dehydration of the resulting tertiary alcohol.

The 2-pyridyl (**6**), 4-pyridyl (**7**), and 2-picolyl (**8**) steroidal acetates (Chart 1) were synthesized similarly to **2** but employing different nucleophilic coupling partners and modifying the conditions accordingly. The reagents used to prepare **6** and **8** were 2-pyridyl- and 2-picolylzinc chloride, respectively. In the latter case the intermediate **8** was converted without isolation directly into **11** in good overall yield (79%). An attempt to prepare the 3-picolyl analog of **11** using 3-picolylzinc chloride was unsuccessful due to homocoupling of this reagent. In the synthesis of the 2- (**6**) and 4- (**7**) pyridyl steroid acetates, the novel palladium catalyst bromo-(isopropenyl)bis(triphenylphosphine)palladium(II) was employed. Its use enabled the coupling reaction to be carried out at ambient temperature, thereby avoiding side reactions, and **6** was obtained in 74% yield from which hydrolysis gave the required 2-pyridyl analog **9**. The catalyst had been developed to enable low-temperature cross-coupling reactions for the stereoselective synthesis of (*E*)-4-hydroxytamoxifen^{21,22} and was prepared from 2-bromopropene and tetrakis(triphenylphos-

Scheme 2^a

^a (a) C₇F₈, CsF, DMF; (b) Tf₂O, base; (c) 3-PyBEt₂, Pd(PPh₃)₂Cl₂, THF, H₂O, Na₂CO₃; (d) HCl, H₂O, EtOH; (e) Al(O-*i*-Pr)₃.


Chart 2




phine)palladium(0) by a procedure analogous to that used to make benzylchlorobis(triphenylphosphine)palladium(II).²³ When the coupling reaction was performed using 4-pyridylzinc chloride, prepared from 4-bromopyridine, only a low yield (18%) of the 4-pyridyl steroid acetate **7** was obtained. Instability of 4-halopyridines can restrict the use of 4-pyridylmagnesium and -zinc halides in palladium cross-coupling reactions, and diethyl(4-pyridyl)borane has been used as an alternative reagent.²⁴ Here, lithium trimethoxy(4-pyridyl)boronate, an intermediate in the synthesis²⁵ of 4-pyridylboronic acid, was the organoboron reagent used, and the coupled product thus obtained, **7**, was hydrolyzed directly to give the 4-pyridyl steroid **10** in 53% yield overall from **1**.

The preparation of **15**, starting from androstenedione (Scheme 2), required selective protection of the 3-keto function, to prevent the formation of a 3-dienol triflate.²⁶ Protection as the perfluorotolyl enol ether **12** by reaction with octafluorotoluene in the presence of cesium fluoride has proved to be a convenient one-step procedure.²⁷ The perfluoroaryl group was stable to the subsequent steps needed to insert the pyridyl substituent and was then cleaved by acidic hydrolysis. It was later found that **15** was more conveniently prepared directly from **3** by Oppenauer oxidation using cyclohexanone and aluminum isopropoxide.

Several 3-pyridyl derivatives (**18**, **21**, **22**; Chart 2) exemplifying further structural variation in rings A and B were prepared using procedures analogous to those already described. Adrenosterone was the starting point for the synthesis of a ring-C-substituted variant, **26** (Scheme 3), which was prepared in good overall yield (60%). The formation of the *tert*-butyldimethylsilyl dienol ether **23** provided an alternative protecting

Scheme 3^a

^a (a) *t*-BDMSOTf, base; (b) (*i*-Pr)₂NLi, PhNTf₂; (c) 3-PyBEt₂, Pd(PPh₃)₂Cl₂, THF, H₂O, Na₂CO₃; (d) Bu₄NF, THF, H₂O.

Scheme 4^a

^a (a) N₂H₄, AcOH, EtOH, air; (b) Red-Al, ZnCl₂, THF.

strategy for the 3-keto function. The chemical shifts for the two vinylic protons in this product were very similar to those previously reported for the silyl dienol ether formed from a testosterone derivative,²⁸ and the present product is therefore similarly formulated as the 2,4-dienol ether. In the following step, *N*-phenyltriflimide²⁹ was employed to prepare the enol triflate **24** since use of triflic anhydride resulted in desilylation and 3-dienol triflate formation. This also enabled selective formation of the 17-enol triflate without affecting the 11-keto function by preparing the intermediate lithium enolate under kinetic conditions at low temperature.

Lastly, analogs containing a saturated D-ring were prepared from the corresponding 16,17-ene compounds. Reduction of **3** using diimide, generated *in situ* from hydrazine hydrate, gave the 17β-(3-pyridyl) steroid **27** (Scheme 4). Reduction of the 16,17-double bond of the 4-pyridyl steroid (**10**) utilized the electron-withdrawing influence of the 4-pyridyl substituent under electrophilic activation by zinc chloride to achieve direct hydride reduction with Red-Al to produce the 17β-(4-pyridyl) steroid **28**. The β-orientation of the pyridyl ring in compounds **27** and **28** was confirmed by ¹H-NMR spectroscopy which showed an apparent triplet with a coupling constant of 10 Hz for the 17α-proton which is characteristic of 17β-substituted steroids.^{20,30} Attempts at preparing the corresponding 17α-(4-pyridyl) analog, by either direct reduction of **10** or epimerization of **28**, were unsuccessful.

Inhibition of Human Testicular 17α-Hydroxylase and C_{17,20}-Lyase. Structure-Activity Relationships. We have identified as potent inhibitors of human testicular steroidal 17α-hydroxylase-C_{17,20}-lyase a variety of pyridyl steroids having as their common structural feature the 17-(3-pyridyl) 16,17-ene moiety (Table 1). Although it might be expected that the most potent compounds would be those (**3**, **15**) with structures most closely related to natural substrates, there was an unexpected tolerance for structural variation in this respect. Comparing **3** and **15** with analogs (**18**, **21**, **22**,

Table 1. Enzyme Inhibition Data

compound	IC ₅₀ (nM) ^a		IC ₅₀ (μM) ^a	
	C _{17,20} -lyase	17α-hydroxylase	aromatase	5α-reductase
2	17	18		
3	2.9	4	>20	>50
5	5.6	12.5		
9	76	270	>20	
10	1000	4000	>20	
11	>10 000	>10 000		
14	>10 000	>10 000		
15	2.1	2.8	1.8	10
18^b	1.8	2.6		
21	2.5	4.3		
22	3	4.7		>50
26	2.9	13		
27	23	47		
28	53	160		
ketoconazole	26	65		

^a The standard errors were usually <10% of the IC₅₀ value. The concentration of enzyme in the assays for lyase/hydroxylase inhibition was estimated to be about 4–5 nM, except in the assays of **9**, **11**, **14**, and ketoconazole for which the concentration was ca. 25 and 10 nM for the lyase and hydroxylase assays, respectively. ^b Other biological activity: estrogen receptor binding affinity (estradiol = 100), 4.9.

26) synthesized from other naturally occurring steroid precursors, there was little variation (from 1.8 to 3.0 nM) in the IC₅₀ values for inhibition of the lyase component. The absence of any functionality at the 3-position in the steroid skeleton leads to a modest drop in potency (compound **5**). The markedly lower potency of the acetoxy derivative **2** compared with **3** could reflect a limited bulk tolerance at the 3-position, as indicated by the total loss in activity for the much more sterically demanding perfluorotolyl derivative **14** of the potent inhibitor **15**. The stringent requirement for the 17-(3-pyridyl) 16,17-ene functionality for good inhibition was in marked contrast to the relative flexibility in relation to other features discussed and is reflected in the marked reduction, or abolition of activity, on relocating the pyridyl nitrogen (compounds **9**, **10**) or on reducing the 16,17-double bond of **3** to give the 17β-pyridyl derivative **27**. In contrast, reduction of the 4-pyridyl derivative **10** gave a product, **28**, with markedly improved inhibitory potency over its parent.

The most inhibitory compounds in the present study were far more potent than any inhibitor of hydroxylase/lyase for which comparable data have been previously described. The K_{i,app} for **3** was <1 nM, whereas the most potent inhibitor, also steroidal, reported to date is 17β-(cyclopropylamino)androst-5-en-3β-ol³¹ with a K_{i,app} of 90 nM. Another steroidal compound, 4-pregnen-3-one-20β-carboxaldehyde oxime has been developed as a combined inhibitor of this enzyme and testosterone 5α-reductase.³² Though a potent inhibitor (K_i = 16 nM) of the reductase, it was much less inhibitory toward the rat hydroxylase/lyase, being comparable to ketoconazole. 17β-Ureido-substituted steroids with potent activity toward the rat hydroxylase/lyase enzyme have been described.^{33,34} Though the data are presented in a way not easily comparable with the results of the present study, one of these compounds, 17β-ureido-1,4-androstadien-3-one, markedly suppressed testosterone levels and ablated androgen-dependent organs in the rat. Liarazole is a nonsteroidal imidazole derivative having activity toward the rat testicular enzyme very similar³⁵ to that of ketoconazole. No example among our previously mentioned^{12,14} esters of 4- and 3-pyridylacetic acid compares in potency with the best of the present steroidal derivatives.

Other Biological Activities. While inhibition of other targets was not explored in detail in the present study, limited evaluations have been carried out (Table 1), particularly where such activity might be anticipated, from structural analogy with compounds known to interact with the target in question. Thus **15**, structurally related to androstenedione, a substrate for aromatase, was a moderate inhibitor of aromatase. Likewise the inhibition by **15** of testosterone 5 α -reductase might reflect its structural resemblance to the natural substrate testosterone, whereas **22**, correspondingly related to the product 5 α -dihydrotestosterone, was not an inhibitor. Notably, compound **3** inhibited neither aromatase nor testosterone 5 α -reductase at the highest concentration tested, respectively 20 and 50 μ M. Lastly, the estradiol-related analog **18** had an appreciable binding affinity for the estrogen receptor, 5% of that of estradiol itself.

Concluding Remarks

Two of the compounds described here, namely **2** (as a prodrug for **3**) and **15**, have been evaluated *in vivo* in the WHT mouse.³⁶ Each markedly reduced the weights of androgen-dependent organs, and **2** depressed testosterone to undetectable levels. The adrenals were unaffected, implying that **3** and **15**, unlike ketoconazole, do not inhibit enzymes in the pathway leading to corticosterone. This evidence for selective inhibition of testosterone biosynthesis, together with the further evidence for selectivity of action provided here for **3** in particular, makes **3** a strong candidate for further development as a potential drug for the treatment of prostatic carcinoma in humans.

Experimental Section

Chemical Methods. ¹H-NMR spectra (250 MHz) (internal Me₄Si = δ 0) were determined in CDCl₃ (unless otherwise indicated) using a Bruker AC 250 spectrometer. Infrared spectra were determined with a Perkin-Elmer 1720X spectrometer. Mass spectra (electron impact, 70 eV) were obtained by direct insertion with a VG 7070H spectrometer and VG 2235 data system. Melting points were determined with a Reichert micro hot stage apparatus and are uncorrected. Chromatography refers to column chromatography on silica gel (Merck Art. 15111) with the solvent indicated applied under positive pressure. Light petroleum refers to the fraction with bp 60–80 °C. 3-Pyridyl(diethyl)borane was purchased from Aldrich Chemical Co., Gillingham, Dorset, U.K. Elemental analyses were determined by CHN Analysis Ltd., South Wigston, Leicester, England.

3 β -Acetoxyandrosta-5,16-dien-17-yl Trifluoromethanesulfonate (1) and Androsta-3,5,16-trien-17-yl Trifluoromethanesulfonate (4). To a stirred solution of dehydroepiandrosterone 3-acetate (24.8 g, 75 mmol) in dry CH₂Cl₂ (500 mL) containing 2,6-di-*tert*-butyl-4-methylpyridine (18.5 g, 90 mmol) was added trifluoromethanesulfonic anhydride (12.6 mL, 75 mmol). After 12 h the mixture was filtered, washed with water (50 mL), and dried (MgSO₄) and the solvent evaporated. Chromatography, on elution with CH₂Cl₂–light petroleum (1:6), gave first **4** (3.02 g, 10%) as an oil: ν_{\max} for C=O str absent; ¹H NMR δ 0.99 (s, 3, H-18), 1.02 (s, 3, H-19), 5.39 (m, 1, H-6), 5.59 (m, 1, H-16), 5.62 (m, 1, H-3), 5.93 (dm, 1, *J* = 9.4 Hz, H-4); *m/z* 402 (M⁺).

Further elution with CH₂Cl₂–light petroleum (1:3) afforded **1** (20.1 g, 58%): mp 75–76 °C (from hexane); ν_{\max} 1734 cm⁻¹ (C=O str); ¹H NMR δ 1.00 (s, 3, H-18), 1.06 (s, 3, H-19), 2.04 (s, CH₃CO), 4.59 (m, 1, H-3 α), 5.39 (dm, 1, *J* = 4.9 Hz, H-6), 5.58 (m, 1, H-16); *m/z* 402 (M⁺ – AcOH). Anal. (C₂₂H₂₉O₅F₃S) C, H, F, S.

3 β -Acetoxy-17-(3-pyridyl)androsta-5,16-diene (2). Diethyl(3-pyridyl)borane (3.3 g, 23 mmol) was added to a stirred

solution of **1** (6.94 g, 15 mmol) in THF (75 mL) containing bis-(triphenylphosphine)palladium(II) chloride (0.105 g, 0.15 mmol). An aqueous solution of Na₂CO₃ (2 M, 30 mL) was then added and the stirred mixture heated at 80 °C for 1 h and then partitioned between Et₂O and H₂O. The organic phase was dried (Na₂CO₃), filtered through a short column of silica gel, and concentrated. Chromatography, on elution with Et₂O–light petroleum (1:2), afforded **2** (4.95 g, 84%): mp 144–145 °C (from hexane); ν_{\max} 1732 cm⁻¹ (C=O str); ¹H NMR δ 1.05 (s, 3, H-19), 1.08 (s, 3, H-18), 2.04 (s, 3, CH₃CO), 4.60 (m, 1, H-3 α), 5.42 (dm, 1, *J* = 4.7 Hz, H-6), 5.99 (m, 1, H-16), 7.23 (dd, 1, *J*_{5,4} = 8.1 Hz, *J*_{5,6} = 3.9 Hz, pyridyl H-5), 7.65 (ddd, 1, *J*_{4,2} = 2.0 Hz, *J*_{4,6} = 1.6 Hz, pyridyl H-4), 8.46 (dd, 1, pyridyl H-6), 8.62 (d, 1, pyridyl H-2); *m/z* 392 (M⁺ + H). Anal. (C₂₆H₃₃NO₂) C, H, N.

17-(3-Pyridyl)androsta-5,16-dien-3 β -ol (3). To a solution of **2** (4.90 g, 12.5 mmol) in methanol (50 mL) was added 2.5 M NaOH (10 mL), and the mixture was stirred at 80 °C for 5 min and then allowed to cool, poured into water, neutralized with 1 M HCl, reacidified with saturated aqueous NaHCO₃, and extracted with hot toluene (3 \times 100 mL). The toluene extracts were dried (Na₂CO₃) and concentrated. Chromatography, on elution with Et₂O–toluene (1:2), gave **3** (3.45 g, 79%): mp 228–229 °C (from toluene); ν_{\max} 3351 (OH str); ¹H NMR δ 1.05 (s, 3, H-19), 1.07 (s, 3, H-18), 3.54 (m, 1, H-3 α), 5.40 (dm, 1, *J* = 5.0 Hz, H-6), 5.99 (m, 1, H-16), 7.22 (dd, 1, pyridyl H-5), 7.65 (ddd, 1, pyridyl H-4), 8.46 (dd, 1, pyridyl H-6), 8.62 (d, 1, pyridyl H-2); *m/z* 349 (M⁺). Anal. (C₂₄H₃₁NO) C, H, N.

17-(3-Pyridyl)androsta-3,5,16-triene (5). The method followed that described for **2** but used **4** (2.01 g, 5.0 mmol). Chromatography, on elution with CH₂Cl₂, gave **5** (1.39 g, 84%): mp 110–112 °C (from hexane); ¹H NMR δ 1.02 (s, 3, H-19), 1.07 (s, 3, H-18), 5.44 (m, 1, H-6), 5.61 (m, 1, H-3), 5.95 (dm, 1, *J* = 9.8 Hz, H-4), 6.01 (m, 1, H-16), 7.23 (dd, 1, pyridyl H-5), 7.66 (ddd, 1, pyridyl H-4), 8.46 (dd, 1, pyridyl H-6), 8.63 (d, 1, pyridyl H-2); *m/z* 331 (M⁺). Anal. (C₂₄H₂₉N) H, N; C: calcd 86.96; found, 86.24.

3 β -Acetoxy-17-(2-pyridyl)androsta-5,16-diene (6). To Et₂O (6 mL), at –18 °C, was added *n*-butyllithium (0.96 mL, 2.5 M solution in hexanes) followed dropwise by 2-bromopyridine (0.228 mL, 2.4 mmol) in Et₂O (2 mL). The resulting blood-red solution of 2-pyridyllithium was added dropwise to a solution of ZnCl₂ (382 mg, 2.8 mmol) in THF, cooled to –18 °C, and the orange-brown solution of 2-pyridylzinc chloride was stirred for a further 30 min. For the preparation of the palladium catalyst, a solution of tetrakis(triphenylphosphine)palladium(0) (1.16 g, 1 mmol) in benzene (10 mL) was treated with 2-bromopropene (0.18 mL, 242 mg, 2 mmol) and the mixture stirred for 16 h at ambient temperature, whereupon the initially orange suspension became a yellow solution. The solvent was removed under vacuum, the residue was triturated with Et₂O, and the pale yellow product (0.70 g) bromo-(isopropenyl)bis(triphenylphosphine)palladium(II) was recovered by filtration: ¹H NMR δ 0.81 (s, CH₃), 4.6 (m, C=CH₂), 7.2–7.8 (m, arom H).

To a solution of **1** (926 mg, 1 mmol) in THF (10 mL) containing the palladium catalyst (76 mg, ca. 0.1 mmol) was added the solution of 2-pyridylzinc chloride, and the mixture was stirred at ambient temperature. After 1 h, the mixture was partitioned between Et₂O and H₂O and the organic phase was dried (Na₂CO₃) and concentrated. Chromatography, on elution with Et₂O–light petroleum (1:4), gave **6** (0.583 g, 74%): mp 189–190 °C (from light petroleum); ν_{\max} 1734 cm⁻¹ (C=O str); ¹H NMR δ 1.09 (s, 3, H-19), 1.15 (s, 3, H-18), 2.04 (s, 3, CH₃CO), 4.62 (m, 1, H-3 α), 5.42 (dm, 1, H-6), 6.37 (m, 1, H-16), 7.09 (dd, 1, *J*_{5,4} = 7.9 Hz, *J*_{5,6} = 4.1 Hz, pyridyl H-5), 7.38 (d, 1, *J*_{3,4} = 7.9 Hz, pyridyl H-3), 7.59 (t, *J* = 7.7 Hz, 1, pyridyl H-4), 8.55 (d, 1, pyridyl H-6); *m/z* 391 (M⁺). Anal. (C₂₆H₃₃NO₂) C, H, N.

3 β -Acetoxy-17-(4-pyridyl)androsta-5,16-diene (7). 4-Bromopyridine (4.5 g) was liberated from its hydrochloride (5 g, 26 mmol) using the procedure previously applied to 4-chloropyridine³⁷ but keeping solutions below 10 °C during concentration to prevent polymerization. The free base was twice concentrated from Et₂O (to remove residual CHCl₃), and then a solution of the freshly prepared 4-bromopyridine (1.58 g, 10

mmol) in Et₂O (8 mL) was added dropwise during 30 min to a mixture of *n*-butyllithium (2.5 M in hexanes, 4 mL, 10 mmol) and Et₂O (20 mL) at -20 °C. Of the resulting solution of 4-pyridyllithium, 15 mL (4.5 mmol) was added to a stirred solution of anhydrous ZnCl₂ (681 mg, 5 mmol) in dry THF (25 mL). After 1 h at ambient temperature, 30 mL of the red solution of 4-pyridylzinc chloride was added to a solution of **1** (925 mg, 2 mmol) in THF containing bromo(isopropenyl)bis(triphenylphosphine)palladium(II) (see **6** above; 75 mg, 0.1 mmol) and the mixture stirred at ambient temperature overnight and then filtered through Celite. Chromatography, on elution with Et₂O–light petroleum (1:2), gave **7** (140 mg, 18%): mp 175–177 °C (from light petroleum); ν_{\max} 1732 cm⁻¹ (C=O str); ¹H NMR δ 1.08 (s, 3, H-19), 1.63 (s, 3, H-18), 2.05 (s, 3, CH₃CO), 4.63 (m, 1, H-3 α), 5.42 (dm, 1, H-6), 6.18 (m, 1, H-16), 7.26 (d, 2, *J* = 6.0 Hz, pyridyl H-3, H-5), 8.50 (d, 2, pyridyl H-2, H-6); *m/z* 331 (M⁺ – AcOH). Anal. (C₂₆H₃₃NO₂) C, H, N.

17-(2-Pyridyl)androsta-5,16-dien-3 β -ol (9). The method followed that described for **3** but used **6** (392 mg, 1 mmol), except that on completion of the reaction the product was extracted with Et₂O followed by benzene and crystallized without prior chromatography giving **9** (273 mg, 78%): mp 206–207 °C (from benzene–light petroleum); ν_{\max} 3390 cm⁻¹ (OH str); ¹H NMR δ 1.08 (s, 3, H-18), 1.15 (s, 3, H-19), 3.56 (m, 1, H-3 α), 5.41 (m, 1, H-6), 6.38 (m, 1, H-16), 7.10 (m, 1, pyridyl H-5), 7.38 (d, 1, *J* = 7.8 Hz, pyridyl H-3), 7.59 (m, 1, pyridyl H-4), 8.55 (d, 1, *J* = 4.2 Hz, pyridyl H-6); *m/z* 349 (M⁺). Anal. (C₂₄H₃₁NO) C, H, N.

17-(4-Pyridyl)androsta-5,16-dien-17-ol (10). A solution of 4-bromopyridine (from the hydrochloride; 25 g, 129 mmol; see **7** above) in Et₂O (80 mL) was added dropwise to a stirred mixture of *n*-butyllithium (51.6 mL, 2.5 M in hexanes, 129 mmol) and Et₂O (200 mL) at -76 °C. The resulting solution of 4-pyridyllithium was added by transfer needle to a cooled (-76 °C) solution of trimethyl borate (13.4 g, 14.6 mL, 129 mmol) in Et₂O (75 mL); the mixture was stirred for 20 min and then allowed to reach ambient temperature. Water (10 mL) was added, and the resulting light brown precipitate of lithium trimethoxy(4-pyridyl)boronate (22.04 g, ca. 90%) was collected by filtration, washed with Et₂O, and dried *in vacuo*. This product (2.83 g, ca. 15 mmol) was added to a solution of **1** (1.21 g, 5 mmol) in THF (30 mL) containing bis(triphenylphosphine)palladium(II) chloride (175 mg, 0.25 mmol), followed by 2 M aqueous Na₂CO₃ (12.5 mL), and the mixture heated at 80 °C, for 6 h, and then partitioned between Et₂O and H₂O. The organic phase was dried (Na₂CO₃) and filtered through a short column of silica gel to give crude **7** which was used for the next step without further purification. The method for converting **7** into **10** followed that described for **2** → **3**. Chromatography, eluting with Et₂O–toluene (1:2), gave **10** (928 mg, 53% from **1**): mp 226–228 °C (from toluene); ¹H NMR δ 1.08 (s, 3, H-19), 1.62 (s, 3, H-18), 3.55 (m, 1, H-3 α), 5.40 (dm, 1, H-6), 6.18 (m, 1, H-16), 7.26 (d, 2, *J* = 6.1 Hz, pyridyl H-3, H-5), 8.51 (d, 2, pyridyl H-2, H-6). Anal. (C₂₄H₃₁NO) C, H, N.

17-(2-Pyridylmethyl)androsta-5,16-dien-17-ol (11). To a solution of 2-picoline (7.45 g, 7.9 mL, 80 mmol) in THF (42 mL) at -20 °C was added *n*-butyllithium (50 mL, 1.6 M in hexanes, 80 mmol) during 30 min. The red solution of 2-picolyllithium³⁸ (10 mL) was added with vigorous stirring under argon to anhydrous ZnCl₂ (1.09 g, 8 mmol), followed by benzene (10 mL). The resulting homogeneous solution of 2-picolylzinc chloride (15 mL) was added to a solution of **1** (925 mg, 2 mmol) in THF (6 mL) containing bis(triphenylphosphine)palladium(II) chloride (70 mg, 0.1 mmol), and the resulting yellow solution was heated at 70 °C for 2 h and then partitioned between Et₂O and H₂O. The organic phase was concentrated and the crude 3 β -acetoxy-17-(2-pyridylmethyl)-androsta-5,16-diene (**8**) (650 mg) used directly for the next step. The method followed that described for **2** → **3**. Chromatography, eluting with Et₂O–light petroleum (1:1), gave **11** (460 mg, 79%): mp 86–88 °C (from light petroleum–toluene); ν_{\max} 3330 cm⁻¹ (OH str); ¹H NMR δ 0.82 (s, 3, H-19), 1.04 (s, 3, H-18), 3.5 (s + m, 3, benzyl H + H-3 α), 5.10 (m, 1, H-16), 5.35 (m, 1, H-6), 7.12 (dd, 1, *J*_{5,4} = 6.5 Hz, *J*_{5,6} = 4.7 Hz, pyridyl H-5), 7.24 (d, 1, *J*_{3,4} = 7.8 Hz, pyridyl H-3), 7.62 (dd, 1, pyridyl

H-4), 8.54 (d, 1, pyridyl H-6); *m/z* 363 (M⁺). Anal. (C₂₅H₃₃NO·H₂O) C, H, N.

3-[2,3,5,6-Tetrafluoro-4-(trifluoromethyl)phenoxy]androsta-3,5,16-trien-17-yl Trifluoromethanesulfonate (13). The method followed that described for **1** but used 3-[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]androsta-3,5-dien-17-one²⁷ (**12**; 5.03 g, 10 mmol). Chromatography, on elution with CH₂Cl₂–light petroleum (1:10), gave **13** (1.93 g, 30%): mp 106–107 °C (from EtOH); ¹H NMR δ 1.02 (s, 6, H-18, H-19), 5.16 (s, 1, H-4), 5.28 (m, 1, H-6), 5.59 (m, 1, H-16); *m/z* 634 (M⁺). Anal. (C₂₇H₂₄F₁₀O₄S) H, S; C: calcd, 51.10; found, 50.61. F: calcd, 29.94; found, 29.14.

3-[2,3,5,6-Tetrafluoro-4-(trifluoromethyl)phenoxy]-17-(3-pyridyl)androsta-3,5,16-triene (14). The method for **2** was followed, using **13** (1.27 g, 2.0 mmol). Chromatography, on elution with Et₂O–light petroleum (1:3), gave **14** (0.82 g, 73%): mp 166–166.5 °C (from hexane); ¹H NMR δ 1.05 (s, 3, H-19), 1.07 (s, 3, H-18), 5.18 (s, 1, H-4), 5.32 (m, 1, H-6), 6.01 (m, 1, H-16), 7.23 (dd, 1, pyridyl H-5), 7.66 (ddd, 1, pyridyl H-4), 8.47 (dd, 1, pyridyl H-6), 8.63 (d, 1, pyridyl H-2); *m/z* 563 (M⁺). Anal. (C₃₁H₂₈F₇NO) C, H, F, N.

17-(3-Pyridyl)androsta-4,16-dien-3-one (15). (a) From **14**. To a solution of **14** (0.423 g, 0.75 mmol) in THF (5 mL) was added EtOH (5 mL) followed by 1 M HCl (5 mL), and the mixture was heated with stirring at 65 °C for 48 h. The mixture was then poured into H₂O (20 mL), neutralized with 1 M NaOH, and extracted with Et₂O (3 × 30 mL). The ether extracts were combined, dried (Na₂CO₃), and concentrated. Chromatography, on elution with Et₂O, gave **15** (185 mg, 71%): mp 148–150 °C (from Et₂O); ν_{\max} 1674 cm⁻¹ (C=O str); ¹H NMR δ 1.07 (s, 3, H-18), 1.24 (s, 3, H-19), 5.76 (s, 1, H-4), 5.99 (m, 1, H-16), 7.23 (dd, 1, pyridyl H-5), 7.64 (ddd, 1, pyridyl H-4), 8.47 (d, 1, pyridyl H-6), 8.62 (d, 1, pyridyl H-2); *m/z* 347 (M⁺).

(b) From **3**, by Oppenauer Oxidation. From a solution of **3** (4.17 g, 12 mmol) in dry toluene (300 mL) and cyclohexanone (60 mL) was distilled off part of the solvent (80 mL) to eliminate moisture. After allowing to cool to 90 °C, Al(O-*i*-Pr)₃ (4.08 g, 20 mmol) was added and the mixture heated under reflux for 90 min and then allowed to cool, diluted with Et₂O (250 mL), washed with aqueous trisodium citrate (15%, w/v; 2 × 30 mL), dried (Na₂CO₃), and concentrated. Chromatography, on elution with MeOH–toluene (1:20), afforded **15** (3.4 g, 82%), identical to the product obtained by method a above. Anal. (C₂₄H₂₉NO) C, N; H: calcd, 8.09; found 7.67.

3-Acetoxyestra-1,3,5[10],16-tetraen-17-yl Trifluoromethanesulfonate (16). The method followed that described for **1** but used estrone 3-acetate (4.69 g, 15 mmol). Chromatography, on elution with CH₂Cl₂–light petroleum (1:3), gave **16** (5.21 g, 78%) as a waxy solid; ν_{\max} 1766 cm⁻¹ (C=O str); ¹H NMR δ 1.00 (s, 3, H-18), 2.29 (s, 3, CH₃CO), 5.62 (m, 1, H-16), 6.81 (bs, 1, arom H-4), 6.85 (dd, 1, *J*_{2,1} = 8.5 Hz, *J*_{2,4} = 2.6 Hz, arom H-2), 7.26 (d, 1, arom H-1); *m/z* 445 (M⁺ + H).

3-Acetoxy-17-(3-pyridyl)estra-1,3,5[10],16-tetraene (17). The method followed that described for **2** but used **16** (3.56 g, 8.0 mmol). Chromatography, on elution with Et₂O–light petroleum (1:2), gave **17** (2.40 g, 80%) as an oil; ν_{\max} 1738 cm⁻¹ (C=O str); ¹H NMR δ 1.04 (s, 3, H-18), 2.29 (s, 3, CH₃CO), 6.03 (m, 1, H-16), 6.82 (s, 1, arom H-4), 6.85 (d, 1, *J*_{2,1} = 8.4 Hz, arom H-2), 7.24 (m, 1, pyridyl H-5), 7.29 (d, arom H-1), 7.69 (m, 1, pyridyl H-4), 8.48 (dd, 1, pyridyl H-6), 8.65 (d, 1, pyridyl H-2); *m/z* 373 (M⁺).

17-(3-Pyridyl)estra-1,3,5[10],16-tetraen-3-ol (18). The method followed that described for **3** but used **17** (2.36 g, 6.3 mmol). Chromatography, on elution with MeOH–toluene (1:8), gave **18** (1.40 g, 67%): mp 256–258 °C (from toluene); ¹H NMR (Me₂SO-*d*₆) δ 1.01 (s, 3, H-18), 6.15 (m, 1, H-16), 6.47 (s, 1, arom H-4), 6.52 (d, 1, *J*_{2,1} = 8.4 Hz, arom H-2), 7.04 (d, 1, arom H-1), 7.35 (dd, 1, pyridyl H-5), 7.79 (m, 1, pyridyl H-4), 8.45 (d, 1, pyridyl H-6), 8.62 (s, 1, pyridyl H-2). Anal. (C₂₈H₂₅NO) C, H, N.

3 α -Acetoxy-17-(3-pyridyl)-5 α -androst-16-ene (20). The method followed that described for **2** but used 3 α -acetoxy-5 α -androst-16-en-17-yl trifluoromethanesulfonate (**19**; 3.44 g, 7.4 mmol), prepared from 3 α -acetoxy-5 α -androst-17-one as described for **1**. Chromatography, on elution with Et₂O–light petroleum (1:2), gave **20** as an oil (2.39 g, 82%): ¹H NMR δ

0.85 (s, 3, H-19), 1.01 (s, 3, H-18), 2.06 (s, 3, CH₃CO), 5.02 (m, 1, H-3β), 6.00 (m, 1, H-16), 7.24 (dd, 1, pyridyl H-5), 7.68 (ddd, 1, pyridyl H-4), 8.47 (dd, pyridyl H-6), 8.63 (dd, 1, pyridyl H-2); *m/z* 393 (M⁺).

17-(3-Pyridyl)-5α-androst-16-en-3α-ol (21). The method followed that described for **3** but used **20** (2.33 g, 5.9 mmol). Chromatography, on elution with MeOH-toluene (1:40), gave **21** (1.62 g, 78%): mp 198–199 °C (from toluene); ¹H NMR δ 0.84 (s, 3, H-19), 1.00 (s, 3, H-18), 4.06 (bs, 1, H-3β), 5.97 (m, 1, H-16), 7.21 (dd, 1, pyridyl H-5), 7.64 (ddd, 1, pyridyl H-4), 8.45 (dd, 1, pyridyl H-6), 8.61 (d, 1, pyridyl H-2); *m/z* 351 (M⁺). Anal. (C₂₃H₃₁NO) C, H, N.

17-(3-Pyridyl)-5α-androst-16-en-3-one (22). The method essentially followed method b for the formation of **15**, from **3** by Oppenauer oxidation, but used **21** (1.05 g, 3.0 mmol). Chromatography, on elution with MeOH-toluene (1:40), gave **22** (0.90 g, 86%): mp 190–192 °C (from toluene); *ν*_{max} 1713 cm⁻¹ (C=O str); ¹H NMR δ 1.04 (s, 3, H-19), 1.07 (s, 3, H-18), 5.98 (m, 1, H-16), 7.22 (dd, 1, pyridyl H-5), 7.64 (ddd, 1, pyridyl H-4), 8.46 (dd, 1, pyridyl H-6), 8.61 (d, 1, pyridyl H-2); *m/z* 349 (M⁺). Anal. (C₂₄H₂₉NO) H, N; C: calcd, 82.47; found, 82.00.

3-(tert-Butyldimethylsiloxy)androsta-2,4-diene-11,17-dione (23). To a solution of adrenosterone (6.0 g, 20 mmol) in dry CH₂Cl₂ (120 mL) was added Et₃N (8.4 mL, 60 mmol) followed by *tert*-butyldimethylsilyl trifluoromethanesulfonate (5.0 mL, 22 mmol), and the mixture was stirred at ambient temperature for 3 h and concentrated and the residue dissolved in Et₂O (100 mL). After a further 30 min, the supernatant was decanted from the oil which had separated and residual solvent was removed under reduced pressure to give crude **23** which was used directly in the following step: *ν*_{max} 1705, 1747 cm⁻¹ (C=O str); ¹H NMR δ 0.12 (s, 6, Me₂Si), 0.85 (s, 3, H-18), 0.92 (s, 9, *t*-BuSi), 1.17 (s, 3, H-19), 4.73 (dm, 1, J = 6.9 Hz, H-2), 5.36 (app t, J = 2.1 Hz, 1, H-4); *m/z* 373 (M⁺ - C₃H₇).

3-(tert-Butyldimethylsiloxy)-11-oxoandrosta-2,4,16-trien-17-yl Trifluoromethanesulfonate (24). To a solution of the crude **23** in dry THF (100 mL), cooled to -78 °C, was added a freshly prepared solution of lithium diisopropylamide [prepared by adding *n*-butyllithium (1.6 M in hexane, 13.8 mL, 22 mmol) to a solution of diisopropylamine (3.1 mL, 22 mmol) in dry THF (25 mL) at -18 °C], and the resultant yellow solution was stirred at -78 °C for 30 min. A solution of *N*-phenyltrifluoromethanesulfonamide (7.15 g, 20 mmol) in dry THF (20 mL) was then added, and after an additional 1 h at -78 °C, the mixture was allowed to reach ambient temperature. The reaction mixture was poured into water (200 mL) and extracted with Et₂O (2 × 200 mL). The extracts were washed with H₂O (20 mL), dried (Na₂CO₃), and concentrated to give crude **24** which was used directly in the next step: *ν*_{max} 1710 cm⁻¹ (C=O str); ¹H NMR δ 0.13 (s, 6, Me₂Si), 0.92 (s, 9, *t*-BuSi), 1.35 (2s, 6, H-18, H-19), 4.75 (m, 1, H-2), 5.38 (bs, 1, H-4), 5.68 (m, 1, H-16); *m/z* 505 (M⁺ - C₃H₇).

3-(tert-Butyldimethylsiloxy)-17-(3-pyridyl)androsta-2,4,16-trien-11-one (25). The method essentially followed that described for **2** but used crude **24**. The crude **25** so obtained was used directly in the following step: *ν*_{max} 1705 cm⁻¹ (C=O str); ¹H NMR δ 0.13 (s, 6, Me₂Si), 0.93 (s, 9, *t*-BuSi), 0.99 (s, 3, H-18), 1.18 (s, 3, H-19), 4.75 (dm, 1, H-2), 5.37 (app t, J = 2.1 Hz, 1, H-4), 6.09 (m, 1, H-16), 7.26 (dd, 1, pyridyl H-5), 7.62 (ddd, 1, pyridyl H-4), 8.50 (dd, 1, pyridyl H-6), 8.60 (d, 1, pyridyl H-2); *m/z* 475 (M⁺).

17-(3-Pyridyl)androsta-4,16-diene-3,11-dione (26). To a solution of the crude **25** in wet THF (60 mL) was added a solution of tetra-*n*-butylammonium fluoride (1.0 M, 10 mL, 10 mmol) in THF, and the mixture was stirred at room temperature for 12 h and then partitioned between Et₂O and H₂O and basified with saturated aqueous NaHCO₃ and the organic phase dried (Na₂CO₃) and concentrated. Chromatography, on elution with Et₂O, gave **26** (4.30 g, 60% overall yield from adrenosterone); mp 181–183 °C (from Et₂O); *ν*_{max} 1669, 1703 cm⁻¹ (C=O str); ¹H NMR δ 1.02 (s, 3, H-18), 1.45 (s, 3, H-19), 5.76 (s, 1, H-4), 6.08 (m, 1, H-16), 7.24 (dd, 1, pyridyl H-5), 7.59 (ddd, 1, pyridyl H-4), 8.50 (dd, 1, pyridyl H-6), 8.59 (d, 1, pyridyl H-2); *m/z* 361 (M⁺). Anal. (C₂₄H₂₆NO₂) C, H, N.

17β-(3-Pyridyl)androst-5-en-3β-ol (27). To a solution of **3** (350 mg, 1 mmol) in EtOH (60 mL) were added hydrazine

hydrate (1.6 mL, 5 mmol) and AcOH (1 mL). The mixture was heated at 80 °C for 16 h while a stream of air was passed through the solution and then partitioned between toluene and aqueous NaHCO₃. Chromatography of the organic extract, on elution with MeOH-toluene (1:40), gave **27** (290 mg, 83%) as white crystals: mp 217–218 °C (from toluene); *ν*_{max} 3368 cm⁻¹ (OH str) ¹H NMR δ 0.44 (s, 3, H-18), 0.95 (s, 3, H-19), 2.62 (app t, 1, J = 9.6 Hz, H-17α), 3.50 (m, 1, H-3α), 5.32 (m, 1, H-6), 7.18 (dd, 1, J_{5,4} = 7.8 Hz, J_{5,6} = 4.8 Hz, pyridyl H-5), 7.50 (ddd, 1, pyridyl H-4), 8.39 (m, 2, pyridyl H-2, H-6); *m/z* 351 (M⁺). Anal. (C₂₄H₃₃NO) H, N; C: calcd, 82.00; found, 81.58.

17β-(4-Pyridyl)androst-5-en-3β-ol (28). To a suspension of **10** (699 mg, 2 mmol) in dry THF (10 mL) was added dropwise by syringe a solution of Red-Al (2.4 mL, 3.4 M in toluene), then ZnCl₂ (545 mg, 4 mmol) was added, and the mixture was stirred at ambient temperature for 2 h. Excess hydride was quenched by dropwise addition of H₂O, and then the mixture was poured into water and extracted first with Et₂O and then with hot toluene. The extracts were concentrated, and the white crystalline residue was recrystallized from hot toluene to give slightly impure **28** (672 mg, 96%) which was purified by further recrystallization from toluene: mp 272–273 °C; ¹H NMR δ 0.47 (s, 3, H-18), 1.00 (s, 3, H-19), 2.65 (app t, 1, J = 9.6 Hz, H-17α), 3.54 (m, 1, H-3α), 5.37 (m, 1, H-6), 7.12 (d, 2, J = 6.0 Hz, pyridyl H-3, H-5), 8.48 (d, 2, pyridyl H-2, H-6). Anal. (C₂₄H₃₃NO) C, H, N.

Enzyme Preparation and Assay Procedure for the 17α-Hydroxylase-C_{17,20}-lyase Enzyme. The ³H-labeled compounds were obtained from NEN Products, Stevenage, Herts, U.K. The biochemical reagents were from Boehringer Mannheim, U.K., Lewes, East Sussex, U.K., or Sigma Chemical Co. Ltd., Poole, Dorset, U.K. The chemicals were of analytical grade.

A microsomal fraction was prepared by the method of Chasalow³⁹ from human testes removed at orchietomy from previously untreated patients with cancer of the prostate. The microsomes were resuspended in 50 mM sodium phosphate buffer (pH 7.4)-glycerol (3:1) at the equivalent of 1 mL/g of fresh tissue and stored in liquid nitrogen until use.

The assay was based on that of Chasalow,⁴⁰ and the assay mixture contained 3 μM ³H-labeled substrate (1–3 μCi/nmol), 250 μM NADPH, 10 mM D-glucose 6-phosphate, 1 mM MgCl₂, 2 U/mL D-glucose 6-phosphate dehydrogenase, 0.1 mM dithiothreitol, 0.2 mM EDTA, 1% ethanol, 1% DMSO, 3% glycerol, and 95% 50 mM sodium phosphate buffer (pH 7.4). The test compounds were prepared in 50% DMSO, the controls receiving just 50% DMSO. The reaction was carried out at 37 °C. It was started by the addition of the microsomal preparation and stopped by the addition of 2 volumes of MeCN-MeOH (1:2) containing unlabeled steroids (ca. 100 μM). The samples were stored at -20 °C until analysis. The reaction was linear with time, and the rate was proportional to the protein concentration under the conditions used (data not shown).

HPLC Analysis. (a) Hydroxylase Activity. For measurement of the hydroxylase activity, the substrate was progesterone and the unlabeled steroids added at the end of the assay were progesterone, 17α-hydroxyprogesterone, androstenedione, testosterone, and 16α-hydroxyprogesterone. The samples were injected onto a 10 cm Nucleosil 5-μM C18 column fitted with an Uptight guard column filled with Nucleosil C18 packing. The mobile phase was 60% MeOH at a flow rate of 1 mL min⁻¹. The effluent was monitored at 240 nm before being mixed with Ecoscint A containing 25% MeCN and monitored for ³H using a Berthold LB506C detector. Activity was measured as the production of 17α-hydroxyprogesterone. No androstenedione nor testosterone were produced until the substrate was depleted, and the reaction was not carried out long enough for this to occur.

(b) Lyase Activity. For the measurement of the C_{17,20}-lyase activity, the substrate was 17α-hydroxyprogesterone and the unlabeled steroids added at the end of the assay were 17α-hydroxyprogesterone, androstenedione, and testosterone. The samples were injected onto a 10 cm Apex 5 μM C18 column fitted with an Uptight guard column filled with PELL ODS packing. The mobile phase was H₂O-MeCN-MeOH (4:1:3) at a flow rate of 1 mL min⁻¹. The effluent was monitored at

240 nm before being mixed with Ecoscint A containing 5% MeCN, 5% MeOH and monitored for ^3H using a Berthold LB506C detector. Activity was measured as the production of androstenedione and testosterone.

Inhibitory Activity. For ease of dissolution, test compounds were first converted into their hydrochlorides. In a typical procedure, HCl gas was passed through a solution of the base in Et_2O and the hydrochloride which precipitated was recovered by filtration and desiccated. Variations on the procedure included the use of toluene (**22**) or MeOH–toluene (**18**) as solvent followed by addition of Et_2O to precipitate the salt. Each compound was tested at a minimum of four different concentrations, and the data were fitted by nonlinear regression to the median effect equation of Chou:⁴¹

$$f_a/f_u = (I/IC_{50})^n$$

where f_a = the fraction of activity affected, f_u = the fraction of activity unaffected, I = the concentration of inhibitor, IC_{50} = inhibitor concentration giving 50% inhibition, and n depends on the sigmoid shape of the curve ($n = 1$ for systems obeying Michaelis–Menten kinetics). The correlation coefficients were all greater than 0.96. This method of analysis was chosen as it is valid for calculating IC_{50} values whatever the IC_{50} /(enzyme concentration) ratio. In contrast, methods based on the Michaelis–Menten equation become invalid for values of $IC_{50} < 100$ (enzyme concentration).⁴²

The estimates of the enzyme concentration were obtained by fitting some of the data to the equation derived by Henderson⁴³ for tight binding inhibitors. For a tight binding inhibitor:

$$IC_{50} = K_{i,app} + 0.5(\text{enzyme concentration})$$

Other Biological Activities. The reagents and conditions for the assays for inhibition of aromatase enzyme from the microsomal fraction of human placenta⁴⁴ and of testosterone 5α -reductase from human benign prostatic tissue⁴⁵ were as previously described. The estrogen receptor binding assay, using immature rat uterine cytosol, was a modification of that described by Wakeling⁴⁶ and is described elsewhere.⁴⁷

Acknowledgment. This work was supported by grants from the Cancer Research Campaign and the Medical Research Council. Financial support (to S.E.B. and G.A.P.) from the British Technology Group and (to S.E.B.) from Cancer Research Campaign Technology is also gratefully acknowledged. We thank Mr. J. Houghton and Mr. M. H. Baker for skilled technical assistance.

References

- Boring, C. C.; Squires, T. S.; Tong, T. Cancer Statistics. *CA Cancer J. Clin.* **1993**, *43*, 7–26.
- Smith, J. A. New Methods of Endocrine Management of Prostatic Cancer. *J. Urol.* **1987**, *137*, 1–10.
- Forti, G.; Salerno, R.; Moneti, G.; Zoppi, S.; Fiorelli, G.; Marinon, T.; Natali, A.; Constantini, A.; Serio, M.; Martini, L.; Motta, M. Three-Month Treatment with a Long Acting Gonadotrophin-releasing Hormone Agonist of Patients with Benign Prostatic Hyperplasia: Effects on Tissue Androgen Concentration, 5α -Reductase Activity and Androgen Receptor Content. *J. Clin. Endocrinol. Metab.* **1989**, *68*, 461–468.
- Crawford, E. D.; Eisenberger, M. A.; McLeod, D. G.; Spaulding, J. T.; Benson, R.; Dorr, F. A.; Blumenstein, B. A.; Davies, M. A.; Goodman, P. J. A Controlled Trial of Leuprolide With and Without Flutamide in Prostatic Carcinoma. *N. Engl. J. Med.* **1989**, *321*, 419–424.
- Labrie, F.; Belanger, A.; Simard, J.; Labrie, C.; Dupont, A. Combination Therapy for Prostate Cancer. *Cancer Suppl.* **1993**, *71*, 1059–1067.
- Rajfer, J.; Sikka, S. C.; Rivera, F.; Handelsman, D. J. Mechanism of Inhibition of Human Testicular Steroidogenesis by Oral Ketoconazole. *J. Clin. Endocrinol. Metab.* **1986**, *63*, 1193–1198.
- Trachtenberg, J.; Pont, A. Ketoconazole Therapy for Advanced Prostatic Cancer. *Lancet II* **1984**, 433–435.
- Van Wauwe, J. P.; Janssen, P. A. J. Is there a Case for P-450 Inhibitors in Cancer Treatment? *J. Med. Chem.* **1989**, *32*, 2231–2239.
- Jubelirer, S. J.; Hogan, T. High Dose Ketoconazole for the Treatment of Hormone Refractory Metastatic Prostate Carcinoma: 16 Cases and Review of the Literature. *J. Urol.* **1989**, *142*, 89–91.
- Hasselund, S.; Norman, N.; Sander, S. Ketoconazole High Dose in the Hormonal Treatment of Advanced Carcinoma of the Prostate. *Scand. J. Urol. Nephrol.* **1987**, *21*, 273–276.
- Muscato, J. J.; Ahmann, T. A.; Johnson, K. M.; Wilding, W.; Monaghan, G.; Schlossman, D. M. Optimal Dosing of Ketoconazole and Hydrocortisone Leads to Long Responses in Hormone Refractory Prostate Cancer. *Proc. Am. Assoc. Cancer Res.* **1994**, *13*, 22.
- McCague, R.; Rowlands, M. G.; Barrie, S. E.; Houghton, J. Inhibition of Enzymes of Estrogen and Androgen Biosynthesis by Esters of 4-Pyridylacetic Acid. *J. Med. Chem.* **1990**, *33*, 3050–3055.
- Laughton, C. A.; Neidle, S. Inhibitors of the P450 Enzymes Aromatase and Lyase. Crystallographic and Molecular Modelling Studies Suggest Structural Features of Pyridylacetic Acid Derivatives Responsible for Differences in Enzyme Inhibitory Activity. *J. Med. Chem.* **1990**, *33*, 3055–3060.
- Barrie, S. E.; Jarman, M.; Potter, G. A.; Rowlands, M. G. Preparation of Bridged Cycloalkyl 3-Pyridylalkanoates and Analogs as Hydroxylase/Lyase Inhibitors. U.K. Pat Appl. GB 2,253,851, 1992; *Chem. Abstr.* **1993**, *118*, 101815e.
- Hosaka, M.; Oshima, H.; Troen, P. Studies of the Human Testis. XIV. Properties of C17–C20 Lyase. *Acta Endocrinol.* **1980**, *94*, 389–396.
- Oh-e, T.; Miyaura, N.; Suzuki, A. Palladium-Catalyzed Cross-Coupling Reaction of Organoboron Compounds with Organic Triflates. *J. Org. Chem.* **1993**, *58*, 2201–2208.
- Terashima, M.; Kakimi, H.; Ishikura, M.; Kamata, K. Dialkyl-(3-pyridyl)boranes. *Chem. Pharm. Bull.* **1983**, *31*, 4573–4577.
- Ishikura, M.; Kamada, M.; Terashima, M. An Efficient Synthesis of 3-Heteroarylpyridines via Diethyl-(3-pyridyl)-borane. *Synthesis* **1984**, 936–938.
- Wicha, J.; Masnyk, M. Synthesis of 17β -Pyridyl- and 17β -Pyridonyl-androstane Derivatives. *Heterocycles* **1981**, *16*, 521–524.
- Wicha, J.; Masnyk, M.; Duddeck, H. Cardiotonic Steroids. Part 7. Synthesis of 17β -Pyridyl-androstane Derivatives. *Bull. Pol. Acad. Sci. Chem.* **1984**, *32*, 75–83.
- Potter, G. A. Applications of Organotransition Metal Chemistry to the Synthesis of Novel Anticancer Agents. Ph.D. Thesis, University of London, 1990; p 153.
- Potter, G. A.; McCague, R. Highly Stereoselective Access to an (E)-Vinyl Bromide from an Aryl Ketone Leads to Short Syntheses of (Z)-Tamoxifen and Important Substituted Derivatives. *J. Org. Chem.* **1990**, *55*, 6184–6187.
- Fitton, P.; McKeon, J. E.; Ream, B. C. Preparation of Benzylpalladium(II) Derivatives and their Reactions with Metal Acetates in Acetic Acid. *J. Chem. Soc., Chem. Commun.* **1969**, 370–371.
- Ishikura, M.; Ohta, T.; Terashima, M. A Novel Synthesis of 4-Aryl- and 4-Heteroarylpyridines via Diethyl(4-pyridyl)borane. *Chem. Pharm. Bull.* **1985**, *33*, 4755–4763.
- Fischer, F. C.; Havinga, E. Pyridineboronic Acids. *Recl. Trav. Chim. Pays-Bas* **1965**, *84*, 439–440.
- Cacchi, S.; Morera, E.; Ortari, G. Palladium-catalyzed Reduction of Vinyl Trifluoromethanesulfonates to Alkenes: Cholesta-3,5-diene. *Org. Synth.* **1990**, *68*, 138–147.
- Jarman, M.; McCague, R. The Selective Protection of the 3-Ketone Functions of Steroids as Heptafluoro-*p*-tolyl Enol Ethers. *J. Chem. Soc., Perkin Trans. 1* **1987**, 1129–1133.
- Tanabe, M.; Crowe, D. F. Generation and Reaction of 2,4-Dienolate Ions from Δ^4 -3-Keto-steroids with Lithium Hexamethyldisilazane. *J. Chem. Soc., Chem. Commun.* **1973**, 564–565.
- McMurry, J. E.; Scott, W. J. A Method for the Regiospecific Synthesis of Enol Triflates by Enolate Trapping. *Tetrahedron Lett.* **1983**, *24*, 979–982.
- Bridgeman, J. E.; Cherry, P. C.; Clegg, A. S.; Evans, J. M.; Ewart, E. R.; Kasal, A.; Kumar, V.; Meakins, G. D.; Morisawa, Y.; Richards, E. E.; Woodgate, P. D. Proton Magnetic Resonance Spectra of Ketones, Alcohols and Acetates in the Androstane, Pregnane, and Estrane Series. *J. Chem. Soc. C* **1970**, 250–257.
- Angelastro, M. R.; Laughlin, M. E.; Schatzman, G. L.; Bey, P.; Blohm, T. R. 17β -(Cyclopropylamino)androst-5-en- 3β -ol, a Selective Mechanism-based Inhibitor of Cytochrome P450_{17 α} (Steroid 17α -Hydroxylase/C_{17,20} Lyase). *Biochem. Biophys. Res. Commun.* **1989**, *162*, 1571–1577.
- Li, J.; Li, Y.; Son, C.; Banks, P.; Brodie, A. 4-Pregnene-3-one-20 β -carboxaldehyde: a Potent Inhibitor of 17α -Hydroxylase/C₁₇–20 Lyase and of 5α -Reductase. *J. Steroid Biochem. Mol. Biol.* **1992**, *42*, 313–320.
- Arth, G. E.; Patchett, A. A.; Jefopoulos, T.; Bugianesi, R. L.; Peterson, L. H.; Ham, E. A.; Kuehl, F. A., Jr.; Brink, N. G. Steroidal Androgen Biosynthesis Inhibitors. *J. Med. Chem.* **1971**, *14*, 675–680.

- (34) Goldman, A. S.; Eavey, R. D.; Baker, M. K. Production of Male Pseudohermaphroditism in Rats by Two New Inhibitors of Steroid 17 α -hydroxylase and C 17–20 Lyase. *J. Endocrinol.* **1976**, *71*, 289–297.
- (35) Vanden Bossche, H.; Willemsens, G.; Bellens, D.; Roels, I.; Janssen, P. A. J. From 14 α -Demethylase Inhibitors in Fungal Cells to Androgen and Oestrogen Biosynthesis Inhibitors in Mammalian Cells. *Biochem. Soc. Trans.* **1990**, *18*, 10–13.
- (36) Barrie, S. E.; Potter, G. A.; Goddard, P. M.; Haynes, B. P.; Dowsett, M.; Jarman, M. Pharmacology of Novel Steroidal Inhibitors of Cytochrome P450_{17α} (17 α -Hydroxylase/C17–20 Lyase). *J. Steroid Biochem. Mol. Biol.* **1994**, *50*, 267–273.
- (37) Chan, F. C. Y.; Potter, G. A.; Jarman, M. The Direct Formation of Esters of 2-(4-Pyridyl)propanoic Acid by Reaction of a Meldrum's Acid Derivative with Lithium Alkoxides. *J. Chem. Res., Synop.* **1993**, 454–455.
- (38) Beumel, O. F., Jr.; Smith, W. N.; Rybalka, B. Preparation of 2- and 4-Picolylolithium. *Synthesis* **1974**, 43–45.
- (39) Chasalow, F. I. Mechanism and Control of Rat Testicular Steroid Synthesis. *J. Biol. Chem.* **1979**, *254*, 3000–3005.
- (40) Chasalow, F. I.; Marr, H.; Taylor, G. A New Assay and A Solubilisation Procedure for Steroid 17,20-lyase From Rat Testes. *Steroids* **1982**, *30*, 617–630.
- (41) Chou, T. C. Derivation and Properties of Michaelis-Menten Type and Hill Type Equations for Reference Ligands. *J. Theor. Biol.* **1976**, *39*, 253–276.
- (42) Goldstein, A. The Mechanism of Enzyme-Inhibitor-Substrate Reactions. *J. Gen. Physiol.* **1944**, *27*, 529–580.
- (43) Henderson, P. J. F. Steady-state Enzyme Kinetics with High-affinity Substrates or Inhibitors. *Biochem. J.* **1973**, *135*, 101–107.
- (44) Foster, A. B.; Jarman, M.; Leung, C.-S.; Rowlands, M. G.; Taylor, G. N. Analogues of Aminoglutethimide: Selective Inhibition of Cholesterol Side-Chain Cleavage. *J. Med. Chem.* **1983**, *26*, 50–54.
- (45) Jarman, M.; Barrie, S. E.; Houghton, J.; Rowlands, M. G.; Mann, J.; Haase-Held, M.; Hatzis, M. Evaluation of Some 4-Fluoro- and 4-Cyano Derivatives of $\Delta^4,3$ -Ketosteroids as Inhibitors of Testosterone 5 α -Reductase. *J. Enzyme Inhib.* **1994**, *8*, 17–23.
- (46) Wakeling, A. E. In *Steroid Hormones, a Practical Approach*; Green, B., Leake, R. E., Eds.; IRL Press Ltd.: Oxford, 1987; pp 219–236.
- (47) Hardcastle, I. R.; Rowlands, M. G.; Houghton, J.; Parr, I. B.; Potter, G. A.; Jarman, M.; Edwards, K. J.; Laughton, C. A.; Neidle, S. Rationally Designed Analogues of Tamoxifen with Improved Calmodulin Antagonism. *J. Med. Chem.* **1995**, *38*, 241–248.

JM950035N