pubs.acs.org/OrgLett

Letter

Preference of *cis*-Thioamide Structure in *N*-Thioacyl-*N*-methylanilines

Jin Zhang,* Zhulin Liu, Zheng Yin, Xiufang Yang, Yangmin Ma, Roman Szostak, and Michal Szostak*



ABSTRACT: The thioamide group represents a highly attractive isostere of the amide bond. We report a combined structural and computational study on *cis*-thioamide conformation of *N*-thioacyl-*N*-methylanilines. Amide to thioamide replacement in a class of anilides that are highly valuable as conformational locks results in a higher preference for *cis* conformation in a unique compacted template intrinsic to the thioamide structure. The study strongly supports the use of *N*-methylthioanilides as *cis*-conformational locks in various facets of chemistry.

In their classic study, Itai and co-workers demonstrated the cis-preference of the amide bond in N-methylbenzanilides (Figure 1A).¹ Although acyclic amides strongly prefer trans conformation around the N–C(O) bond,^{2,3} as exemplified by closely related benzanilides, their N-methylated counterparts undergo a conformational switch arising from avoidance of unfavorable steric interactions^{4,5} while retaining substantial double bond character of the amide bond through $n_N \rightarrow \pi^*_{C=O}$ conjugation.^{6,7} This unexpected finding opened the door for the implementation of *trans* to *cis* conformational switch to control substituent geometry around the amide bond in a variety of fields utilizing amides, including medicinal chemistry, biochemistry, molecular recognition, conformation relays, synthetic switches, and organic synthesis.^{8–11}

Recently, significant attention has been given to thioamides as attractive amide bond isosteres.¹² The oxygen to sulfur replacement is particularly attractive considering (1) the privileged role of sulfur-based functional groups in pharmaceuticals, wherein more than 25% of APIs contain sulfur as the key component,¹³ and (2) the unique electronic properties inherent to thioamides vs amides, including long van der Waals radius of sulfur vs oxygen (1.85 Å vs 1.40 Å), elongated C=X bonds (1.64 Å vs 1.19 Å, HC=XNH₂), and significantly lower polarization of the C=X bond (electronegativity, S: 2.58 vs O: 3.44).¹⁴ A prominent feature of thioamides is their role in increasing the stabilization of proteins owing to strong $n \rightarrow \pi^*$ interactions¹⁵ and a plethora of thioamide-containing natural products.¹⁶ Elegant studies on macrocyclic amide to thioamide replacement have attracted considerable interest over the last years.^{17,18}

As part of our program on amide bonds,^{19,20} herein we report a combined structural and computational study on *cis*thioamide conformation of *N*-thioacyl-*N*-methylanilines. Most importantly, we demonstrate that amide to thioamide replacement results in a higher preference for *cis* conformation in a unique compacted template intrinsic to the thioamide structure. The present study strongly supports the use of *N*methylthioanilides as highly valuable *cis*-conformational locks^{8–11} in various facets of chemistry.

Guided by the study by Itai and our own studies in amide bond chemistry, we selected six thioamides shown in Figure 2 (1-6). These thioamides are readily synthesized from the corresponding amides using Lawesson's reagent (see the Supporting Information, SI). The selected compounds mirror the amide counterparts used by Itai (Figure 2, 7-12).

We commenced by obtaining X-ray structures of Nmethylthioanilides (Figure 3). All thioamides 1-5 feature *cis* conformation in the crystal. Selected structural parameters relevant to the thioamide geometry are presented in Table 1 and the SI. Compound 6 is a liquid and is not suitable for crystallographic analysis (vide infra). There are several instructive structural correlations between thioamides 1-5

Received: October 21, 2020





(synthesis, medicinal chemistry, peptide backbones, natural products)

Figure 1. (a) Classical study on *cis*-preference of anilides, selected applications. (b) This work: conformational preference of thioanilides.



Figure 2. Structures of thioamides used in the present study.

and their amide counterparts 7–11 obtained from crystallographic studies, including (1) a significant decrease of the N– C(X) bond length (avg 1.337 Å, X = S; avg 1.352 Å, X = O); (2) a significant increase of the C=X bond length (avg 1.660 Å, X = S; avg 1.228 Å, X = O); (3) an increase in C–C(Ph) bond length (avg 1.443 Å, X = S; avg 1.439 Å, X = O); and (4) an excellent linear correlation between the C–C(S) and C– C(O) bond lengths (R^2 = 0.92, see the SI for correlation plots).²¹

Next, we performed computational studies to gain insight into the structures of thioamides 1-6 and compare them with their oxygen counterparts 7-12. As shown by our previous studies,¹⁹ computed data provide vastly improved correlations vs solid state structures, and this approach is particularly effective when considering multiple series of compounds using



Figure 3. Crystal structures of 1-5. See SI for expanded structures. 50% ellipsoids. Crystallographic data have been deposited with the Cambridge Crystallographic Data Center.

Table 1. Selected Crystallographic Structural Parameters of Thioamides $1-5^a$

| no. | thioamide (R) | N-С (Å) | C S (Å) | C-C(S) (Å) | N—Ph (Å) | N–Me (Å) |
|---|-----------------------|------------|-----------------------|---------------|-------------|-------------|
| 1 | Ph | 1.342 | 1.658 | 1.485 | 1.440 | 1.463 |
| 2 | $C(CH_3)$ = CH_2 | 1.333 | 1.655 | 1.490 | 1.447 | 1.466 |
| 3 | Су | 1.332 | 1.664 | 1.481 | 1.441 | 1.467 |
| 4 | <i>i</i> -Pr | 1.338 | 1.659 | 1.519 | 1.442 | 1.469 |
| 5 | t-Bu | 1.341 | 1.662 | 1.546 | 1.444 | 1.476 |
| ^{<i>a</i>} This study. For X-ray structures, see the SI for details. | | | | | | |

X-ray structures for geometry optimization. Geometry optimization was performed at the B3LYP/6-311++G(d,p) level. Extensive studies have shown that this level is accurate in predicting structural and energetic properties of amides.¹⁹

Computations closely parallel the experimental properties of the amide to thioamide bond substitution and allow us to include noncrystalline compounds **6** and **12** in the correlation (Figure 4 and SI). Thus, amide to thioamide replacement is accompanied by a reinforced n_N to $\pi^*_{C=X}$ conjugation as evidenced by a shortening of the N–C(X) bond. This effect is accompanied by C = X bond elongation as well as C–C(X) bond shortening and C–C_(Ph) bond elongation, consistent with a weakened $n_N \rightarrow Ar$ conjugation. Overall, these effects lead to a compacted *cis* amide bond geometry in the thioanilide template.

Next, resonance energies of the thioamide bond in 1-6 were calculated using the COSNAR method.²² Resonance energy in 1-6 ranges between 9.6 and 19.9 kcal/mol and is higher than in the corresponding oxygen counterparts 7-12 (9.8–15.4 kcal/mol) (Figure 5A and SI). Remarkably, there is an excellent inverse linear correlation between a plot of log(ΔRE) of the sulfur and oxygen analogues and the steric Charton (ν) value (Figure 5B, $R^2 = 0.91$). Overall, the energetic parameters in 1-6 vs 7-12 indicate a reinforced bond resonance upon O to S replacement. The resonance manifests as a function of the R substitution. This unexpected effect likely arises from the increased van der Waals radius of sulfur vs oxygen atoms minimizing steric interactions.

To gain further insight into the effect of O to S replacement, we obtained a detailed rotational profile of the parent thioamide 1 by systematic rotation along the X-C-N-C dihedral angle (Figure 5C). The rotation was performed in

Organic Letters

pubs.acs.org/OrgLett







Figure 4. (a) Plot of N–C bond length (Å) in thioamides 1–6 vs amides 7–12. (b) Plot of N–Ph bond length (Å) in thioamides 1–6 vs amides 7–12. (c) Correlation of C–C(S) bond length (Å) to C–C(O) bond length (Å)SI for additional details.



Figure 5. (a) Resonance energies for thioamides 1–6 (kcal/mol). (b) Plot of log(ΔRE) vs Charton parameter in thioamides 1–6 vs amides 7–12. Note that the ν value for C(CH₃)=CH₂ is not available. (c) Correlation of ΔE [kcal/mol] to X–C–C–C (deg) in 1 and 7 (B3LYP/6-311++G(d,p)). See the SI for additional details.

both directions using the X-ray structure of 1 as the starting geometry. Figure 5C shows rotational profile of 1 (X = S) in comparison with 7 (X = O). The rotational profile in 1 confirms a significantly higher barrier to rotation than in the oxygen counterpart with a planar thioamide bond conformation (the energy minimum at ca. 0° S-C-N-C angle; the energy maximum at ca. 90° S-C-N-C angle). The rotational barrier was determined to be 14.02 kcal/mol (90° S-C-N-C angle) and 10.84 kcal/mol (70° O-C-N-C angle). It is worth noting that as suggested by the resonance energies thioamide 6 shows one conformer in the NMR spectrum. This effect is driven by minimization of steric interactions between the R substituent and the sulfur atom. Both steric and electronic effects of the substituents play a key role in adopting the *cis* conformation.

Several additional studies were conducted (Scheme 1 and see SI). (1) To further confirm the intrinsic preference of *N*methylthioanilides to exist in the *cis* conformation, the relative stabilities of the *trans* isomers for both sulfur and oxygen analogues were determined at the B3LYP/6-311++G(d,p) level, showing that *cis* isomers are more stable in all cases (S: avg 3.1 kcal/mol; O: avg 2.7 kcal/mol, see SI). (2) The difference in proton affinities (Δ PA) in representative thioamide 1 indicates that these thioanilides strongly favor protonation at sulfur¹⁹ (Δ PA = 17.0 kcal/mol, S_{PA} vs N_{PA}), which can be compared with the oxygen analogue 7 (Δ PA = 11.0 kcal/mol, O_{PA} vs N_{PA})^{19e} and is in agreement with the findings on the enhanced n_N $\rightarrow \pi^*_{C=X}$ conjugation in thioamides (Scheme 1A). (3) To push the prospects of resonance alteration, RE of derivatives 13–16 varying by a



single substituent on each aromatic ring were determined (Scheme 1B) (13: R = 4-NMe₂/R' = 4-NO₂; 14: R = 4-MeO/ R' = 4-CN; 15: R = 4-CN/R' = 4-MeO; 16: R = 4-NO₂/R' = 4-NMe₂). The RE in 13–16 of 8.7, 11.1 19.6, 20.6 kcal/mol spans the range of 11.9 kcal/mol, which supersedes the effects observed in the oxygen counterparts of 9.5 kcal/mol determined earlier.^{19e} Overall, these studies strongly support significantly enhanced conformational buttressing effects upon O to S substitution.

In summary, amide bond architecture represents one of the most vital motifs in organic chemistry and biology. Although typical acyclic amides exist in the *trans* conformation, *N*-methylanilides undergo a conformation switch to the *cis* geometry, an effect that has been engaged as a conformational anchor in many systems across chemical disciplines.^{8–11} The present study demonstrates and quantifies that the amide to

thioamide replacement in these systems results in a higher preference for *cis* conformation in a unique compacted template intrinsic to the thioamide structure. The effect of sterics, substitution, resonance effects, barriers to rotation, and proton affinities have been discussed and quantified. The attractive properties of thioamides, including photoisomerization, should facilitate the widespread use of *N*-methylthioanilides as *cis*-conformational locks in various facets of chemistry. Future studies will focus on further determination of conformational preferences of amides and thioamides by spectroscopic and computational methods.

ASSOCIATED CONTENT

3 Supporting Information

The Supporting Information is available free of charge at https://pubs.acs.org/doi/10.1021/acs.orglett.0c03512.

Experimental procedures and characterization data. Cartesian coordinates and energies. Detailed description of computational methods used. CIF files for 1-5 (PDF)

Accession Codes

CCDC 2039785, 2039803, and 2039815–2039817 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/ data_request/cif, or by emailing data_request@ccdc.cam.ac. uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Authors

Jin Zhang – College of Chemistry and Chemical Engineering, Key Laboratory of Chemical Additives for China National Light Industry, Shaanxi University of Science and Technology, Xi'an 710021, China; orcid.org/0000-0002-6616-7087; Email: zhangjin@sust.edu.cn

Michal Szostak – Department of Chemistry, Rutgers University, Newark, New Jersey 07102, United States; orcid.org/0000-0002-9650-9690; Email: michal.szostak@rutgers.edu

Authors

- Zhulin Liu College of Chemistry and Chemical Engineering, Key Laboratory of Chemical Additives for China National Light Industry, Shaanxi University of Science and Technology, Xi'an 710021, China
- Zheng Yin College of Chemistry and Chemical Engineering, Key Laboratory of Chemical Additives for China National Light Industry, Shaanxi University of Science and Technology, Xi'an 710021, China
- Xiufang Yang College of Chemistry and Chemical Engineering, Key Laboratory of Chemical Additives for China National Light Industry, Shaanxi University of Science and Technology, Xi'an 710021, China
- Yangmin Ma College of Chemistry and Chemical Engineering, Key Laboratory of Chemical Additives for China National Light Industry, Shaanxi University of Science and Technology, Xi'an 710021, China
- Roman Szostak Department of Chemistry, Wroclaw University, Wroclaw 50-383, Poland

Complete contact information is available at:

https://pubs.acs.org/10.1021/acs.orglett.0c03512

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

J.Z. thanks the CSC (No. 201808610096). Support was provided by the Foundation for Young Scholars of SUST (No. BJ12-26). M.S. thanks Rutgers University and the NSF (CHE-1650766). We thank the Wroclaw Center for Networking and Supercomputing (Grant No. WCSS159, R.S.).

REFERENCES

(1) Itai, A.; Toriumi, Y.; Saito, S.; Kagechika, H.; Shudo, K. Preference of *cis*-Amide Structure in *N*-Acyl-*N*-methylanilines. *J. Am. Chem. Soc.* **1992**, *114*, 10649–10650.

(2) Pauling, L. The Nature of the Chemical Bond; Oxford University Press: London, 1940.

(3) Greenberg, A.; Breneman, C. M.; Liebman, J. F., Eds. The Amide Linkage: Structural Significance in Chemistry, Biochemistry, and Materials Science; Wiley: New York, 2000.

(4) For further studies, see: (a) Zimmerman, S. S.; Scheraga, H. A. Stability of Cis, Trans, and Nonplanar Peptide Groups. Macromolecules 1976, 9, 408-416. (b) Zimmerman, S. S.; Scheraga, H. A. Influence of local interactions on protein structure. I. Conformational energy studies of N-acetyl-N'-methylamides of pro-X and X-pro dipeptides. Biopolymers 1977, 16, 811-843. (c) Wedemeyer, W. J.; Welker, E.; Scheraga, H. A. Proline Cis-Trans Isomerization and Protein Folding. Biochemistry 2002, 41, 14637-14644. (d) Dugave, C.; Demange, L. Cis-Trans Isomerization of Organic Molecules and Biomolecules: Implications and Applications. Chem. Rev. 2003, 103, 2475–2532. (e) Eisenberg, D. The discovery of the α -helix and β sheet, the principal structural features of proteins. Proc. Natl. Acad. Sci. U. S. A. 2003, 100, 11207-11210. (f) Forbes, C.; Beatty, A. M.; Smith, B. D. Using Pentafluorophenyl as a Lewis Acid To Stabilize a Cis Secondary Amide Conformation. Org. Lett. 2001, 3, 3595-3598. (g) Laursen, J. S.; Engel-Andreasen, J.; Fristrup, P.; Harris, P.; Olsen, C. A. Cis-Trans Amide Bond Rotamers in β -Peptoids and Peptoids: Evaluation of Stereoelectronic Effects in Backbone and Side Chains. J. Am. Chem. Soc. 2013, 135, 2835-2844. (h) Nguyen, K.; Iskandar, M.; Rabenstein, D. L. Kinetics and Equilibria of Cis/Trans Isomerization of Secondary Amide Peptide Bonds in Linear and Cyclic Peptides. J. Phys. Chem. B 2010, 114, 3387-3392.

(5) For recent elegant studies on enforcing amide bond conformation by steric effects, see: (a) Tani, K.; Stoltz, B. M. Synthesis and structural analysis of 2-quinuclidonium tetrafluoroborate. *Nature* **2006**, *441*, 731–734. (b) Komarov, I. V.; Yanik, S.; Ishchenko, A. Y.; Davies, J. E.; Goodman, J. M.; Kirby, A. J. The Most Reactive Amide As a Transition-State Mimic For *cis*-trans Interconversion. *J. Am. Chem. Soc.* **2015**, *137*, 926–930. (c) Adachi, S.; Kumagai, N.; Shibasaki, M. Pyramidalization/twisting of the amide functional group via remote steric congestion triggered by metal coordination. *Chem. Sci.* **2017**, *8*, 85–90. (d) Otani, Y.; Liu, X.; Ohno, H.; Wang, S.; Zhai, L.; Su, A.; Kawahata, M.; Yamaguchi, K.; Ohwada, T. Amide Nitrogen Pyramidalization Changes Lactam Amide Spinning. *Nat. Commun.* **2019**, *10*, 461. (e) Takezawa, H.; Shitozawa, T.; Fujita, M. Enhanced Reactivity of Twisted Amides Inside a Molecular Cage. *Nat. Chem.* **2020**, *12*, 574–578.

(6) For selected theoretical studies on amide bonds, see: (a) Kemnitz, C. R.; Loewen, M. J. Amide Resonance" Correlates with a Breadth of C-N Rotation Barriers. J. Am. Chem. Soc. 2007, 129, 2521-2528. (b) Mujika, J. I.; Mercero, J. M.; Lopez, X. Water-Promoted Hydrolysis of a Highly Twisted Amide: Rate Acceleration Caused by the Twist of the Amide Bond. J. Am. Chem. Soc. 2005, 127, 4445-4453. (c) Glover, S. A.; Rosser, A. A. Reliable Determination of Amidicity in Acyclic Amides and Lactams. J. Org. Chem. 2012, 77, 5492-5502. (d) Morgan, J.; Greenberg, A.; Liebman, J. F. Paradigms and Paradoxes: O- and N-Protonated Amides, Stabilization Energy and Resonance Energy. *Struct. Chem.* **2012**, *23*, 197–199. (e) Morgan, J. P.; Weaver-Guevara, H. M.; Fitzgerald, R. W.; Dunlap-Smith, A.; Greenberg, A. Ab initio computational study of 1-methyl-4silatranone and attempts at its conventional synthesis. *Struct. Chem.* **2017**, *28*, 327–331.

(7) For selected studies on proteolysis of activated amides, see: (a) Elashal, H. E.; Raj, M. Site-Selective Chemical Cleavage of Peptide Bonds. *Chem. Commun.* **2016**, *52*, 6304–6307. (b) Elashal, H. E.; Cohen, R. D.; Elashal, H. E.; Raj, M. Oxazolidinone-Mediated Sequence Determination of One-Bead One-Compound Cyclic Peptide Libraries. *Org. Lett.* **2018**, *20*, 2374–2377. (c) Mahesh, S.; Tang, K. C.; Raj, M. Amide Bond Activation of Biological Molecules. *Molecules* **2018**, *23*, 2615.

(8) (a) Nowick, J. S. Chemical Models of Protein β -Sheets. Acc. Chem. Res. **1999**, 32, 287–296. (b) Kilburn, J. P.; Kehler, J.; Langgard, M.; Erichsen, M. N.; Leth-Petersen, S.; Larsen, M.; Christoffersen, C. T.; Nielsen, J. N-Methylanilide and N-methylbenzamide derivatives as phosphodiesterase 10A (PDE10A) inhibitors. *Bioorg. Med. Chem.* **2013**, 21, 6053–6062.

(9) (a) Curran, D. P.; Liu, W.; Chen, C. H. T. Transfer of Chirality in Radical Cyclizations. Cyclization of o-Haloacrylanilides to Oxindoles with Transfer of Axial Chirality to a Newly Formed Stereocenter. J. Am. Chem. Soc. **1999**, *121*, 11012–11013. (b) Curran, D. P.; Hale, G. R.; Geib, S. J.; Balog, A.; Cass, Q. B.; Degani, A. L. G.; Hernandes, M. Z.; Freitas, L. C. G. Rotational features of carbonnitrogen bonds in axially chiral o-tert-butyl anilides and related molecules. Potential substrates for the 'prochiral auxiliary' approach to asymmetric synthesis. *Tetrahedron: Asymmetry* **1997**, *8*, 3955–3975. (c) Guthrie, D. B.; Geib, S. J.; Curran, D. P. Radical and Heck Cyclization of Diastereomeric o-Haloanilide Atropisomers. J. Am. Chem. Soc. **2011**, *133*, 115–122.

(10) (a) Yamaguchi, K.; Matsumura, G.; Kagechika, H.; Azumaya, I.; Ito, Y.; Itai, A.; Shudo, K. Aromatic architecture. Use of the Nmethylamide structure as a molecular splint. J. Am. Chem. Soc. 1991, 113, 5474-5475. (b) Azumaya, I.; Yamaguchi, K.; Okamoto, I.; Kagechika, H.; Shudo, K. Total asymmetric transformation of an Nmethylbenzamide. J. Am. Chem. Soc. 1995, 117, 9083-9084. (c) Tanatani, A.; Yamaguchi, K.; Azumaya, I.; Fukutomi, R.; Shudo, K.; Kagechika, H. N-Methylated Diphenylguanidines: Conformations, Propeller-Type Molecular Chirality, and Construction of Water-Soluble Oligomers with Multilayered Aromatic Structures. J. Am. Chem. Soc. 1998, 120, 6433-6442. (d) Yamasaki, R.; Tanatani, A.; Azumaya, I.; Saito, S.; Yamaguchi, K.; Kagechika, H. Amide Conformational Switching Induced by Protonation of Aromatic Substituent. Org. Lett. 2003, 5, 1265-1267. (e) Okamoto, I.; Nabeta, M.; Hayakawa, Y.; Morita, N.; Takeya, T.; Masu, H.; Azumaya, I.; Tamura, O. Acid-Induced Molecular Folding and Unfolding of N-Methyl Aromatic Amide Bearing 2,6-Disubstituted Pyridines. J. Am. Chem. Soc. 2007, 129, 1892-1893. (f) Katoono, R.; Tanaka, Y.; Fujiwara, K.; Suzuki, T. A Foldable Cyclic Oligomer: Chiroptical Modulation through Molecular Folding upon Complexation and a Change in Temperature. J. Org. Chem. 2014, 79, 10218-10225.

(11) (a) Shah, N. H.; Butterfoss, G. L.; Nguyen, K.; Yoo, B.; Bonneau, R.; Rabenstein, D. L.; Kirshenbaum, K. Oligo(N-aryl glycines): A New Twist on Structured Peptoids. *J. Am. Chem. Soc.* **2008**, 130, 16622–16632. (b) Knipe, P. C.; Thompson, S.; Hamilton, A. D. Ion-mediated conformational switches. *Chem. Sci.* **2015**, *6*, 1630–1639.

(12) (a) Mahanta, N.; Szantai-Kis, D. M.; Petersson, E. J.; Mitchell, D. A. Biosynthesis and Chemical Applications of Thioamides. ACS Chem. Biol. 2019, 14, 142–163. (b) Murai, T. The Chemistry of Thioamides; Springer: Singapore, 2019.

(13) (a) Ilardi, E. A.; Vitaku, E.; Njardarson, J. T. Data-Mining for Sulfur and Fluorine: An Evaluation of Pharmaceuticals To Reveal Opportunities for Drug Design and Discovery. *J. Med. Chem.* **2014**, *57*, 2832–2842. (b) Beno, B. R.; Yeung, K. S.; Bartberger, M. D.; Pennington, L. D.; Meanwell, N. A. A Survey of the Role of Noncovalent Sulfur Interactions in Drug Design. *J. Med. Chem.* **2015**, Letter

58, 4383–4438. (c) Scott, K. A.; Njardarson, J. T. Analysis of US FDA-Approved Drugs Containing Sulfur Atoms. *Top Curr. Chem.* 2019, 376, 1–34.

(14) Choudhary, A.; Raines, R. T. An Evaluation of Peptide-Bond Isosteres. *ChemBioChem* **2011**, *12*, 1801–1807.

(15) Newberry, R. W.; Van Veller, B.; Guzei, I. A.; Raines, R. T. $n \rightarrow \pi^*$ interactions of amides and thioamides: implications for protein stability. *J. Am. Chem. Soc.* **2013**, *135*, 7843–7846.

(16) (a) Kjaerulff, L.; Sikandar, A.; Zaburannyi, N.; Adam, S.; Herrmann, J.; Koehnke, J.; Müller, R. Thioholgamides: Thioamide-Containing Cytotoxic RiPP Natural Products. *ACS Chem. Biol.* **2017**, *12*, 2837–2841. (b) See also ref 12a.

(17) (a) Khatri, B.; Majumder, P.; Nagesh, J.; Penmatsa, A.; Chatterjee, J. Increasing protein stability by engineering the $n \rightarrow \pi^*$ interaction at the β -turn. *Chem. Sci.* **2020**, *11*, 9480–9487. (b) Verma, H.; Khatri, B.; Chakraborti, S.; Chatterjee, J. Increasing the bioactive space of peptide macrocycles by thioamide substitution. *Chem. Sci.* **2018**, *9*, 2443–2451. (c) Mukherjee, S.; Verma, H.; Chatterjee, J. Efficient Site-Specific Incorporation of Thioamides into Peptides on a Solid Support. Org. Lett. **2015**, *17*, 3150–3153.

(18) For selected studies on thioamides, see: (a) Saito, M.; Murakami, S.; Nanjo, T.; Kobayashi, Y.; Takemoto, Y. Mild and Chemoselective Thioacylation of Amines Enabled by the Nucleophilic Activation of Elemental Sulfur. J. Am. Chem. Soc. 2020, 142, 8130-8135. (b) Fukagawa, S.; Kato, Y.; Tanaka, R.; Kojima, M.; Yoshino, T.; Matsunaga, S. Enantioselective C(sp³)-H Amidation of Thioamides Catalyzed by a CobaltIII/Chiral Carboxylic Acid Hybrid System. Angew. Chem., Int. Ed. 2019, 58, 1153-1157. (c) Jagodziński, T. S. Thioamides as Useful Synthons in the Synthesis of Heterocycles. Chem. Rev. 2003, 103, 197-228. (d) Reiner, A.; Wildemann, D.; Fischer, G.; Kiefhaber, T. Effect of Thioxopeptide Bonds on α -Helix Structure and Stability. J. Am. Chem. Soc. 2008, 130, 8079-8084. (e) Barrett, T. M.; Chen, X. S.; Liu, C.; Giannakoulias, S.; Phan, H. A. T.; Wang, J.; Keenan, E. K.; Karpowicz, R. J., Jr; Petersson, E. J. Studies of Thioamide Effects on Serine Protease Activity Enable Two-Site Stabilization of Cancer Imaging Peptides. ACS Chem. Biol. 2020, 15, 774-779. (f) Walters, C. R.; Szantai-Kis, D. M.; Zhang, Y.; Reinert, Z. E.; Horne, W. S.; Chenoweth, D. M.; Petersson, E. J. The Effects of Thioamide Backbone Substitution on Protein Stability: A Study in Alpha-Helical, Beta-Sheet, and Polyproline Ii Helical Contexts. Chem. Sci. 2017, 8, 2868-2877. (g) Goldberg, J. M.; Batjargal, S.; Petersson, E. J. Thioamides as Fluorescence Quenching Probes: Minimalist Chromophores to Monitor Protein Dynamics. J. Am. Chem. Soc. 2010, 132, 14718-14720. (h) Zhang, J.; Liu, Y.; Jia, Q.; Wang, Y.; Ma, Y.; Szostak, M. Ruthenium(II)-Catalyzed C-H Arylation of N,N-Dialkyl Thiobenzamides with Boronic Acids by Sulfur Coordination in 2-MeTHF. Org. Lett. 2020, 22, 6884-6890. (19) (a) Meng, G.; Shi, S.; Lalancette, R.; Szostak, R.; Szostak, M. Reversible Twisting of Primary Amides via Ground State N-C(O)Destabilization: Highly Twisted Rotationally Inverted Acyclic Amides. J. Am. Chem. Soc. 2018, 140, 727-734. (b) Szostak, R.; Szostak, M. N-Acyl-Glutarimides: Resonance and Proton Affinities of Rotationally-Inverted Twisted Amides Relevant to N-C(O) Cross-Coupling. Org. Lett. 2018, 20, 1342-1345. (c) Liu, C.; Li, G.; Shi, S.; Meng, G.; Lalancette, R.; Szostak, R.; Szostak, M. Acyl and Decarbonylative Suzuki Coupling of N-Acetyl Amides: Electronic Tuning of Twisted, Acyclic Amides in Catalytic Carbon-Nitrogen Bond Cleavage. ACS Catal. 2018, 8, 9131-9139. (d) Liu, C.; Shi, S.; Liu, Y.; Liu, R.; Lalancette, R.; Szostak, R.; Szostak, M. The Most Twisted Acyclic Amides: Structures and Reactivity. Org. Lett. 2018, 20, 7771-7774. (e) Szostak, R.; Meng, G.; Szostak, M. Resonance Destabilization in N-Acylanilines (Anilides): Electronically-Activated Planar Amides of Relevance in N-C(O) Cross-Coupling. J. Org. Chem. 2017, 82, 6373-6378. (f) Li, G.; Ma, S.; Szostak, M. Amide Bond Activation: The Power of Resonance. Trends Chem. 2020, 2, 914-928.

(20) For selected synthetic studies, see: (a) Liu, C.; Szostak, M. Twisted Amides: From Obscurity to Broadly Useful Transition-Metal-Catalyzed Reactions by N–C Amide Bond Activation. *Chem. - Eur. J.* **2017**, *23*, 7157–7173. (b) Meng, G.; Szostak, M. General Olefin

Synthesis by the Palladium-Catalyzed Heck Reaction of Amides: Sterically Controlled Chemoselective N-C Activation. Angew. Chem., Int. Ed. 2015, 54, 14518-14522. (c) Shi, S.; Meng, G.; Szostak, M. Synthesis of Biaryls through Nickel-Catalyzed Suzuki-Miyaura Coupling of Amides by Carbon-Nitrogen Bond Cleavage. Angew. Chem., Int. Ed. 2016, 55, 6959-6963. (d) Liu, C.; Szostak, M. Decarbonylative Phosphorylation of Amides by Palladium and Nickel Catalysis: The Hirao Cross-Coupling of Amide Derivatives. Angew. Chem., Int. Ed. 2017, 56, 12718-12722. (e) Zhang, J.; Hou, Y.; Ma, Y.; Szostak, M. Synthesis of Amides by Mild Palladium-Catalyzed Aminocarbonylation of Arylsilanes with Amines Enabled by Copper-(II) Fluoride. J. Org. Chem. 2019, 84, 338-345. (f) Zhang, J.; Ma, Y.; Ma, Y. Synthesis of Secondary Amides through the Palladium(II)-Catalyzed Aminocarbonylation of Arylboronic Acids with Amines or Hydrazines and Carbon Monoxide. Eur. J. Org. Chem. 2018, 2018, 1720-1725.

(21) (a) Greenberg, A.; Venanzi, C. A. Structures and energetics of two bridgehead lactams and their N- and O-protonated forms: an ab initio molecular orbital study. *J. Am. Chem. Soc.* **1993**, *115*, 6951–6957. (b) Greenberg, A.; Moore, D. T.; DuBois, T. D. Small and Medium-Sized Bridgehead Bicyclic Lactams: A Systematic ab Initio Molecular Orbital Study. *J. Am. Chem. Soc.* **1996**, *118*, 8658–8668.

(22) (a) Wiberg, K. B. The Interaction of Carbonyl Groups with Substituents. Acc. Chem. Res. **1999**, 32, 922–929. (b) Wiberg, K. B.; Rush, D. J. Solvent Effects on the Thioamide Rotational Barrier: An Experimental and Theoretical Study. J. Am. Chem. Soc. **2001**, 123, 2038–2046.