

## Research Article

# Syntheses of perdeuterated indoles and derivatives as probes for the biosyntheses of crucifer phytoalexins

M. Soledade C. Pedras\* and Denis P. O. Okinyo

*Department of Chemistry, University of Saskatchewan, 110 Science Place, Saskatoon, SK S7N 5C9, Canada*

## Summary

A simple two-step preparation of [ $^2\text{H}_4$ ]indole, a starting material necessary for the synthesis of various crucifer metabolites, starting with readily available  $^1\text{H}$  NMR solvent [ $^2\text{H}_5$ ]nitrobenzene (99% deuterated) was developed. [4,5,6,7- $^2\text{H}_4$ ]Indole 99% deuterated at the specified positions was then used to synthesize [4',5',6',7'- $^2\text{H}_4$ ]indolyl-3-acetaldoxime, [4',5',6',7'- $^2\text{H}_4$ ]1-methoxyindolyl-3-acetaldoxime, [1'',1'',1'',4',5',6',7'- $^2\text{H}_7$ ]1-methoxyindolyl-3-acetaldoxime, [4',5',6',7'- $^2\text{H}_4$ ]1-methoxybrassinin, and [3,3,3,4',5',6',7'- $^2\text{H}_7$ ]1-methoxybrassinin. Copyright © 2005 John Wiley & Sons, Ltd.

**Key Words:** 1-methoxybrassinin; 1-methoxyindolyl-3-acetaldoxime; brassinin; indolyl-3-acetaldoxime; tetradeuterated indole

## Introduction

The interesting array of phytoalexins, i.e. chemical defenses, produced by plants of the family Cruciferae (Brassicaceae) has prompted diverse studies dealing with their syntheses, biosynthesis, biotransformation, and biological activity.<sup>1</sup> In particular, the correlation between phytoalexins and defense pathways operating in plants<sup>2</sup> explains both the interest in and significance of the biosynthetic pathway of crucifer phytoalexins. Knowledge of such a pathway(s) is of great importance in the genetic manipulation of secondary metabolites of crucifers. Crucifer crops are the third largest source of vegetable oils (canola, rapeseed, and mustard oils) and include as well a great variety of vegetables used worldwide as staple food (broccoli, cauliflower, cabbage,

\*Correspondence to: M. Soledade C. Pedras, Department of Chemistry, University of Saskatchewan, 110 Science Place, Sask., Canada SK S7N 5C9. E-mail: s.pedras@usask.ca

Contract/grant sponsor: Natural Sciences and Engineering Research Council of Canada


Contract/grant sponsor: University of Saskatchewan



## Results and discussion

The use of perdeuterated compounds in biosynthetic studies is invaluable as extremely small amounts of deuterated metabolites can be detected unambiguously by mass spectrometry, e.g. ESI-HRMS or APCI-HRMS.<sup>9</sup> The indole derivatives of immediate interest in biosynthetic studies of phytoalexins need to contain at least three deuterium atoms, as this number will allow the unambiguous detection and assignment of labeled phytoalexins using HRMS analysis of  $m/z$   $[M+3]^{+/-}$  and corresponding fragment ions  $m/z$   $[M-X+3]^{+/-}$ . The target compounds were selected containing different deuterium composition:  $[4',5',6',7'-^2\text{H}_4]$ indolyl-3-acetaldoxime (**2a**),  $[4',5',6',7'-^2\text{H}_4]$ 1-methoxyindolyl-3-acetaldoxime (**3a**),  $[1'',1'',1'',4',5',6',7'-^2\text{H}_7]$ 1-methoxyindolyl-3-acetaldoxime (**3b**),  $[4',5',6',7'-^2\text{H}_4]$ 1-methoxybrassinin (**5a**), and  $[3,3,3,4',5',6',7'-^2\text{H}_7]$ 1-methoxybrassinin (**5b**). The required starting material for these syntheses,  $[4,5,6,7-^2\text{H}_4]$ indole (**17a**), needed to be at least 99% deuterated; however, commercially available perdeuterated indoles either did not offer this percentage of deuterium content at specific sites or were extremely expensive. Subsequently, after consideration of a previous synthesis of 99% tetradeuterated indole,<sup>10</sup> which included the preparation of  $[^2\text{H}_4]$ 2-nitrophenylacetonitrile from  $[^2\text{H}_8]$ toluene,<sup>11</sup> a rather difficult and somewhat hazardous preparation, we searched for a shorter route to  $[^2\text{H}_4]$ 2-nitrophenylacetonitrile that could potentially provide access to  $[2,3-^{13}\text{C}_2]$ indole as well. The well-known vicarious nucleophilic substitution of stabilized *R*-chlorocarbanions with nitrobenzene<sup>12</sup> appeared to provide a ready entry to various isotopically labeled indoles. Thus,  $[^2\text{H}_4]$ 2-nitrophenylacetonitrile was obtained from  $[^2\text{H}_5]$ nitrobenzene and provided access to  $[4,5,6,7-^2\text{H}_4]$ indole (**17a**), from which the readily accessible key intermediate  $[^2\text{H}_4]$ indolyl-3-acetonitrile (**11a**) could be prepared.<sup>13,14</sup> The required deuterated oxime **2a** could be readily obtained from **11a**,<sup>15</sup> but the synthesis of perdeuterated *N*<sub>6</sub>-acetyl-1-methoxytryptamine (**19a**, **19b**) from which the corresponding oximes could be obtained through oxidation, required a higher number of steps.  $[^2\text{H}_4]$ Indole (**17a**) was the starting material also used to prepare deuterated methoxyindole (**21a**) and methoxybrassinins **5a** and **5b** via  $[^2\text{H}_4]$ indole-3-carboxaldehyde oxime (**22a**), using a well-established route for non-labeled materials.<sup>1</sup>


Subsequently,  $[4,5,6,7-^2\text{H}_4]$ indole (**17a**) was obtained in two steps and 24% overall yield, starting with the readily available  $^1\text{H}$  NMR solvent  $[2,3,4,5,6-^2\text{H}_5]$ nitrobenzene (**16a**, 99%,  $^2\text{H}_5$ ) and 2-chloroacetonitrile,<sup>11</sup> as summarized in Scheme 2 and described in the Experimental section.  $[4',5',6',7'-^2\text{H}_4]$ Indolyl-3-acetonitrile (**11a**) was prepared by reaction of indolyl-magnesium iodide (prepared from  $[^2\text{H}_4]$ indole (**17a**), and Mg/methyl iodide) with bromoacetonitrile.<sup>13,14</sup> Reduction of indolyl-3-acetonitrile (**11a**) to indolyl-3-acetaldehyde employing DIBAL-H, followed by hydrolysis<sup>15</sup> and treatment with  $\text{HONH}_2 \cdot \text{HCl}$  and NaOAc yielded the desired



Scheme 2. Synthesis of perdeuterated indolyl-3-acetaldoximes **2a**, **3a** and **3b**

[4',5',6',7'- $^2\text{H}_4$ ]indolyl-3-acetaldoxime (**2a**) in acceptable yields. [ $^2\text{H}_7$ ] $N_b$ -acetyl 1-methoxytryptamine (**19b**) was prepared from **18a** after reduction with  $\text{NaBH}_3\text{CN}$ , followed by oxidation with  $\text{Na}_2\text{WO}_4/\text{H}_2\text{O}_2$ ,<sup>16</sup> and methylation of the 1-hydroxyindolyl intermediate with  $(\text{CD}_3\text{O})_2\text{SO}_2$ . [ $^2\text{H}_4$ ] $N_b$ -acetyl 1-methoxytryptamine (**19a**) was prepared similar to the [ $^2\text{H}_7$ ] compound **19b** but using  $(\text{CH}_3\text{O})_2\text{SO}_2$ ,<sup>16</sup> instead of the hexadeuterated methylating reagent. Perdeuterated 1-methoxyindolyl-3-acetaldoximes **3a** and **3b** were then obtained by hydrolysis of the  $N_b$ -acetyl group followed by oxidation of the amine to oxime using  $\text{Na}_2\text{WO}_4/\text{H}_2\text{O}_2$ .<sup>17</sup> This oxidation step proved rather difficult, mostly due to decomposition of the product under the reaction conditions used, however no simple and efficient methods were found to oxidize aliphatic amines to oximes. For example,  $\text{CH}_3\text{ReO}_3/\text{H}_2\text{O}_2$  catalyzed oxidation of primary alkylamines possessing the  $\alpha\text{-C-H}$  bond was found to yield mixtures of oximes, nitroso dimers, and azoxy compounds.<sup>18</sup>

Next, perdeuterated 1-methoxybrassinins **5a** and **5b** were obtained from [4,5,6,7- $^2\text{H}_4$ ]indole as shown in Scheme 3. First, reduction of [ $^2\text{H}_4$ ]indole (**17a**) with  $\text{NaBH}_3\text{CN}$  in AcOH followed by oxidation with  $\text{Na}_2\text{WO}_4/\text{H}_2\text{O}_2$  and



**Scheme 3.** Synthesis of perdeuterated 1-methoxybrassinins **5a** and **5b**

methylation with (CH<sub>3</sub>O)<sub>2</sub>SO<sub>2</sub><sup>16</sup> yielded [<sup>2</sup>H<sub>4</sub>]1-methoxyindole (**21a**) in moderate yield. Standard Vilsmeier-Haack conditions yielded [<sup>2</sup>H<sub>4</sub>]1-methoxyindole-3-carboxaldehyde, which was first treated with HONH<sub>2</sub>·HCl and NaOAc followed by reduction of the resulting oxime with NaBH<sub>3</sub>CN, TiCl<sub>3</sub>, and NH<sub>4</sub>OAc in MeOH to yield [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-methoxyindolyl-3-methanamine (**23a**).<sup>19</sup> The synthesis of [<sup>2</sup>H<sub>4</sub>] and [<sup>2</sup>H<sub>7</sub>]1-methoxybrassinins **5a** and **5b**, followed previously set routes,<sup>1</sup> which afforded the desired compounds in reasonable yields.

All the synthesis described above afforded perdeuterated products 99% labeled at the indicated sites.

## Experimental

### General

All chemicals were purchased from Sigma-Aldrich Canada Ltd., Oakville, Ont., except for deuterated solvents that were purchased from Cambridge Isotopes Laboratories Inc., Andover, MA. All solvents were high-performance liquid chromatograph (HPLC) grade and used as such, except CHCl<sub>3</sub> and CH<sub>2</sub>Cl<sub>2</sub> that were redistilled. Preparative TLC: (Merck, Kieselgel 60 F<sub>254</sub>) 20 × 20 cm × 0.25 mm; analytical TLC (Merck, Kieselgel 60 F<sub>254</sub>, aluminum sheets) 5 × 2.5 cm × 0.2 mm; compounds were visualized by exposure to UV and by dipping the plates in a 5% aqueous (w/v) phosphomolybdic acid solution containing a trace of ceric sulfate and 4% (v/v) H<sub>2</sub>SO<sub>4</sub>, followed by heating at 200°C. Flash column chromatography (FCC): silica gel Merck, grade 60, mesh size 230–400, 60 Å. HPLC analysis was carried out with an

HPLC equipped with quaternary pump, automatic injector and diode array detector (wavelength range 190–600 nm), degasser and a Hypersil ODS column (5  $\mu$ m particle size silica, 4.6 i.d.  $\times$  200 mm) equipped with an in-line filter. HPLC mobile phase H<sub>2</sub>O–CH<sub>3</sub>CN: 75%/25%–0%/100%, for 45 min, linear gradient and flow rate 1.0 ml/min. NMR spectra were recorded on Bruker 500 MHz Avance spectrometers; for <sup>1</sup>H (500 MHz), values were referenced to CHCl<sub>3</sub> (7.27 ppm) or CHD<sub>2</sub>CN (1.94 ppm), for <sup>13</sup>C (125.8 MHz) referenced to CHCl<sub>3</sub> (77.2 ppm) or CD<sub>3</sub>CN (118.7 ppm). Mass spectra (MS) were obtained on a VG 70 SE mass spectrometer (high resolution (HR), electron impact (EI)), employing a solids probe.

*[4',5',6',7'-<sup>2</sup>H<sub>4</sub>]Indolyl-3-acetaldoxime (2a)*

A solution of DIBAL-H (850  $\mu$ l, 1.28 mmol) in toluene was added drop wise to a solution of [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]indolyl-3-acetonitrile (100 mg, 0.63 mmol) in dry toluene (8.0 ml) cooled to –78°C under atmosphere of argon. The reaction mixture was allowed to stir at –78°C for 10 min, was diluted with ice-cold HCl (10.0 ml, 2 M) and immediately extracted in EtOAc (20.0 ml  $\times$  3).<sup>15</sup> The combined organic extract was washed with H<sub>2</sub>O (10.0 ml  $\times$  2), was dried over Na<sub>2</sub>SO<sub>4</sub>, and was concentrated under reduced pressure to yield crude [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]indolyl-3-acetaldehyde, which was used for the next step without purification. A solution of HONH<sub>2</sub>·HCl (111 mg, 1.60 mmol) and CH<sub>3</sub>COONa (131 mg, 1.60 mmol) in water (1.0 ml) was added to a cooled solution (1°C) of crude [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]indolyl-3-acetaldehyde in EtOH (8.5 ml). The reaction mixture was allowed to stir at 1°C for 10 min, and then at r.t. for 15 min. The reaction mixture was concentrated, the residue was dissolved in H<sub>2</sub>O (15.0 ml), extracted with EtOAc (15.0 ml  $\times$  4), the combined organic extract was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure. Separation by FCC (CH<sub>2</sub>Cl<sub>2</sub>–MeOH, 97:3, v/v) yielded [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]indolyl-3-acetaldoxime (**2a**, 36.8 mg, 33% yield from indolyl-3-acetonitrile (**11a**)).

<sup>1</sup>H NMR (500 MHz CD<sub>3</sub>CN): (two isomers, 1.1:0.9) major isomer:  $\delta$  3.60 (d,  $J$  = 6 Hz, H<sub>2</sub>-2), 7.12 (d,  $J$  = 2 Hz, H-2'), 7.49 (t,  $J$  = 6 Hz, H-1), 8.37 (s, OH), 9.17 (br s, NH); minor isomer:  $\delta$  3.78 (d,  $J$  = 5 Hz, H<sub>2</sub>-2), 6.83 (t,  $J$  = 5 Hz, H-1), 7.15 (d,  $J$  = 1 Hz, H-2'), 8.85 (s, OH), 9.17 (br s, NH). HREIMS  $m/z$  measured 178.1044 (178.1044 calculated for C<sub>10</sub>H<sub>6</sub>D<sub>4</sub>N<sub>2</sub>O); EIMS  $m/z$  (relative abundance) 178 [M]<sup>+</sup> (76), 134 (100).

*[4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-Methoxyindolyl-3-acetaldoxime (3a)*

Ten percent Pd/C (80 mg) was added to a solution of [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]indolyl-3-acetonitrile (80 mg, 0.50 mmol) in Ac<sub>2</sub>O (4.0 ml), and the mixture was stirred at r.t. under H<sub>2</sub> atmosphere (balloon pressure). After 14 h the catalyst was filtered

off, the filtrate was diluted with toluene and concentrated under reduced pressure, the residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (15.0 ml), was washed with a 10% solution of  $\text{NaHCO}_3$  (8.0 ml),  $\text{H}_2\text{O}$  (8.0 ml  $\times$  2), and dried over  $\text{Na}_2\text{SO}_4$ . After removal of the solvent under reduced pressure, crude  $[4',5',6',7'\text{-}^2\text{H}_4]\text{N}_b$ -acetyltryptamine was obtained (87.9 mg, 85%) in sufficient purity to use directly in the next step. To a solution of  $[4',5',6',7'\text{-}^2\text{H}_4]\text{N}_b$ -acetyltryptamine (100 mg, 0.495 mmol) in glacial acetic acid (2.0 ml) at r.t.,  $\text{NaBH}_3\text{CN}$  (47 mg, 0.74 mmol) was added in portions through a canula. The reaction mixture was allowed to stir for 3 h at room temperature, was diluted with  $\text{H}_2\text{O}$  (5.0 ml), was basified with  $\text{NaOH}$ , and was extracted with  $\text{Et}_2\text{O}$  (15.0 ml  $\times$  3). The combined organic extract was dried over  $\text{Na}_2\text{SO}_4$  and concentrated under reduced pressure to yield crude  $[4',5',6',7'\text{-}^2\text{H}_4]\text{N}_b$ -acetyl-2,3-dihydrotryptamine (117 mg, 99%), which was used in the next step without purification. A solution of  $\text{Na}_2\text{WO}_4 \cdot 2\text{H}_2\text{O}$  (17.2 mg, 0.0943 mmol) in  $\text{H}_2\text{O}$  (199  $\mu\text{l}$ ) was added to a solution of  $[4',5',6',7'\text{-}^2\text{H}_4]\text{N}_b$ -acetyl-2,3-dihydrotryptamine (112.8 mg, 0.542 mmol) in  $\text{MeOH}$  (2.2 ml) cooled to  $-20^\circ\text{C}$  under stirring, followed by drop wise addition of a solution of  $\text{H}_2\text{O}_2$  (520  $\mu\text{l}$ , 5.42 mmol) in  $\text{MeOH}$  (540  $\mu\text{l}$ ). After being stirred at r.t. for 10 min,  $\text{K}_2\text{CO}_3$  (601 mg, 4.44 mmol) and  $(\text{CH}_3\text{O})_2\text{SO}_2$  (78  $\mu\text{l}$ , 0.87 mmol) were added under vigorous stirring at r.t.<sup>16</sup> After 60 min, the reaction mixture was diluted with  $\text{H}_2\text{O}$  (12.0 ml), was extracted in  $\text{Et}_2\text{O}$  (20.0 ml  $\times$  3), the combined organic extract was dried over  $\text{Na}_2\text{SO}_4$  and was concentrated under reduced pressure to yield crude  $[4',5',6',7'\text{-}^2\text{H}_4]\text{N}_b$ -acetyl-1-methoxytryptamine (**19a**, 79.4 mg, 62%). Crude  $[4',5',6',7'\text{-}^2\text{H}_4]\text{N}_b$ -acetyl-1-methoxytryptamine (79.4 mg) was dissolved in 15% methanolic solution of  $\text{NaOH}$  (20.0 ml) and the mixture was allowed to reflux for 24 h. After removing the solvent under reduced pressure, the residue was dissolved in water, was extracted with  $\text{CH}_2\text{Cl}_2$ – $\text{MeOH}$  (95:5, v/v), was dried over  $\text{Na}_2\text{SO}_4$  and was concentrated under reduced pressure to yield crude  $[4',5',6',7'\text{-}^2\text{H}_4]$ 1-methoxytryptamine (41 mg, 63%). An aqueous solution of  $\text{Na}_2\text{WO}_4 \cdot 2\text{H}_2\text{O}$  (1.4 mg, 0.0042 mmol) in  $\text{H}_2\text{O}$  (60  $\mu\text{l}$ ) was added to the solution of  $[4',5',6',7'\text{-}^2\text{H}_4]$ 1-methoxytryptamine (41 mg, 0.21 mmol) in  $\text{MeOH}$  (400  $\mu\text{l}$ ), the mixture was cooled to  $-15^\circ\text{C}$  and  $\text{H}_2\text{O}_2$  (48  $\mu\text{l}$ , 0.5 mmol) was added under stirring.<sup>17</sup> After being stirred for 60 min at r.t., the mixture was diluted with  $\text{H}_2\text{O}$  (5.0 ml), was basified with  $\text{NaOH}$ , was extracted with  $\text{CH}_2\text{Cl}_2$  (20.0 ml  $\times$  3), the combined organic extract was dried over  $\text{Na}_2\text{SO}_4$  and the solvent was removed under reduced pressure to afford a crude residue (24.3 mg). The crude residue was separated by preparative TLC ( $\text{CH}_2\text{Cl}_2$ – $\text{MeOH}$ , 95:5, v/v) to yield  $[4',5',6',7'\text{-}^2\text{H}_4]$ 1-methoxyindolyl-3-acetaldoxime (**3a**, 4.8 mg, 5% over five steps).

$^1\text{H}$  NMR (500 MHz  $\text{CDCl}_3$ ): (two isomers, 1.1:0.9) major isomer  $\delta$  3.64 (d,  $J$  = 6 Hz,  $\text{H}_2$ -2), 4.08 (s,  $\text{OCH}_3$ ), 7.15 (s,  $\text{H}$ -2'), 7.60 (t,  $J$  = 6 Hz,  $\text{H}$ -1); minor isomer:  $\delta$  3.83 (d,  $J$  = 5 Hz,  $\text{H}_2$ -2), 4.09 (s,  $\text{OCH}_3$ ), 6.93 (t,  $J$  = 5 Hz,

H-1), 7.17 (s, H-2'). HREIMS  $m/z$  (relative abundance) measured 208.1142 (208.1149 calculated for  $C_{11}H_8D_4N_2O_2$ ); EIMS  $m/z$  (relative abundance) 208  $[M]^+$  (100), 190 (25), 164 (45), 159 (54), 132 (70).

*[1'',1'',1'',4',5',7'-<sup>2</sup>H<sub>7</sub>]1-Methoxyindolyl-3-acetaldoxime (3b)*

A solution of  $Na_2WO_4 \cdot 2H_2O$  (32.3 mg, 0.098 mmol) in  $H_2O$  (206  $\mu$ l) was added to a solution of  $[4',5',6',7'-^2H_4]N_b$ -acetyl-2,3-dihydrotryptamine (117 mg, 0.563 mmol) in MeOH (2.3 ml) under stirring.<sup>16</sup> The mixture was cooled to  $-20^\circ C$  and a solution of  $H_2O_2$  (541  $\mu$ l, 5.63 mmol) in MeOH (562  $\mu$ l) was added drop wise. After being stirred for 10 min at r.t.,  $K_2CO_3$  (623 mg, 4.50 mmol) and  $(C^2H_3O)_2SO_2$  (85  $\mu$ l, 0.90 mmol) were added to the reaction mixture under vigorous stirring. After being stirred for 60 min at r.t., the reaction mixture was diluted with  $H_2O$  (12.0 ml), was extracted with  $Et_2O$  (20.0 ml  $\times$  3), the combined organic extract was dried over  $Na_2SO_4$  and was concentrated under reduced pressure to yield crude  $[1'',1'',1'',4',5',6',7'-^2H_7]N_b$ -acetyl-1-methoxytryptamine (84.8 mg, 63%). Crude  $[1'',1'',1'',4',5',6',7'-^2H_7]N_b$ -acetyl-1-methoxytryptamine (84.8 mg) was treated as reported above for  $[4',5',6',7'-^2H_4]N_b$ -acetyl-1-methoxytryptamine (**19a**) to yield  $[1'',1'',1'',4',5',6',7'-^2H_7]$ 1-methoxyindolyl-3-acetaldoxime (**3b**, 4.2 mg, 4% over five steps).

$^1H$  NMR (500 MHz  $CDCl_3$ ): (two isomers, 1.1:0.9) major isomer  $\delta$  3.64 (d,  $J$  = 6 Hz,  $H_2$ -2), 7.15 (s, H-2'), 7.60 (t,  $J$  = 6 Hz, H-1); minor isomer:  $\delta$  3.83 (d,  $J$  = 5 Hz,  $H_2$ -2), 6.93 (t,  $J$  = 5 Hz, H-1), 7.17 (s, H-2'). HREIMS  $m/z$  (relative abundance) measured 211.1335 (211.1338 calculated for  $C_{11}H_5D_7N_2O_2$ ); EIMS  $m/z$  (relative abundance) 211  $[M]^+$  (100), 193 (29), 167 (48), 159 (51), 132 (61).

*[4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-Methoxybrassinin (5a)*

$NaBH_3CN$  (195 mg, 1.68 mmol) was added to a solution of  $[4,5,6,7-^2H_4]$ indole (**17a**, 135 mg, 1.12 mmol) in glacial acetic acid (2.0 ml) under an argon atmosphere. The reaction mixture was stirred at r.t. for 60 min, was diluted with  $H_2O$  (4.0 ml), was basified with NaOH, was extracted with  $Et_2O$  (30.0 ml  $\times$  3), the combined organic extract was dried over  $Na_2SO_4$  and was concentrated under reduced pressure. The crude material was separated by FCC ( $CH_2Cl_2$ ) to yield  $[4,5,6,7-^2H_4]$ indoline (**20a**, 113.2 mg, 82%). A solution of  $Na_2WO_4 \cdot 2H_2O$  (53 mg, 0.16 mmol) in  $H_2O$  (0.35 ml) was added to the stirred solution of  $[4,5,6,7-^2H_4]$ indoline (113.2 mg, 0.920 mmol) in MeOH (3.0 ml) and the suspension was then cooled to  $-20^\circ C$ . A solution of  $H_2O_2$  (766  $\mu$ l, 7.96 mmol) in MeOH (1.0 ml) was added drop wise to the cooled suspension, the reaction mixture was stirred at r.t. for 10 min, after which solid  $K_2CO_3$  (1.02 g, 7.36 mmol) and  $(CH_3O)_2SO_2$  (133  $\mu$ l, 1.47 mmol) were added



under vigorous stirring.<sup>16</sup> After being stirred for 10 min, the reaction mixture was diluted with H<sub>2</sub>O (20.0 ml), was extracted with Et<sub>2</sub>O (30.0 ml × 3), the combined organic extract was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure. The crude material was separated by FCC (hexane–CH<sub>2</sub>Cl<sub>2</sub>, 70:30, v/v) to yield [4,5,6,7-<sup>2</sup>H<sub>4</sub>]1-methoxyindole (**21a**, 73 mg, 52%). Freshly distilled POCl<sub>3</sub> (69 µl, 0.75 mmol) was added to a solution of [4,5,6,7-<sup>2</sup>H<sub>4</sub>]1-methoxyindole (102 mg, 0.68 mmol) in DMF (0.6 ml) and the reaction mixture was stirred at r.t. After 40 min, a solution of aqueous NH<sub>3</sub> (4.0 ml, 28%) was added to the reaction mixture, the reaction mixture was extracted with Et<sub>2</sub>O (10.0 ml × 3), the combined organic extract was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure to yield [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-methoxyindole-3-carboxaldehyde (113 mg, 93%). A solution of HONH<sub>2</sub>·HCl (80.6 mg, 1.16 mmol) and Na<sub>2</sub>CO<sub>3</sub> (61.5 mg, 0.580 mmol) in H<sub>2</sub>O (810 µl) was added to a solution of [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-methoxyindole-3-carboxaldehyde (103 mg, 0.580 mmol) in EtOH (2.7 ml). The resulting reaction mixture was refluxed at 85°C for 60 min, was diluted with H<sub>2</sub>O (3.0 ml) and was extracted with Et<sub>2</sub>O (20.0 ml × 3). The combined organic extract was washed with brine (10.0 ml × 2), was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure to yield [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-methoxyindole-3-carboxaldehyde oxime (**22a**, 122.1 mg, 99%). NaBH<sub>3</sub>CN (163 mg, 2.58 mmol) and NH<sub>4</sub>OAc (219 mg, 2.84 mmol) were added to a solution of [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-methoxyindole-3-carboxaldehyde oxime (50 mg, 0.26 mmol) in MeOH (640 µl) cooled to 0°C and the resulting mixture was treated with TiCl<sub>3</sub> in 2 M HCl (795 µl, 2.04 mmol) neutralized with NaOH.<sup>19</sup> After being stirred for 15 min at 0°C, the reaction mixture was diluted with H<sub>2</sub>O (2.2 ml), was basified with NaOH and was extracted with CH<sub>2</sub>Cl<sub>2</sub> (15.0 ml × 3). The combined organic extract was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure to yield crude [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-methoxyindolyl-3-methanamine (**23a**, 40.3 mg) which was used in the next step without purification. Et<sub>3</sub>N (34 µl, 0.25 mmol) and CS<sub>2</sub> (13 µl, 0.22 mmol) were added to a solution of [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-methoxyindolyl-3-methanamine (40.3 mg, 0.224 mmol) in pyridine (105 µl) cooled to 0°C. After being stirred for 60 min at 0°C, CH<sub>3</sub>I (14 µl, 0.22 mmol) was added to the reaction mixture which was kept at 5°C for 90 min. The reaction mixture was diluted with H<sub>2</sub>O (2.0 ml), was extracted with Et<sub>2</sub>O (10.0 ml × 2), the combined organic extract was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure. The reaction mixture was separated by FCC (hexane–CH<sub>2</sub>Cl<sub>2</sub>) to yield [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-methoxybrassinin (**5a**, 46.5 mg, 67% from [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-methoxyindole-3-carboxaldehyde oxime (**22a**)).

<sup>1</sup>H NMR (500 MHz CDCl<sub>3</sub>): δ 2.67 (s, SCH<sub>3</sub>), 4.12 (s, OCH<sub>3</sub>), 5.04 (d, *J* = 4.5 Hz, H<sub>2</sub>-1), 7.04 (br s, NH), 7.35 (s, H-2'). HREIMS *m/z* (relative abundance) measured 270.0794 (270.0795 calculated for C<sub>12</sub>H<sub>10</sub>D<sub>4</sub>N<sub>2</sub>OS<sub>2</sub>);

EIMS  $m/z$  (relative abundance), 270  $[M]^+$  (6), 239 (64), 222 (15), 191 (8), 164 (100), 149 (15), 133 (43).

*[3,3,3,4',5',6',7'-<sup>2</sup>H<sub>7</sub>]1-Methoxybrassinin (5b)*

NaBH<sub>3</sub>CN (179 mg, 2.84 mmol) and NH<sub>4</sub>OAc (241 mg, 3.13 mmol) were added to a solution of [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-methoxyindole-3-carboxaldehyde oxime (**22a**, 55 mg, 0.284 mmol) in MeOH (700  $\mu$ l) cooled to 0°C. The resulting reaction mixture was treated with a solution of TiCl<sub>3</sub> in 2 M HCl (876  $\mu$ l, 2.25 mmol) neutralized with NaOH.<sup>19</sup> After being stirred for 15 min at 0°C, the reaction mixture was diluted with H<sub>2</sub>O (2.4 ml), was basified with NaOH and was extracted with CH<sub>2</sub>Cl<sub>2</sub> (15.0 ml  $\times$  3). The combined organic extract was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure to yield [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-methoxyindolyl-3-methanamine (**23a**, 44 mg) which was used in the next step without purification. Et<sub>3</sub>N (37  $\mu$ l, 0.27 mmol) and CS<sub>2</sub> (15  $\mu$ l, 0.24 mmol) were added to a solution of [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-methoxyindolyl-3-methanamine (44 mg, 0.24 mmol) in pyridine (118  $\mu$ l) cooled to 0°C. After being stirred for 60 min at 0°C, CD<sub>3</sub>I (16  $\mu$ l, 0.244 mmol) was added and the reaction mixture was kept at 5°C for 90 min. The reaction mixture was diluted in H<sub>2</sub>O (2.0 ml), was extracted with Et<sub>2</sub>O (10.0 ml  $\times$  2), the combined organic extract was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure. The reaction mixture was separated by FCC (hexane–CH<sub>2</sub>Cl<sub>2</sub>) to yield [3,3,3,4',5',6',7'-<sup>2</sup>H<sub>7</sub>]1-methoxybrassinin (**5b**, 52.7 mg, 68% yield from [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]1-methoxyindole-3-carboxaldehyde oxime (**22a**)).

<sup>1</sup>H NMR (500 MHz CDCl<sub>3</sub>):  $\delta$  4.11 (s, (OCH<sub>3</sub>), 5.02 (br s, H<sub>2</sub>-1), 7.04 (br s, NH), 7.32 (br s, H-2'). HREIMS  $m/z$  (relative abundance) measured 273.0984 (273.0987 calculated for C<sub>12</sub>H<sub>7</sub>D<sub>7</sub>N<sub>2</sub>OS<sub>2</sub>); EIMS  $m/z$  (relative abundance) 273  $[M]^+$  (7), 242 (100), 191 (12), 164 (85), 149 (19), 133 (40).

*[4',5',6',7'-<sup>2</sup>H<sub>4</sub>]Indolyl-3-acetonitrile (11a)*

CH<sub>3</sub>I (400  $\mu$ l, 2.80 mmol) was added to Mg (turnings, 80 mg, 3.3 mmol) in dry Et<sub>2</sub>O (5.0 ml) with stirring, under an argon atmosphere. After the Mg was consumed, excess CH<sub>3</sub>I was distilled off and fresh dry Et<sub>2</sub>O (4.0 ml) was added. A solution of [4,5,6,7-<sup>2</sup>H<sub>4</sub>]indole (100 mg, 0.830 mmol) in dry Et<sub>2</sub>O (700  $\mu$ l) was added drop wise and the mixture was allowed to stir for 15 min at r.t., followed by drop wise addition of bromoacetonitrile (160  $\mu$ l, 2.70 mmol).<sup>13,14</sup> After being stirred for 20 min, the reaction mixture was concentrated under reduced pressure and the residue was dissolved in water H<sub>2</sub>O (40.0 ml). The resulting mixture was neutralized with HCl (2.0 ml, 1 M) and was extracted with CH<sub>2</sub>Cl<sub>2</sub> (30.0 ml  $\times$  3). The combined organic extract was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated to dryness under reduced pressure.

Separation by FCC (hexane–CH<sub>2</sub>Cl<sub>2</sub>, 1:1–1:4) yielded [4',5',6',7'-<sup>2</sup>H<sub>4</sub>]indolyl-3-acetonitrile (**11a**, 75.7 mg, 57% yield).

<sup>1</sup>H NMR (500 MHz CDCl<sub>3</sub>): δ 3.86 (s, H<sub>2</sub>-2), 7.24 (t, *J* = 1 Hz, H-2'), 8.23 (br s, NH). HREIMS *m/z* (relative abundance) measured 160.0940 (160.0939 calculated for C<sub>10</sub>H<sub>4</sub>D<sub>4</sub>N<sub>2</sub>); EIMS *m/z* (relative abundance) 160 [M]<sup>+</sup> (62), 159 (100), 134 (32).

#### [4,5,6,7-<sup>2</sup>H<sub>4</sub>]Indole (**17a**)

A solution of [2,3,4,5,6,-<sup>2</sup>H<sub>5</sub>]nitrobenzene (**16a**, 4.20 ml, 39.4 mmol) and chloroacetonitrile (250 μl, 3.94 mmol) in DMSO (3.9 ml) was added drop wise over 30 min to a thoroughly stirred suspension of NaOH (1.60 g, 39.4 mmol) in DMSO (3.9 ml) at 20°C.<sup>12</sup> After 60 min, the reaction mixture was poured into ice-cold HCl (25.0 ml, 1 M), was diluted with H<sub>2</sub>O (10.0 ml), and was extracted with CHCl<sub>3</sub> (50.0 ml × 3). The combined organic extract was washed with H<sub>2</sub>O (70.0 ml), was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure. Separation by FCC (hexane–CH<sub>2</sub>Cl<sub>2</sub>, 100:0–25:75) yielded a mixture of [3,4,5,6,-<sup>2</sup>H<sub>4</sub>]2-nitrophenylacetonitrile and [2,3,5,6,-<sup>2</sup>H<sub>4</sub>]4-nitrophenylacetonitrile (in a 10:1 ratio, 40% yield). Next, 10% Pd on C (220 mg) was added to the solution of [<sup>2</sup>H<sub>4</sub>]2- and [<sup>2</sup>H<sub>4</sub>]4-nitrophenylacetonitriles (200 mg) in EtOAc (20.0 ml) and the resulting reaction mixture was stirred under H<sub>2</sub> atmosphere (balloon pressure) at r.t. After 40 h, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. The residue was taken in CH<sub>2</sub>Cl<sub>2</sub> (15.0 ml), was washed with HCl (10.0 ml × 2, 1 M), was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure to yield [4,5,6,7-<sup>2</sup>H<sub>4</sub>]indole (**17a**, 97.7 mg, 61% yield).

<sup>1</sup>H NMR (500 MHz CDCl<sub>3</sub>): δ 6.61 (br s, H-3), 7.23 (br s, H-2), 8.12 (br s, NH). HREIMS *m/z* (relative abundance) measured 121.0827 (121.0830 calculated for C<sub>8</sub>H<sub>3</sub>D<sub>4</sub>N); EIMS *m/z* (relative abundance) 121 [M]<sup>+</sup> (100), 94 (22), 71 (11), 69 (12), 57 (17).

## Conclusion

To the best of our knowledge, only one synthesis of perdeuterated indole (**17a**) and indole containing phytoalexins with 99% incorporation at multiple sites has been reported to date;<sup>10</sup> however, this new route to indole starting with [<sup>2</sup>H<sub>5</sub>]nitrobenzene (**16a**) is simpler, less hazardous and more efficient, providing an accessible method for the synthesis of perdeuterated brassinins (**5a**, **5b**). Furthermore, a simple route to perdeuterated indolyl-3-acetonitrile (**11a**), a key intermediate for the preparation of perdeuterated indolyl-3-acetaldoxime **2a** and 1-methoxyindolyl-3-acetaldoximes **3a** and **3b** is described for the first time. The availability of 99% labeled indole derivatives **2a**, **3a**, **3b**,

**5a**, **5b** and **11a** will facilitate probing the various biosynthetic relationships among crucifer phytoalexins using LC-HRMS techniques. As well, considering that indolyl-3-acetaldoxime (**2**) is an intermediate in other important metabolic pathways of Brassicaceae, such as those of plant hormones (auxins) and indole glucosinolates,<sup>20</sup> availability of synthetic methodology to make these perdeuterated compounds will have additional applications.

## Acknowledgements

Support for the authors' work was obtained in part from the Natural Sciences and Engineering Research Council of Canada (Discovery grant to MSCP) and the University of Saskatchewan (Graduate Teaching Assistantship to DPOO). We would like to thank K. Brown and K. Thoms, Department of Chemistry, University of Saskatchewan, for technical assistance to obtain NMR and MS data, respectively.

## References

1. (a) Pedras MSC, Okanga FI, Zaharia IL, Khan AQ. *Phytochemistry* 2000; **53**: 161–176 (review); (b) Pedras MSC, Jha M, Ahiahonu PWK. *Curr Org Chem* 2003; **7**: 1635–1647 (review).
2. Bailey JA, Mansfield JW (eds). *Phytoalexins*. Blackie & Son: Glasgow, UK, 1982.
3. (a) Jeffery EH, Jarrell V. In *Handbook of Nutraceuticals and Functional Foods*, Wildman REC (ed.). CRC Press LLC: Boca Raton, FL, 2001; 169–191 (review); (b) Talalay P, Fahey JW. *J Nutr* 2001; **131**: 3027S–3033S (review).
4. Bonnesen C, Stephensen PU, Andersen O, Sorensen H, Vang O. *Nutr Cancer* 1999; **33**: 178–187.
5. Stephensen PU, Bonnesen C, Schaldach C, Andersen O, Bjeldanes LF, Vang O. *Nutr Cancer* 2000; **36**: 112–121.
6. Glawischnig E, Hansen BG, Olsen CE, Halkier BA. *Proc Natl Acad Sci USA* 2004; **101**: 8245–8250.
7. Monde K, Takasugi MJ, Ohnishi T. *J Am Chem Soc* 1994; **116**: 6650–6657.
8. Pedras MSC, Montaut S. *Chem Commun* 2004; 452–453.
9. (a) Kostianen R, Kotiaho T, Kuuranen T, Auriola S. *J Mass Spectrom* 2003; **38**: 357–372 (review); (b) Papac DI, Shahrokh Z. *Pharm Res* 2001; **18**: 131–145 (review).
10. Pedras MSC, Loukaci A, Okanga FI. *Bioorg Med Chem Lett* 1998; **8**: 3037–3038.
11. Van den Berg EMM, Baldew AU, De Goede ATJW, Raap J, Lugtenburg J. *Recl Trav Chim Pays-Bas* 1988; **107**: 73–81.
12. Makosza M, Winiarski J. *J Org Chem* 1984; **49**: 1494–1499.
13. Filler R, Woods SM, White WL. *Can J Chem* 1989; **67**: 1837–1841.
14. Pedras MSC, Chumala PB, Suchy M. *Phytochemistry* 2003; **64**: 949–956.
15. Miyashita K, Kondoh K, Tsuchiya K, Miyabe H, Imanishi T. *Chem Pharm Bull* 1997; **45**: 932–935.

16. Somei M, Kawasaki T. *Heterocycles* 1989; **29**: 1251–1254.
17. Burckard P, Fleury J-P, Weiss F. *Bull Soc Chim Fr* 1965; **10**: 2730–2733.
18. Yamazaki S. *Bull Chem Soc Jpn* 1997; **70**: 877–883.
19. Pedras MSC, Zaharia I. *Phytochemistry* 2000; **55**: 213–216.
20. Mikkelsen M, Petersen BL, Olsen CE, Halkier BA. *Amino Acids* 2002; **22**: 279–295 (review).