

Synthesis of Guaianolide Analogues with a Tunable α -Methylene- γ -lactam Electrophile and Correlating Bioactivity with Thiol Reactivity

Paul A. Jackson, Henry A. M. Schares, Katherine F. M. Jones, John C. Widen, Daniel P. Dempe, Francois Grillet, Matthew E. Cuellar, Michael A. Walters, Daniel A. Harki,* and Kay M. Brummond*

Cite This: <https://dx.doi.org/10.1021/acs.jmedchem.0c01464>

Read Online

ACCESS |




Metrics & More



Article Recommendations



Supporting Information



ABSTRACT: α -Methylene- γ -lactones are present in ~3% of known natural products, and compounds comprising this motif display a range of biological activities. However, this reactive lactone limits informed structure–activity relationships for these bioactive molecules. Herein, we describe chemically tuning the electrophilicity of the α -methylene- γ -lactone by replacement with an α -methylene- γ -lactam. Guaianolide analogues having α -methylene- γ -lactams are synthesized using the allenic Pauson–Khand reaction. Substitution of the lactam nitrogen with electronically different groups affords diverse thiol reactivity. Cellular NF- κ B inhibition assays for these lactams were benchmarked against parthenolide and a synthetic α -methylene- γ -lactone showing a positive correlation between thiol reactivity and bioactivity. Cytotoxicity assays show good correlation at the outer limits of thiol reactivity but less so for compounds with intermediate reactivity. A La assay to detect reactive molecules by nuclear magnetic resonance and mass spectrometry peptide sequencing assays with the La antigen protein demonstrate that lactam analogues with muted nonspecific thiol reactivities constitute a better electrophile for rational chemical probe and therapeutic molecule design.

■ INTRODUCTION

α -Methylene- γ -lactones are present in ~3% of all known natural products.¹ Given this abundance, and their validated biological function, it is curious that not a single compound with this motif has received FDA approval.^{2–4} This absence is due in part to the electrophilicity of the α,β -unsaturated carbonyl of the lactone that reacts rapidly with biothiols via a hetero-Michael addition, which limits the value of structure–activity relationship (SAR) studies and can result in off-target effects.^{1,2,5–8} We hypothesized that substituting the α -methylene- γ -lactone ($X = O$) of a compound predisposed to noncovalent binding to a particular protein target ($K_i = k_{\text{off}}/k_{\text{on}}$), with an α -methylene- γ -lactam ($X = NR$), would afford analogues with muted thiol reactivity (k_{inact}), leading to useful SAR data (Figure 1).⁹ Our approach to tuning the electrophilic reactivity of the exocyclic methylene group utilizes the lactam nitrogen that is substituted with electronically different R groups.^{10–15} Using this R group on the nitrogen, which is distal from the reacting methylenyl group, serves as a way to parse out electronic effects from steric effects on k_{inact} .

To test this hypothesis we turned to the guaianolides, a subclass of sesquiterpene lactone natural products with rich bioactivity largely attributed to the α -methylene- γ -lactone.^{1–4,16–18} Moreover, guaianolides have been shown to target the NF- κ B p65 protein at Cys38, making the NF- κ B signaling pathway an excellent model for testing the bioactivity of these novel α -methylene- γ -lactam-containing compounds.^{19–22} This assertion is further supported by results from our laboratories, where several guaianolide analogues were prepared and tested for inhibition of NF- κ B signaling.^{23,24} Compound 1 represents one member from a series of guaianolide analogues differing in substitution at C14 and stereochemistry at C6, C7, and C8 (Figure 2A). All analogues

Received: September 2, 2020



ACS Publications

© XXXX American Chemical Society

A

<https://dx.doi.org/10.1021/acs.jmedchem.0c01464>
J. Med. Chem. XXXX, XXX, XXX–XXX




Figure 1. Initial noncovalent binding specificity ($K_i = k_{\text{off}}/k_{\text{on}}$) of a covalent inhibitor to a target protein and second-order rate constant (k_{inact}) for covalent bond formation.




Figure 2. Guaianolide analogues: potent inhibitors of NF- κ B signaling. (A) Example structure of a guaianolide analogue, where the series varies by the substitution at the C14 position, and the stereochemistry at the C6, C7, and C8 positions. (B) Retrosynthetic analysis suggests that lactam 3 can be prepared by an allenic Pauson–Khand reaction (APKR) of allene-yne 5 based upon precedent established for the preparation of 2 from 4.


from this series demonstrated high levels of inhibition of the NF- κ B signaling pathway at low micromolar concentrations but with little evidence of SAR.²⁴ Furthermore, the criticality of the α -methylene- γ -lactone for inhibition was demonstrated by loss of all inhibitory activity when the α -methylene group of one of these analogues was reduced to a methyl substituent.²⁴

Testing our hypothesis, where an electrophilic α -methylene- γ -lactone is replaced with an electronically tunable α -methylene- γ -lactam to control thiol reactivity, required synthetic access to lactam-containing guaianolide analogues. Previously, we demonstrated that the APKR could be used to prepare the carbocyclic skeleton of lactone 2 from allene-yne 4 tethered with an α -methylene- γ -lactone (Figure 2B).^{23,25} Expanding the scope of the APKR to include allene-yne 5 equipped with an α -methylene- γ -lactam would provide rapid access to the molecularly complex and electronically tunable 5,7,5-fused ring system 3. Additionally, this APKR approach offers access to lactam analogues structurally similar to previously studied lactones, enabling direct comparisons of electrophilic reactivity toward thiols and inhibitory activity of the NF- κ B signaling pathway.^{23,24}

RESULTS AND DISCUSSION


Synthesis of α -Methylene- γ -lactam Guaianolide Analogues. Our approach to the 5,7,5-fused structure 6 was designed around the intramolecular APKR of lactam-containing allene-yne 7 (Scheme 1).^{23,25} Functionalization of the nitrogen of α -methylene- γ -lactam 7 late in the synthetic sequence was deemed necessary as it minimizes potential reactivity issues imparted by an electrophilic group. Naturally occurring guaianolides typically possess a methyl group at C15; however, here we chose to incorporate aryl and silyl groups ($R^1 = \text{Ar}$ or SiR_3) at this position because of their electronic and steric versatility. The allenyl group of 7 should be available from the methyl ketone 8 using a three-step reaction sequence involving a tertiary propargyl carboxylate.^{26,27} In turn, access to

Scheme 1. APKR Approach to α -Methylene- γ -lactam Guaianolide Analogues



the α -methylene- γ -lactam 8 should be possible by reaction of the imine 10, formed *in situ* from 3-phenyl- or 3-silyl-2-propynal and ammonium hydroxide and allylboronate 9. While we have utilized a similar reaction for the preparation of the corresponding lactones, formation of lactam 8 by this approach will represent the first example of a consecutive allylboronation and lactamization process involving the imine of a 2-alkynal.^{23,28}

To test the feasibility of this APKR strategy for the preparation of α -methylene- γ -lactam guaianolide analogues, synthesis of allene-yne 24a and 25a begins with alkynoate 11 prepared from 2,4-pentanedione (S1) in three steps on a multigram scale (Scheme 2A) (see the Supporting Information). Conversion of alkynoate 11 to allylboronate 9 was accomplished by addition of 11 to a solution of methyl lithium, copper iodide, hexamethylphosphoramide (HMPA), and diisobutylaluminum hydride (DIBALH), followed by addition of chloromethyl pinacolboronate.^{29,30} This reaction was performed several times affording allylboronate 9 with an average yield of 81% and a 2:1 Z/E isomeric mixture that also included an alkenoate byproduct S4 (5–10%, see the Supporting Information) resulting from protonation of the intermediate alkenyl metal species.^{23,25,30} Allylboronate isomers 9 and the alkenoate byproduct S4 were taken on as a mixture to the next step as chromatographic separation of the mixture resulted in a substantially reduced yield of the product. Addition of this mixture to 3-phenyl-2-propynal (12) and ammonium hydroxide afforded lactams 14a,b in 58% yield as a 5–2:1 mixture of the trans/cis isomers. Addition of mixture 9 to 3-triisopropylsilyl-2-propynal (13) and ammonium hydroxide afforded lactams 15a,b in an average yield of 71% with a ratio of 3.5–2:1 for the trans/cis isomers. 3-Trimethylsilyl-2-propynal ($R^1 = \text{SiMe}_3$) proved unstable to the lactamization reaction conditions, presumably because of the loss of the trimethylsilyl group under the basic reaction conditions (not


Scheme 2. Synthesis of α -Methylene- γ -lactam Tethered Allene-ynes^a

^aReagents and conditions: (a) CuI, MeLi in Et₂O, THF, −30 °C, 30 min; then toluene, HMPA, DIBALH in hexanes, −30 °C, 2 h; then **11**, −20 °C, 5 h; then 2-(chloromethyl)-4,4,5,5-tetramethyl-1,3-dioxaborolane (PinBCH₂Cl), −20 °C to rt, 16 h, 76% as a 3:1; Z/E mixture along with methyl (Z,E)-5-(2-methyl-1,3-dioxolan-2-yl)pent-2-enoate (**S4**), 10%; (b) **12** or **13**, ammonium hydroxide, ethanol, rt, 16 h; (c) PPTS, acetone/water (15:1), reflux, 16 h; (d) ethynyl magnesium bromide, THF, 0 °C, 3 h; (e) scandium triflate, pivalic anhydride, CH₃CN, rt, 16 h; (f) triphenylphosphine copper hydride hexamer (Stryker's reagent), toluene, −10 °C, 2 h; (g) tetra-*n*-butyl ammonium fluoride, THF, 0 °C, 45 min; (h) sodium hydride, iodomethane, DMF, 0 °C to rt, 15 min; (i) acetic anhydride, Et₃N, DMAP, CH₂Cl₂, 0 °C to rt, 3 h.

shown). These two allylboration/lactamization reactions represent the first examples of a three-component process using 2-alkynals to prepare alkynyl-functionalized α -methylene- γ -lactams.^{28,31} We hypothesize that the (*Z*)-allylboration reacts *via* Zimmerman–Traxler transition state **16**, affording *trans* lactams **14a** and **15a**, while the minor (*E*)-allylboration isomer reacts similarly (not shown) to afford *cis* lactams **14b** and **15b** (Scheme 2A).²⁸ However, because the *trans*/*cis* lactam ratio (**14a**/**14b**, 4:1 and **15a**/**15b**, 3.5:1) is higher than the Z/E ratio for the allylboration precursor, it is likely that for the minor (*E*)-allylboration isomer, the relatively small alkynyl group can adopt and react through an axial orientation as depicted by transition state **17**. Alternatively, boat-like transition states have been invoked for secondary aldimines to explain mixture ratios.^{30,32} These stereochemical assignments of the lactams were initially assigned by analogy to

similar compounds and later confirmed by X-ray crystallography of the APKR product (*vide infra*).^{23,25,28} The alkenoate byproduct **S4** could be readily separated from lactams **14a,b** and **15a,b** by column chromatography. Lactams **14a**, **14b**, **15a**, and **15b** were separated for characterization purposes, but in most cases, the lactams were advanced through the synthetic sequence as mixtures of *cis* and *trans* isomers, then separated near the end of the sequence, and only the *trans* lactam isomers were tested in biological assays.

Next, removal of the ketal-protecting group of **14a,b** and **15a,b** by acid-catalyzed hydrolysis using pyridinium *para*-toluenesulfonate (PPTS) in refluxing acetone and water (15/1, v/v) afforded keto lactams **18a,b** in 83% yield and **19a,b** in 73% yield. Both were afforded as a 4:1 mixture of *trans*/*cis* lactam isomers. Addition of ethynyl magnesium bromide to ketones **18a,b** and **19a,b** in tetrahydrofuran (THF) at 0 °C

Table 1. Optimization of Allene Formation with Stryker's Reagent ($[\text{CuHPPH}_3]_6$)


entry	SM ^a	equiv Stryker's	temp (°C)	time (h)	total yield (%)	(24a/25a)/(31a/32a)/SM	scale (mg)
1	23a	1.8	0	1.5	64	100:trace:0	110 ^b
2	23a	1	0	1	73	60:40:0	110
3	23a	0.8	−10	2	66	63:25:12	110
4	23a	0.9	−20	2	58	80:20:0	110
5	23a	0.9	−10	2	78	77:23:0	220
6	23a	0.9	−10	2	49	72:19:9	440
7	22a	0.9	−10	2	70	70:16:0 ^c	110
8	22a	0.9	−10	1.5	56	90:10:0	122

^aSM = starting material. ^bReaction performed with a previously opened aged container of Stryker's reagent; all other reactions were performed with a newly opened container. ^c14% of the total yield was a product resulting from reduction of the alkyne.

gave tertiary propargyl alcohol **20a,b** in 75% yield and **21a,b** in 82% yield. Each was afforded as a ~1:1 diastereomeric mixture at the newly created stereocenter. Conversion of the tertiary hydroxyl group to pivalate **22a,b** and **23a,b** was carried out using pivalic anhydride and a substoichiometric quantity of scandium triflate [$\text{Sc}(\text{OTf})_3$, 0.4 equiv] at rt for 16 h. These conditions afforded chemoselective pivaloylation of the hydroxyl group over the secondary lactam nitrogen, whereas acetylation conditions (Ac_2O , NEt_3 , DMAP) resulted in the *N*-acetylated product. At this point, the *cis* and *trans* lactam isomers **22a** and **22b** were separated *via* column chromatography to afford a 61% yield of the *trans* isomer **22a** and 16% yield of the *cis* isomer **22b**, each as a 1:1 mixture of diastereomers at C10. Similarly, pivalates **23a** (39%) and **23b** (16%) were separated by column chromatography. Only the *trans* isomers were taken on for the remainder of the synthetic sequence.


Conversion of the propargyl pivalate group to a 3,3-disubstituted allene was investigated. Propargyl pivalate **22a** was reacted with triphenylphosphine copper hydride hexamer (Stryker's reagent) at -10°C to afford a 33% yield of allene **24a**.³³ Propargylic pivalate **23a** gave a 46% yield of desired allene **25a** when subjected to Stryker's reagent. Removal of the TIPS substituent from the alkyne terminus by treatment of lactam **25a** with tetra-*n*-butyl ammonium fluoride (TBAF) provided terminal alkyne **26** in 79% yield. A competing reduction of the α -methylene group of the lactam during allene formation is the reason for the low yields. Progress toward solving this chemoselectivity challenge is discussed in more detail in the following section.

To examine the potential impact of the free NH of the secondary lactam on this reaction sequence, **14a** (containing a small amount of *cis* lactam **14b**) was converted to the corresponding tertiary lactam by the addition of sodium hydride and iodomethane (Scheme 2B). Removal of the ketal-protecting group as described above afforded **27a** in 73% yield. Addition of ethynyl magnesium bromide gave a 62% yield of **28a** as a 7:1 *trans*/*cis* ratio and a 1:1 diastereomeric mixture at the newly formed stereogenic carbon. Conversion of **28a** to the propargyl acetate **29a** was accomplished in 54% yield. Reaction of propargyl acetate **29a** with Stryker's reagent gave a 53% yield of desired allene **30a** with no evidence of reduced lactam

methylene. Thus, the formation of the allene was the only step in the reaction sequence impacted by the free NH of the secondary lactam. We presume that complexation of the less sterically hindered secondary lactam of **22a/23a** with copper increases the electrophilic reactivity of the methylene group over that of the tertiary lactam of **29a**.

Optimizing the Formation of 3,3-Disubstituted Allenes from Propargyl Pivalates. Efforts to increase chemoselectivity and yield for the transformation of propargyl pivalates **22a** and **23a** to allenes **24a** and **25a** involved changing the reaction temperature, time, scale, and equivalents and source of Stryker's reagent (eq 1, Table 1). Performing the reduction with **23a** using a previously opened, aged container of Stryker's reagent at 0°C in degassed toluene (0.04 M) with 2 equiv of water gave a 64% yield of **25a** and only trace amounts of reduced methyl lactam **32a** (Table 1, entry 1). However, when a newly opened container of Stryker's reagent was used under identical conditions, a 60:40 ratio of allene **25a** to methyl lactam **32a** was obtained in 73% yield (entry 2). Lowering the equiv of Stryker's reagent (0.8 equiv, new bottle) and temperature (-10°C) provided a slightly improved ratio of **25a** to **32a** (63:25), but with unreacted starting material (entry 3). Decreasing the temperature further (-20°C) with 0.9 equiv of Stryker's reagent afforded **25a** to **32a** (80:20) in 58% yield with no starting material (entry 4). Performing the reaction on larger scale (220 mg) with 0.9 equiv Stryker's reagent and -10°C (conditions used in remaining entries) gave a 77:23 ratio of **25a** to **32a** in 78% yield (entry 5). Increasing the scale of the reaction to 440 mg provided a ratio of 72:19:9 of **25a/32a/23a** in a 49% yield (entry 6). The phenyl-substituted alkyne **22a** afforded a 70:16 ratio of **24a** to **31a** in a 70% yield, but reduction of the alkyne to a *cis* alkene was observed and accounted for ~14% of the product mixture (entry 7).³³ Decreasing the reaction time (1.5 h) led to a 56% yield of **24a** and **31a** in a 90:10 ratio with minimal reduction of the phenyl alkyne (entry 8).

Thus, optimal conditions minimizing byproduct formation involved 0.9 equiv of Stryker's reagent, 0.04 M solution of propargyl carboxylate in toluene, 2 equiv of water at -10°C , and a reaction time of 1.5 to 2 h. Reaction of *cis*-lactams **22b** or **23b** with Stryker's reagent gave lower yields (30–35%) of the corresponding allenes **24b** or **25b** (see the Supporting

Scheme 3. Functionalization of the Lactam Nitrogen^a


^aReagents and conditions: (a) copper iodide (CuI, 20 mol %), *N,N'*-dimethylethylenediamine (40 mol %), cesium carbonate (Cs₂CO₃), ArI, toluene, 80 °C, 20 h; (b) sodium hydride (NaH, 60% dispersion), *para*-toluenesulfonyl chloride (*p*-TsCl), DMF, 0 °C 2 h; (c) di-*tert*-butyl dicarbonate (boc₂O), 4-(*N,N*-dimethylamino)pyridine (DMAP), Et₃N, CH₂Cl₂, 0 °C to rt, 2 h.

Information) where the over-reduction product was afforded predominantly. A selective reduction of the exocyclic methylene of the unsaturated lactams of **22b** and **23b** may be because of one face of the *cis*-lactam isomers being more accessible to the copper hydride complex. Alternative conditions for the formation of 3,3-disubstituted allenes were examined but were not productive.^{34–36}

Functionalization of the Lactam Nitrogen. Functionalization of the lactam nitrogen of **24a** with electronically different groups was examined. To minimize steric contributions, *para*-substituted aryl groups were selected to tune electrophilicity of the α -methylene- γ -lactam. A Buchwald–Hartwig coupling reaction was performed on lactam **24a** with aryl iodides having an electronically neutral group (**33**, R³ = H), electron-withdrawing groups (**34**, R³ = CN; **35**, R³ = CF₃), and an electron-donating group (**36**, R³ = OMe) (Scheme 3).^{37,38} *N*-Arylation of allene-ynone **24a** with iodobenzene (**33**) gave *N*-phenyl allene-ynone **37** in 44% yield using cesium carbonate as the base; a lower yield (27%) was observed with potassium carbonate. *N*-Arylation of **24a** with 4-iodobenzonitrile (**34**), 4-iodobenzotrifluoride (**35**), or 4-iodoanisole (**36**) provided **38**, **39**, or **40** in 42, 47, or 63% yield, respectively (eq 2). These same conditions were used to *N*-arylate **24a** with 2-iodothiophene (**41**) to afford the *N*-heteroaryl lactam **42** with an average yield of 59% (eq 3). The moderate yields for the *N*-arylation and *N*-heteroarylation products are attributed to the sterically hindered nature of the amide nitrogen.³⁸ Two additional electron-withdrawing groups were introduced by treatment of **24a** with sodium hydride and TsCl to give *N*-Ts lactam **43** in 29% yield (eq 4), and lactam **24a** was converted to a carbamate using di-*tert*-butyl dicarbonate to give the boc-functionalized lactam **44** with an average yield of 47% (eq 5). These *N*-functionalized lactams provide a spectrum of electronically disparate groups informing on the impact of electrophilic reactivity of the methylene lactam.

Examining the Feasibility and Scope of the APKR.

With a number of α -methylene- γ -lactam tethered allene-ynes in hand, the feasibility of the APKR was tested. Because these substrates have a methyl-substituted allene, we used conditions previously developed in our group that minimize dimer formation by slowly adding the allene-ynone to the Rh(I) catalyst.³⁹ First, allene-ynone **24a** was diluted in toluene and added dropwise over 1 h to rhodium biscarbonyl chloride dimer ([Rh(CO)₂Cl]₂, 10 mol %) in toluene under carbon monoxide (1 atm) at 110 °C (eq 6). Heating the reaction for an additional 30 min after the addition of the allene-ynone afforded lactam **45** in 72% yield (Table 2, entry 1). The structure of this compound was confirmed by X-ray crystallography (Figure 3). The reaction of TIPS-substituted alkyne **25a** using similar conditions afforded **46** in 46% yield (entry 2). The reduced yield for **46** compared to lactam **45** is attributed to a developing A^{1,3} strain between the large TIPS group on the terminus of the alkyne and the lactam during cyclization. This steric argument is further supported by a longer reaction time for **25a** compared to **24a** (3 vs 1.5 h). In addition, a structurally related α -methylene- γ -lactone-tethered allene-ynone with a less bulky trimethylsilyl group on the alkyne underwent an APKR in high yield with a reaction time similar to a phenyl-substituted alkyne.²⁵ A terminal alkyne was tolerated in the APKR, as evidenced by the reaction of allene-ynone **26** to provide lactam **47** in 53% yield (entry 3). Next, we examined the impact of the substitution on the lactam nitrogen on the APKR yield and reaction time. Allene-ynone **30a** with an *N*-methyl group afforded the APKR product **48** in 75% yield (entry 4), showing that the free NH has minimal impact on the efficiency of the APKR (compare entries 1 and 4). Allene-ynes **37–40**, where the lactam nitrogen was substituted with an *N*-phenyl, *N*-4-cyanophenyl, *N*-4-trifluoromethyl phenyl, or *N*-4-methoxyphenyl gave yields of 67, 72, 70, and 77% of tricyclic structures **49–52**, respectively (entries 5–8). Under these conditions, neither

Table 2. APKR with α -Methylene- γ -lactam Tether

entry	allene-yne	R ¹	R ²	APKR yield % ^a
1	24a	Ph	H	45 (72) ^{b,c}
2	25a	TIPS	H	46 (46) ^{b,d}
3	26	H	H	47 (53) ^b
4	30a	Ph	Me	48 (75) ^c
5	37	Ph	Ph	49 (67) ^{b,c}
6	38	Ph	4-CN-C ₆ H ₄	50 (72) ^c
7	39	Ph	4-CF ₃ -C ₆ H ₄	51 (70) ^b
8	40	Ph	4-OMe-C ₆ H ₄	52 (77) ^b
9	42	Ph	2-thiophenyl	53 (67) ^b
10	43	Ph	Ts	54 (71) ^{b,c}
11	44	Ph	boc	55 (40) ^{b,c}

^aDropwise addition of allene-yne to rhodium catalyst over 1 h, then heated for an additional 30 min. ^bAverage yield for two runs. ^cSimilar yields were obtained when allene-yne was added in a single portion.

^dReaction was heated for 3 h upon completion of allene-yne addition.

^eYield based upon one run.





Figure 3. Crystal structure of APKR product 45.

the reaction time nor the yield was significantly impacted by the electronic nature of the aryl group on the lactam nitrogen. The *N*-2-thiophenyl-substituted allene-yne 42 provided lactam 53 in 67% yield (entry 9). The APKR of *N*-Ts and *N*-boc allene-yne 43 and 44 gave lactams 54 and 55 in yields of 71 and 40%, respectively (entries 11–12). The lower yield of lactam 55 is attributed to the thermal instability of the boc group at 110 °C.⁴⁰


Next, we tested the impact of the α -methylene group on the efficiency of the APKR using allene-yne α -methyl- γ -lactams 56 and 57 (Scheme 4). *Trans, trans*-lactam 56 (8:1 dr at C11) underwent the APKR in 1.5 h to afford 58 in 78% yield as 8:1 mixture of diastereomers (eq 7). The APKR of α -methyl- γ -lactam 56 afforded a moderately higher yield than that of the α -methylene- γ -lactam 26, each possessing a terminal alkyne (78 vs 62%, respectively). *Cis, trans*-lactam 57 afforded the APKR product 59 in 77% yield as a single diastereomer (eq 8). This latter reaction was performed by the addition of the allene-yne to the rhodium catalyst in a single portion. The stereochemistry at C11 for both 58 and 59 was assigned by comparing calculated coupling constants to experimental values.

Scheme 4. APKR with Methyl Lactams



In summary, the reaction of allene-yne shown in Table 2 represent the first examples of an APKR with an α -methylene- γ -lactam tether. Substituents on the alkyne terminus had a moderate influence on the yield and reaction time (Table 2, entries 1–3). The groups on the nitrogen of the lactam had little impact on the yield or reaction time (entries 4–11), demonstrating that the APKR is tolerant of *N*-alkyl, *N*-aryl (neutral, electron-donating or -withdrawing), *N*-thiophenyl, *N*-sulfonyl, *N*-carbamate, and *N*-H α -methylene- γ -lactam tethers. Comparison of APKR reaction times of 56 and 57 and yields for lactams 58 and 59 suggest that the α -methylenyl group has some influence on the yield when compared with α -methyl lactams.

Electrophilic Reactivity of α -Methylene- γ -lactams toward Thiols. A subset of these newly synthesized α -methylene- γ -lactams was reacted with excess cysteamine to obtain pseudo-first-order reaction kinetics as a way to quantify the electrophilic reactivity of these analogues toward a biologically relevant thiol.¹⁵ Given that ¹³C NMR chemical shifts for the C _{β} resonance of the methylene group can be a good predictor of electrophilicity in some systems,¹² we selected lactams 45 (R¹ = H, ¹³C _{β} δ = 115.3); 49 (R¹ = Ph, ¹³C _{β} δ = 117.1); 51 (R¹ = 4-CF₃-C₆H₄, ¹³C _{β} δ = 118.2); 52 (R¹ = 4-MeO-C₆H₄, ¹³C _{β} δ = 116.7); 54 (R¹ = Ts, ¹³C _{β} δ = 121.4); and 60 (R¹ = Ac, ¹³C _{β} δ = 120.6) that show a range of ¹³C _{β} chemical shifts. Cysteamine was selected as the biologically relevant thiol nucleophile, given the strong preference for the sulfhydryl group reacting over the amino group at pH 7.4 and its compatibility with organic solvents.^{41,42} Each reaction was performed by adding a solution of the lactam and cysteamine (15 equiv) in CDCl₃ to an NMR tube at rt (22 °C). The reaction progress was monitored by ¹H NMR with spectra acquired at regular intervals and the reaction tube held in the autosampler between measurements.⁴² Reaction of α -methylene- γ -lactams 45 and 49 with cysteamine each shows a half-life of 2.1 d (entries 1–2, Table 3). The trifluoromethyl aryl lactam 51 reacted with cysteamine with a half-life of 9.8 h, and the methoxy aryl lactam 52 reacted with a half-life of 8.5 d (entries 3 and 4). *N*-Ts-substituted lactam 54 reacted completely in less than 10 min; attempts were not made to determine half-lives for this lactam (entry 5). Lactam 60 reacted with cysteamine with a half-life of 5 min (entry 6). A chemoselective addition to the lactam α -methylenyl group is supported by ¹H NMR showing the complete disappearance of alkenyl proton resonances (see the Supporting Information). Finally, a structurally related lactone S17 (see the Supporting Information for structure) reacted with cysteamine with a half-

Table 3. Reaction of Cysteamine with α -Methylene- γ -lactams^a

entry	SM	R ¹ , R ²	¹³ C _β ppm	half-life
1	45	Ph, H	115.3	2.1 d ^b
2	49	Ph, Ph	117.1	2.1 d
3	51	Ph, 4-CF ₃ -C ₆ H ₄	118.2	9.8 h
4	52	Ph, 4-OMe-C ₆ H ₄	116.7	8.5 d
5	54	Ph, Ts	121.4	<10 min ^{b,c}
6	60	H, Ac	120.6	5 min
7	S17		120.7	7 min
8	PTL		121.1 ^d	<5 min ^e

^aConditions: HSCH₂CH₂NH₂ (15 equiv), CDCl₃ (0.006 M), rt.

^bReaction was performed with cysteamine that was ~60% oxidized.

^cReaction was complete at this time point. ^dBased on previous literature.⁴⁴ ^eBased on previous literature.⁴²

life of 7 min (entry 7). α -Methylene- γ -butyrolactone parthenolide (PTL) was previously shown to react completely with cysteamine in less than 5 min (entry 8).⁴²

Structural confirmation of thiol adducts was thwarted by their instability to column chromatography. Thus, *N*-methyl lactam 48 was reacted with *tert*-butyl thiol to form the thiol adduct S11 after 30 min in 23% yield (dr 1.6:1). The major diastereomer was assigned by analogy to methyl lactam 58, where calculations predict this isomer to be thermodynamically more stable by ~2 kcal/mol.⁴³ Thiol adduct S11 showed that a single equivalent of thiol was added to the exocyclic alkene of 48.

In summary, groups on the lactam nitrogen greatly impact the reactivity of the α -methylene- γ -lactams toward cysteamine where reaction half-lives ranged from days, for electron-neutral and -donating groups, to minutes for electron-withdrawing groups. The chemical shift for the C_β in the ¹³C NMR is an excellent predictor of reactivity; a slow rate of reaction is observed for compounds with a C_β chemical shift in the range of 115.3–117.1 and requires days to complete one reaction half-life. For compounds with C_β chemical shifts in the range of 120.6–121.4, a reaction half-life is achieved in minutes. Notably, lactam 51, *N*-substituted with a trifluoroaryl group and a C_β chemical shift of 118.2, showed moderate reactivity with a half-life measured in hours.

NF- κ B Bioactivity and Cytotoxicity of α -Methylene- γ -lactams. Several natural products and synthetic compounds containing α -methylene- γ -lactones are known to inhibit the NF- κ B pathway; however, these highly reactive lactones limit the availability of informed SAR as they react readily with accessible thiols.^{5,19,45–50} We have shown with our ¹H NMR studies that the electrophilic reactivity of α -methylene- γ -lactams toward thiols can be modulated depending upon the electronic character of *N*-substituents; this tunable electrophilicity should contribute to a better understanding of the role of covalent adduct formation to inhibition of the NF- κ B signaling pathway. To characterize the biological utility of the α -methylene- γ -lactams, we performed cellular NF- κ B inhibition assays with a representative subset of the

synthesized compounds and benchmarked activities to the known α -methylene- γ -lactone NF- κ B inhibitor PTL.^{50,51} We used two NF- κ B reporter cell lines for our studies: A549 cells bearing a stably transfected NF- κ B-driven luciferase reporter gene and HEK293 cells containing a stably transfected NF- κ B-driven secreted embryonic alkaline phosphatase (SEAP) reporter gene. These orthogonal assays with different readouts were used to corroborate data to ensure that inhibition was not occurring at the enzymatic readout level (*i.e.*, direct inhibition of luciferase or SEAP). Direct inhibition of luciferase by sesquiterpene lactones at high micromolar concentrations has been reported.⁵² Our assays also inform how different substituents containing electron-withdrawing, electron-donating, or electronically neutral functional groups attached to the lactam nitrogen affect inhibition of the NF- κ B pathway. Select compounds were incubated with cells for 30 min at 5–20 μ M for the luciferase reporter assay and 1–7.5 μ M for the SEAP reporter assay, before induction with 15 or 22.5 ng/mL TNF- α , respectively, for 8 h and subsequent readout of the reporter gene (Figure 4).⁵³

Compounds with no substituents on the lactam nitrogen (45 and 47) did not inhibit NF- κ B signaling at 20 μ M, the highest concentration tested; a result supported by the cysteamine study showing low thiol reactivity. Lactams 48, 49, and 52 containing a *N*-methyl, *N*-phenyl, and *N*-4-methoxy phenyl substituents, respectively, also showed no inhibitory activities. However, compounds with electron-withdrawing substituents did inhibit NF- κ B activation, with lactam 54 containing a *N*-Ts substituent displaying potent activity with complete inhibition of induced NF- κ B signaling at 10 μ M in the luciferase reporter assay and at 7.5 μ M in the SEAP reporter assay. Cytotoxicity was observed with 54 at 10 μ M treatment in A549/NF- κ B-luc cells (64% cell viability at 8 h), although less toxicity for 54 was observed in HEK293/NF- κ B-SEAP cells at the concentrations tested (82% cell viability at 7.5 μ M). Derivatives 51 and 60 with *N*-(4-trifluoromethyl)phenyl and *N*-acetyl groups, respectively, showed similar inhibitory activities in the luciferase reporter assay (21% [51] and 28% [60] residual NF- κ B activity at 20 μ M, as well as 62% [51] and 59% [60] residual NF- κ B activity at 10 μ M). Both compounds maintained inhibitory activity in the SEAP reporter assay, but derivative 60 was more potent (44% [51] and 19% [60] residual NF- κ B activity at 7.5 μ M). In A549/NF- κ B-luc cells, significant toxicities were not observed for lactams 51 and 60 (88% [51] and 82% [60] cellular viability at 20 μ M). In HEK293/NF- κ B-SEAP cells, lactam 51 was nontoxic at all concentrations; conversely, lactam 60 demonstrated toxicities at 7.5 μ M (69% cell viability) and 5.0 μ M (76% cell viability) treatments. In comparison, a lactone analogue of 45 was previously reported to reduce NF- κ B activity completely in A549/NF- κ B-luc cells when tested at 20 μ M, with limited cytotoxic effects.²³ The compounds were also benchmarked against the α -methylene- γ -butyrolactone PTL, a known NF- κ B inhibitor. Previously published results show that PTL at 10 μ M concentration reduces NF- κ B to 53% residual activity in the A549/NF- κ B-luc assay with no cytotoxicity.²³ These data are consistent with HEK293/NF- κ B-SEAP assay results, which show a 20% residual NF- κ B activity with a moderate level of cytotoxicity at 7.5 μ M. From these results, we conclude that α -methylene- γ -lactams containing electron-withdrawing substituents increase inhibitory activity toward the NF- κ B pathway in both assays compared to electronically neutral and electron-




Figure 4. Modulation of canonical NF- κ B signaling by α -methylene- γ -lactams. (A) Compounds were tested at 20, 10, and 5 μ M and NF- κ B signaling was induced with TNF- α (15 ng/mL) in A549 cells containing a NF- κ B-driven luciferase gene. (B) Compounds were tested at 7.5, 5.0, 2.5, and 1.0 μ M and NF- κ B signaling was induced with TNF- α (22.5 ng/mL) in HEK293 cells containing a NF- κ B-driven SEAP gene. All wells in both assays were induced with TNF- α except for noninduced (N) control wells. Relative NF- κ B activities (referenced to the induced, I, control set to 100%) are shown in dark colors. Accompanying cellular cytotoxicity measurements were made using Alamar Blue viability dye and are shown behind NF- κ B inhibition in light colors. Cytotoxicity was normalized to the induced control, which was set at 100%. Columns marked with NT (nontoxic) indicate instances in which NF- κ B activity (dark bars) obstructs the cellular cytotoxicity (light bars). Ocluded values range from 88 to 101% (Table S1). Values shown are mean $\pm$ S.D. for $n \geq 3$ biological replicates. PTL = parthenolide. See Supporting Information Tables S1 and S2 for numerical values.

donating substituents, which display minimal or no effect on NF- κ B inhibition.

To further investigate the bioactivity of our α -methylene- γ -lactam derivatives, their cytotoxicity was assessed in a model noncancerous cell line for a longer treatment duration (Table 4). Vero is a kidney epithelial cell line derived from an African green monkey that is commonly used as a standard for cytotoxicity in healthy cells.^{54,55} Lactam derivatives were incubated with cells for 48 h at concentrations ranging from 0.1 to 200 μ M and cellular viability was measured by Alamar Blue assays.⁵³ Derivative 47 showed little or no inhibition of the NF- κ B pathway in either of the two reporter assays and displayed low cytotoxicity. Compounds 49 and 52, which had little NF- κ B inhibitory activity, displayed cytotoxicity to Vero cells (IC_{50} = 21.9 and 44.0 μ M, respectively). Lactams 51 and 60, two of the more potent derivatives in the NF- κ B inhibitory

Table 4. Cellular Toxicity of Lactams to Vero Cells^a

compound	IC_{50} (μ M)
47	>200
49	21.9 $\pm$ 1.9
51	15.7 $\pm$ 2.0
52	44.0 $\pm$ 1.8
60	7.8 $\pm$ 1.9
S17	6.2 $\pm$ 1.8

^aCellular viabilities were measured by Alamar Blue viability dye. IC_{50} values shown are mean $\pm$ S.E.M. for $n \geq 3$ biological replicates.

assays that were also nontoxic to the host cell lines, were cytotoxic to Vero cells (51: IC_{50} = 15.7 μ M, 60: IC_{50} = 7.8 μ M); however, 51 was approximately twofold less toxic than 60 and constitutes the most compelling lactam for further mechanistic and structural optimization studies. A more comprehensive evaluation of the toxicity of 51 will be pursued in future *in vivo* pharmacokinetic studies in mice.

A comparison of the half-life for the reaction of an α -methylene- γ -lactam with cysteamine and NF- κ B inhibition showed good correlation between thiol reactivity of the lactam and its inhibition (Table 3). For example, lactams 45, 49, and 52 displayed low thiol reactivity toward cysteamine with $t_{1/2}$ = 2.1, 2.1, and 8.5 d, respectively, and each demonstrated no NF- κ B inhibition. Lactam 51 displayed intermediate thiol reactivity ($t_{1/2}$ = 9.8 h) and showed 62% residual NF- κ B activity for the luciferase assay (10 μ M) and 44% for the SEAP reporter assay (7.5 μ M). These inhibitory values were balanced by less overall cytotoxicity in the Vero model (IC_{50} = 15.7 μ M) in comparison to lactam 60 ($t_{1/2}$ = 5 min for cysteamine reactivity), which was 2-fold more cytotoxic to Vero cells (IC_{50} = 7.8 μ M), yet similarly potent toward inhibiting NF- κ B signaling (59% residual NF- κ B activity in the luciferase assay [at 10 μ M] and 19% residual NF- κ B activity for the SEAP reporter assay [at 7.5 μ M]). As expected, lactam 52 displaying low thiol reactivity also showed lower relative cytotoxicity (IC_{50} = 44.0 μ M).

Measuring Thiol-Reactivity of Lactam Inhibitors with ALARM NMR. To determine the thiol-reactivity of this series of lactam compounds in a proteinaceous environment, we turned to a La assay to detect reactive molecules by nuclear magnetic resonance (ALARM NMR), an assay using the La antigen, which contains a cysteinyl group (C245) that is highly reactive toward electrophilic compounds.^{56–58} This bioassay involves monitoring the changes in the chemical shifts of 13 C-labeled methyl groups (L249, L294, and L296) by 2D 1 H- 13 C HMQC that occur when C245 is modified. The α -methylene- γ -lactam 51 was selected for this study as it shows moderate reactivity with cysteamine when compared to the other lactams (see Table 3) and is the most balanced lactam with respect to NF- κ B inhibitory activity and cellular cytotoxicity. Reaction of excess 51 with 13 C-labeled La antigen at 37 $^{\circ}$ C for 90 min in the presence and absence of dithiothreitol (DTT, 20 mM) resulted in no changes in the HMQC spectra, similar to that seen for the negative control compound, fluconazole (Figure 5). Thus, lactam 51 is characterized as ALARM NMR negative and nonreactive, as this experiment provides compelling evidence that the La antigen protein does not react with the α -methylene- γ -lactam. Reaction of the analogous α -methylene- γ -lactone S17 shows high reactivity with the La antigen thiol, similar to that seen for the positive control compound, CPM (N-[4-(7-diethylamino-




Figure 5. ALARM NMR thiol-reactivity assays. Shown are the ^1H – ^{13}C HMQC spectra of key ^{13}C -labeled methyl groups of the La antigen protein incubated with DMSO; CPM, positive thiol-reactive compound control; fluconazole, negative thiol-reactive and aggregation compound control; α -methylene- γ -lactone **S17**; and α -methylene- γ -lactam **51**. Compounds tested at 400 μM final concentrations, with a La antigen protein concentration of 50 μM ; spectra for **S17** and **51** are from the $t = 9$ h timepoint, spectra for the DMSO, CPM, and fluconazole spectra are representative.




Figure 6. ALARM MSPS assay analysis of nonspecific compound adduction to La antigen C245. (A) Experimental workflow to identify thiol reactivity and reversible/irreversible covalent binding of compounds to C245 of the La antigen. Compounds added at 50 mM final concentration, La antigen final concentration of 30 μM , incubated for 1 h at room temperature in the dark. (B) Heat map indicating the number of samples that afforded IAD-adducted peptide FSGDLDDQTC²⁴⁵R. Three biological replicates for each compound were run in technical triplicate. Fluconazole was used as a nonreactive negative control, and CPM was used as a thiol-reactive positive control, as in the ALARM NMR assay. Data on the additional peptides analyzed containing C232 can be found in the [Supporting Information](#). ^aThe C245 peptide was not identified in one technical replicate of one of the biological replicates but did appear as IAD-adducted in the other two technical replicates of that sample. Therefore, the biological replicate was considered positive for IAD adduction. ^bThe FSGDLDDQTC²⁴⁵R peptide was not observed in one biological replicate, likely due to low peptide abundance because expected d_2 -IAD adductions were clearly observed in other biological replicates, as well as for C232-containing peptides (see [Table S12](#)).

4-methylcoumarin-3-yl)phenyl]maleimide). This latter result demonstrates that the La protein conformation is perturbed as evidenced by the disappearance of the characteristic chemical shifts of ^{13}C -labeled methyl groups of L249, L294, and L296 in the absence of DTT (Figure 5). Therefore, **S17** reacts with the

La antigen protein and is characterized as ALARM NMR positive and reactive. These experiments support the premise that α -methylene- γ -lactam-containing compounds may react discriminately with proteins containing cysteine residues,

whereas the α -methylene- γ -lactones have the propensity to react more indiscriminately.

Measuring Thiol-Reactivity and Reversibility of α -Methylene- γ -lactam **51 and α -Methylene- γ -lactone **S17** Using Mass Spectrometry Peptide Sequencing.** To corroborate the thiol reactivity properties of lactam **51** and lactone **S17**, a novel **L**a **A**ssay to detect **R**eactive **M**olecules by **M**ass **S**pectrometry **P**eptide **S**equencing (ALARM MSPS) assay of the La antigen was developed, offering a complementary assay to ALARM NMR. There are two possible thiol adduction sites on the La antigen: C232 and C245. Consistent with previously reported literature and with our own experiments, C245 is the more reactive cysteine (further discussed in the [Supporting Information](#)).⁵⁶ Although the La antigen has been previously analyzed by mass spectrometry (MS), whole-protein analysis was performed, which does not distinguish covalent adduction between C232 and C245.⁵⁶ In fact, whole-protein MS studies with La antigen and electrophilic compounds revealed both single- and double-adducted products in the same sample, supporting the hypothesis that there is a difference in thiol reactivity between C232 and C245. Therefore, we turned to MS analysis of tryptic peptides of the La antigen to characterize only those compounds that adduct C245, which is more reactive. Moreover, our new approach, ALARM MSPS, provides a method for evaluation of compounds that may function as reversible covalent inhibitors. Recently, a similar assay with isotopic iodoacetamide (IAD) was reported to study protein dynamics, which supports our approach.⁵⁹

The experimental workflow of the ALARM MSPS assay involves the following steps ([Figure 6](#), panel A): (1) incubation of the La antigen, containing the reactive cysteine C245, with test compounds **51** or **S17**; (2) addition of IAD to carbamidomethylate cysteine residues that were not adducted; (3) protein denaturation; (4) incubation with “heavy” d_4 -IAD to distinguish between two possible reactions, one where C245 did not react with an electrophile (labeled with IAD) versus the other where the adducted C245 of the La antigen then underwent a retro-hetero-Michael addition under the denaturation conditions (where the liberated C245 is subsequently adducted with heavy d_4 -IAD); (5) digestion of adducted antigen with trypsin; and (6) MS/MS analysis of the adducted peptide FSGDLDDQTCR containing the C245 residue (For information regarding the peptide containing C232, see the [Supporting Information](#)).

Subjecting lactone **S17** to the ALARM MSPS assay afforded only the heavy (d_4 -IAD) C245 adduct and none of the IAD adduct ([Figure 6B](#), green color), providing support that α -methylene- γ -lactone **S17** reacted with C245 but undergoes a retro-Michael addition upon protein denaturation. The same results were obtained for the positive control experiment using CPM, where only the d_4 -IAD-C245 adduct was observed. In contrast, α -methylene- γ -lactam **51** showed only IAD-labeled C245 ([Figure 6B](#), red color), indicating that **51** either does not react with C245 or reacts with very slow kinetics. Similar results were observed for the negative control compound, fluconazole. This newly developed ALARM MSPS assay provides compelling data to support that α -methylene- γ -lactam-containing compounds may react discriminately with cysteinyl-containing proteins, whereas the α -methylene- γ -lactones react more indiscriminately. Our data are consistent with previous ALARM NMR profiling of the covalent inhibitor of Bruton's tyrosine kinase, ibrutinib, which also tests as

ALARM NMR negative.⁵⁸ Therefore, although compounds may engage protein targets covalently, molecules with less reactive electrophiles, such as **51**, can be identified with ALARM-based assays.

CONCLUSIONS

In this report, the scope of the APKR was expanded to allene-ynes tethered by an α -methylene- γ -lactam, and the 3-component allylboration/lactamization reaction sequence extended to 2-propynals. The electrophilic reactivity of α -methylene- γ -lactams toward thiols in the hetero-Michael addition reaction is impacted by the electronic nature of the group on the nitrogen of the lactam. For example, substituents on the lactam nitrogen of the 5,7,5-fused ring system of the APKR products can be used to control thiol reactivity (k_{inact}) of the α -methylene- γ -lactam, where electron-neutral and -donating N-substituents are slow to react with cysteamine, whereas lactams with electron-withdrawing N-substituents are faster to react. This ability to tune the electrophilic reactivity toward thiols contrasts with α -methylene- γ -lactone-containing compounds, where the electrophilic reactivity is high and cannot be readily tuned. We have shown that α -methylene- γ -lactam guaianolide analogues function as small-molecule regulators of the NF- κ B signaling pathway with reasonable cellular toxicity. The NF- κ B inhibitory activity for the lactam analogues was found to positively correlate with thiol reactivity. Cytotoxicity shows a positive correlation at the outer most limits of thiol reactivity. Further studies evaluating SARs of the α -methylene- γ -lactams described herein having intermediate thiol reactivity are warranted. Finally, because guaianolides are well-known NF- κ B inhibitors, these proof-of-concept studies show that modulating the thiol reactivity profile is beneficial from an inhibitor optimization standpoint. Further, these findings support our hypothesis that NF- κ B signaling is an ideal system for evaluating an approach where an electronically tunable methylene lactam combined with a small-molecule scaffold predisposed to protein-target binding can afford discriminant covalent inhibitors. Finally, these results inform our next step to design and synthesize electronically tuned lactam analogues for SAR studies, which are expected to have enhanced potential as lead compounds for drug design.

EXPERIMENTAL SECTION

Chemistry. Commercially available compounds were used as received unless otherwise noted. Dichloromethane (CH_2Cl_2), THF, and diethyl ether (Et_2O) were purified by passing through alumina using the Sol-Tek ST-002 solvent purification system. Triethylamine was distilled from calcium hydride (CaH_2) and stored over 4 Å molecular sieves. Acetic anhydride (Ac_2O) was shaken with phosphorus pentoxide (P_2O_5), decanted, fractionally distilled from anhydrous potassium carbonate (K_2CO_3), and stored over 4 Å molecular sieves. HMPA was vacuum-distilled from CaH_2 and stored over 4 Å molecular sieves. Deuterated chloroform (CDCl_3) was stored over anhydrous K_2CO_3 . All designated temperatures are bath temperatures unless specified otherwise. All reactions are performed under an atmosphere of nitrogen unless indicated otherwise. Silica gel (40–63 μm particle size, 60 Å pore size) purchased from Sorbent Technologies is used for the purification of compounds by flash chromatography. Thin-layer chromatography (TLC) analyses were performed on Silicycle Siliaplate G silica gel glass plates (250 μm thickness). ^1H NMR and ^{13}C NMR spectra were recorded on Bruker AVANCE 300, 400, or 500 MHz spectrometers. Spectra were referenced to residual chloroform (7.26 ppm, ^1H , 77.16 ppm, ^{13}C). Chemical shifts are reported in parts per million, multiplicities are

indicated by s (singlet), br (broad signal), d (doublet), t (triplet), q (quartet), quint (quintet), m (multiplet), and dt (doublet of triplets). Coupling constants, *J*, are reported in Hertz (Hz). All NMR spectra were obtained at rt unless otherwise specified. IR spectra were obtained using a Nicolet Avatar E.S.P. 360 FT-IR. ESI mass spectroscopy was performed on a Waters Q-TOF Ultima API, Micromass UK Limited high-resolution mass spectrometer. The purity of representative final compounds was checked by high-performance liquid chromatography (HPLC) and ranged from 91 to 99% (see Table S11).

Hex-5-yn-2-one (S2). Compound S2 was synthesized according to a literature procedure.⁶⁰ A flame-dried, 250 mL, single-necked, round-bottomed flask equipped with a magnetic stir bar, septum, and nitrogen inlet needle was charged with 2,4-pentanedione (S1, 30 mL, 290 mmol) and ethanol (120 mL, 2.2 M). Propargyl chloride (37 mL, 260 mmol) was added *via* a syringe, followed by potassium carbonate (48 g, 350 mmol) in a single portion. The septum was removed, and a reflux condenser equipped with a septum and nitrogen inlet needle was attached. The reaction was heated to reflux (oil bath temperature 85 °C) for 24 h. Upon completion of the reaction as observed by TLC, the mixture was cooled to rt and filtered *via* vacuum filtration to remove the solids. The solid was washed with ethyl acetate. Ethyl acetate and ethanol were removed by simple distillation at atmospheric pressure. The residue was diluted with diethyl ether (100 mL), transferred to a separatory funnel, washed with deionized water (50 mL) and then brine (50 mL), dried over magnesium sulfate, gravity filtered, and concentrated by simple distillation at atmospheric pressure. The residue was further purified by simple, vacuum distillation (45 mmHg, 85–90 °C) to give 12.5 g of product in a 49% yield. The spectral data matched literature values.⁶⁰

¹H NMR (300 MHz, CDCl₃): δ 2.67, (t, *J* = 7.2 Hz, 2H, CH₂), 2.42 (dt, *J* = 2.7, 7.1 Hz, 2H, ≡CCH₂), 2.16 (s, 3H, CH₃), 1.93 (t, *J* = 2.7 Hz, 1H, ≡CH); ¹³C NMR (100 MHz, CDCl₃): δ 206.3 (C=O), 82.7, 68.6, 42.0, 29.8, 12.8; TLC *R*_f = 0.46 (25% EtOAc/hexanes) [silica gel, UV, KMnO₄].

2-(But-3-ynyl)-2-methyl-1,3-dioxolane (S3). Following a procedure analogous to that previously reported,⁶¹ a flame-dried, 200 mL, single-necked, round-bottomed flask equipped with a magnetic stir bar was charged with hex-5-yn-2-one (S2, 12.5 g, 131 mmol), benzene (130 mL), ethylene glycol (8.8 mL, 160 mmol), and *p*-toluenesulfonic acid (0.497 g, 2.61 mmol). The flask was equipped with a Dean–Stark trap that was attached to a condenser. The solution was heated at reflux (100 °C). After 15 h, TLC showed complete consumption of the starting material. The solution was allowed to cool to rt, diluted with diethyl ether (80 mL), transferred to a separatory funnel, and washed successively with saturated sodium bicarbonate (100 mL) and brine (100 mL). The organic layer was dried over magnesium sulfate and gravity filtered, and diethyl ether and benzene were removed by simple distillation at atmospheric pressure. The residue was purified by vacuum distillation at 30 mmHg. Two fractions were collected: the first contained benzene with trace amounts of product S3 (bp 40–50 °C, 30 mmHg); the second contained product with less than 10% benzene (bp 113–120 °C, 30 mmHg, 13.0 g, 71% yield), as determined by integration of the benzene resonance at 7.19 ppm and terminal alkyne resonance at 1.87 ppm. ¹H NMR (300 MHz, CDCl₃): δ 3.96–3.86 (m, 4H, OCH₂CH₂O), 2.25 (dt, *J* = 2.7, 7.5 Hz, 2H, ≡CCH₂), 1.92–1.88 (m, 2H, CH₂), 1.87 (s, 1H, ≡CH), 1.30 (s, 3H, CH₃); ¹³C NMR (100 MHz, CDCl₃): δ 109.2, 84.5, 67.9, 65.0 (2C), 38.1, 23.9, 13.3; TLC *R*_f = 0.55 (25% EtOAc/hexanes) [silica gel, UV, KMnO₄].

Methyl 5-(2-Methyl-1,3-dioxolan-2-yl)pent-2-ynoate (11). Following a procedure analogous to that previously reported,⁶² a flame-dried, 200 mL, single-necked, round-bottomed flask equipped with a magnetic stir bar, septum, and nitrogen inlet needle was charged with alkyne S3 (2.10 g, 15.0 mmol) and THF (75 mL) and then cooled to –78 °C. *n*-Butyl lithium (1.6 M in hexanes, 11.3 mL, 18.0 mmol) was added dropwise *via* a syringe. Upon completion of addition, the reaction was maintained at –78 °C for an additional 30 min. Methyl chloroformate (1.5 mL, 19.5 mmol) was added dropwise *via* a syringe and maintained at –78 °C for an additional 30 min and then warmed

to rt. After 3 h, TLC showed complete disappearance of the starting material. Saturated aqueous ammonium chloride (30 mL) was added, and the solution was transferred to a separatory funnel. The layers were separated, and the aqueous layer was extracted with diethyl ether (3 × 50 mL). The organic layers were combined, washed with brine (50 mL), dried over magnesium sulfate, gravity filtered, concentrated by rotary evaporation (30 °C), and purified by silica gel flash column chromatography eluting with 20% diethyl ether in hexanes to yield alkynoate 11 (2.14 g, 72% yield) as a clear liquid. ¹H NMR (300 MHz, CDCl₃): δ 3.98–3.89 (m, 4H, OCH₂CH₂O), 3.74 (s, 3H, OCH₃), 2.42 (t, *J* = 7.9 Hz, 2H, CH₂), 1.95 (t, *J* = 7.9 Hz, 2H, CH₂), 1.31 (s, 3H, CH₃); ¹³C NMR (100 MHz, CDCl₃): δ 154.4 (C=O), 108.8, 89.7, 72.7, 65.0 (2C), 52.7, 36.9, 24.0, 13.6; TLC: *R*_f = 0.23 (20% diethyl ether/hexanes) [silica gel, UV, KMnO₄].

2-(Chloromethyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (S5). Following a procedure analogous to that previously reported,⁶³ a flame-dried, single-necked, 250 mL, round-bottomed flask equipped with a stir bar, septum, and nitrogen inlet needle was charged with trimethyl borate (8.6 mL, 77 mmol) and bromochloromethane (5.5 mL, 85 mmol) and cooled to –78 °C. *n*-Butyl lithium (1.6 M in hexanes, 53 mL, 85 mmol) was added dropwise *via* a syringe pump over 35 min. Upon completion of addition, the solution was maintained at –78 °C for an additional 30 min. Chlorotrimethylsilane (12 mL, 92 mmol) was added dropwise at –78 °C and the reaction was warmed to rt by removal of the dry ice/acetone bath. The reaction was allowed to stand for 16 h (without stirring), and then pinacol (10.0 g, 85 mmol) was added under stirring at rt. The reaction was maintained for 1 h and then poured into a separatory funnel containing water (100 mL) and diethyl ether (100 mL). The organic layer was separated, washed with brine (50 mL), dried over magnesium sulfate, gravity filtered, and concentrated by rotary evaporation (37 °C), and the residue was transferred to a 25 mL, single-necked, round-bottomed flask. This residue was fractionally distilled under reduced pressure using a short, jacketed vigreux column connected to a short-path distillation head to give chloromethylpinacol boronate S5 as a colorless liquid (108–115 °C, 14 mmHg, 8.89 g, 65% yield). ¹H NMR (300 MHz, CDCl₃): δ 2.94 (s, 2H, CH₂Cl), 1.28 (s, 12H, (CH₃)₂CC(CH₃)₂); ¹³C NMR (100 MHz, CDCl₃): δ 84.7, 24.8 (4C); ¹¹B NMR (128 MHz, CDCl₃): δ 31.42; IR: 3459, 2980, 2935, 1352, 1273, 1143 cm^{–1}; TLC: *R*_f = 0.45 (10% diethyl ether/hexanes) [silica gel, KMnO₄].

Methyl 5-(2-Methyl-1,3-dioxolan-2-yl)-2-((4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)methyl)pent-2-enoate (9). 9 was prepared using a procedure analogous to that previously reported.^{30,64} Run 1: A flame-dried, 100 mL, single-necked, round-bottomed flask equipped with a magnetic stir bar, septum, and nitrogen inlet needle was charged with copper iodide (48 mg, 0.25 mmol) and THF (8.4 mL) and then cooled to –30 °C. Methyl lithium (1.6 M solution in diethyl ether, 0.16 mL, 0.25 mmol) was added dropwise over 1 min. Upon completion of addition, the solution was maintained at –30 °C for an additional 20 min at which time it turned from dark brown to black. Toluene (18 mL) was added slowly to the reaction mixture over 10 min *via* a syringe, followed by HMPA (0.88 mL, 5.1 mmol). DIBALH (1 M solution in hexanes, 3.8 mL, 3.8 mmol) was added dropwise *via* a syringe over 10 min at –30 °C, a temperature that was maintained for an additional 2 h. Alkynoate 11 (500 mg, 2.52 mmol) in toluene (12 mL) was added dropwise to the reaction mixture over 10 min *via* a syringe, and the solution was allowed to warm to –20 °C, a temperature that was maintained for 5 h. The reaction progress was monitored by ¹H NMR as the alkene byproduct S4 and alkynoate 11 have the same *R*_f by TLC. Freshly distilled 2-(chloromethyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (S5) (623 mg, 3.53 mmol) in toluene (6 mL) was added dropwise *via* a syringe over 5 min at –20 °C. The reaction was allowed to warm to rt and maintained for 16 h. The reaction was diluted with diethyl ether (10 mL), and 1 N hydrochloric acid (2 mL) was added dropwise over 5 min. The layers were separated, and the organic layer was washed sequentially with 1 N hydrochloric acid (3 × 3 mL), saturated aqueous sodium bicarbonate (1 × 5 mL), water (2 × 5 mL), and brine (1 × 10 mL), dried over magnesium sulfate, gravity filtered, concentrated by rotary evapo-

ration, and passed through a plug of silica gel eluting with 20% diethyl ether in hexanes to yield allylboronate **9** (755 mg, 88% yield) as a pale yellow oil as a 2:1 ratio of Z/E isomers. The Z/E ratio was determined by integration of the alkene resonances at 5.92 ppm for the Z-isomer and 6.71 for the E-isomer. The allylboronate **9** was contaminated with about 5% of alkene byproducts **S4** (2:1 Z/E), as determined by integration of the alkene resonances at 6.25 (dt) and 5.75 (dt) ppm for **9Z/S4** and 6.96 (dt) and 5.90 (dt) for **9E/S4**. Attempts to separate allylboronate **9** from **S4** by column chromatography resulted in a greatly reduced yield of allylboronate. Run 2: Alkynoate **11** (3.02 g, 15.1 mmol), chloromethylpinacolboronate **S5** (3.74 g, 21.2 mmol), copper iodide (288 mg, 1.51 mmol), methyl lithium (1.5 M in diethyl ether, 1.1 mL, 1.5 mmol), DIBALH (1 M in hexanes, 23 mL, 23 mmol), HMPA (5.3 mL, 30 mmol), toluene (216 mL), and THF (50 mL) provided allylboronate **9** (3.73 g, 73% yield, 2:1 Z/E containing ~10% of the alkene byproduct **S4**) as a pale yellow oil. ¹H NMR (400 MHz, CDCl₃): δ 5.92 (t, *J* = 7.6 Hz, 1H, =CH), 3.92–3.88 (m, 4H, OCH₂CH₂O), 3.67 (s, 3H, OCH₃), 2.56 (app q, *J* = 7.6 Hz, 2H, CH₂), 1.82 (br s, 2H, CH₂B), 1.77–1.71 (m, 2H, CH₂), 1.30 (s, 3H, CH₃), 1.20 (s, 12H, (CH₃)₂CC(CH₃)₂); E-isomer, where distinguishable: δ 6.71 (t, *J* = 7.6 Hz, 1H, =CH), 2.25–2.19 (m, 2H, CH₂), 1.85 (br s, 2H, CH₂B); ¹³C NMR (100 MHz, CDCl₃): δ 168.2 (C=O), 142.9, 140.6, 129.3, 128.0, 109.9, 83.4, 64.8 (2C), 51.2, 38.6, 24.8, 24.0 (4C) ppm; E-isomer, where distinguishable: δ 168.7 (C=O), 142.9, 141.0, 129.2, 128.0, 109.6, 83.3, 64.8 (2C), 51.7, 37.8, 24.7, 23.7 (4C) ppm; IR (thin film): 2981, 2884, 2240, 1716, 1645, 1437, 1351, 1257, 1145, 1055 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₁₇H₃₀BO₆, 341.2130; found, 341.2118; TLC R_f = 0.31 (25% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde stain.

3-Phenyl-2-propynal (12). Aldehyde **12** was prepared as previously described, and the spectral data matched that reported.⁶⁵ Active γ-manganese dioxide was prepared from manganese sulfate and potassium permanganate according to the previously reported method.⁶⁶ Alternatively, aldehyde **12** was also prepared according to the procedure used for the preparation of 3-triisopropylsilyl-2-propynal (**13**).⁶⁷ ¹H NMR (400 MHz, CDCl₃): δ 9.43 (s, 1H, CHO), 7.62–7.60 (m, 2H, Ph-H), 7.51–7.47 (m, 1H, Ph-H), 7.43–7.39 (m, 2H, Ph-H); ¹³C NMR (100 MHz, CDCl₃): δ 177.0 (C=O), 133.4 (2C), 131.4, 128.9 (2C), 119.6, 95.3, 88.6.

3-Triisopropylsilyl-2-propynal. Aldehyde **13** was prepared as previously described, and the spectral data matched that reported.⁶⁷ ¹H NMR (400 MHz, CDCl₃): δ 9.2 (s, 1H, CHO), 1.12–1.07 (m, 21H, Si(CH(CH₃)₂)₃); ¹³C NMR (100 MHz, CDCl₃): δ 176.8 (C=O), 104.6, 101.0, 18.6, 11.1.

General Procedure A. α-Methylene-γ-lactam Formation from the Allylboration/Lactamization Sequence Using 2-Propynals. This procedure was modified from the procedure reported by Hall and Elford.²⁸ Run 1: A 5 mL, single-necked, round-bottomed flask equipped with a magnetic stir bar, septum, and nitrogen inlet needle was charged with 3-phenyl-2-propynal (**12**) (72 mg, 0.55 mmol) and ethanol (1 mL). Ammonium hydroxide (28–30% ammonia in water, 0.74 mL, 5.5 mmol) was added *via* a syringe in a single portion at rt. The nitrogen inlet needle was removed, and the reaction was maintained for 20 min at rt. Allylboronate **9** (171 mg, 0.50 mmol, 2:1 Z/E) in ethanol (1 mL) was added dropwise over 1 min, and the reaction was stirred in the sealed flask for 5 h at rt. At this time, complete consumption of allylboronate was indicated by TLC, then 1 N HCl (~5 mL) was slowly added to afford a solution with a final pH of 1.5–2 (pH paper). The resulting solution was transferred to a separatory funnel and extracted with diethyl ether (4 × 10 mL). The combined organic layers were washed with brine (10 mL), dried over magnesium sulfate, gravity filtered, concentrated by rotary evaporation, and purified by silica gel flash column chromatography eluting with 30–50% ethyl acetate in hexanes to give lactams **14a** (trans) and **14b** (cis) (96 mg, 59%) in a 4:1 ratio. The *trans* lactam was taken on to the next step as a single isomer or as a mixture. This reaction was performed using an isomeric mixture of allylboronate **9** and alkene **S4** without significant effects on the yield; however, purification was difficult when the alkene comprises >20% of the molar ratio. This

reaction was repeated eleven times with the yields ranging from 45 to 86% with an average yield of 58%. The *trans/cis* lactam ratio ranged from 2:1 to 5:1 even though the allylboronate Z/E ratio was ~2:1. Run 2: Allylboronate **9** (1.66 g, 4.86 mmol, 2:1 Z/E), 3-phenyl-2-propynal (**12**) (696 mg, 5.35 mmol), ammonium hydroxide (28–30% ammonia in water, 7.2 mL, 54.0 mmol), and ethanol (20 mL) provided lactam **14a,b** (1.38 g, 86% yield, 5:1 *trans/cis*) as a brown oil after column chromatography. Run 3: Allylboronate **9** (1.48 g, 4.35 mmol, 2:1 Z/E), 3-phenyl-2-propynal (**12**) (623 mg, 4.79 mmol), ammonium hydroxide (28–30% ammonia in water, 6.7 mL, 48.0 mmol), and ethanol (18 mL) provided *trans* lactam **14a** (349 mg, 24% yield), a mixture of lactams **14a,b** (380 mg, 27% yield, 3:1 *trans/cis*), and a mixture of lactams **14b,a** (75 mg, 5% yield, 2:1 *cis/trans*) after column chromatography.

(4*S**,5*S**)-4-(2-(2-Methyl-1,3-dioxolan-2-yl)ethyl)-3-methylene-5-(phenylethynyl)pyrrolidin-2-one (**14a**). ¹H NMR (300 MHz, CDCl₃): δ 7.40–7.31 (m, 2H, Ph-H), 7.30–7.27 (m, 3H, Ph-H), 7.04 (br s, 1H, NH), 6.10 (d, *J* = 2.8 Hz, 1H, =CH), 5.40 (d, *J* = 1.6 Hz, 1H, =CH), 4.24 (d, *J* = 4.0 Hz, 1H, CH), 3.98–3.89 (m, 4H, OCH₂CH₂O), 3.11–3.09 (m, 1H, CH), 1.88–1.71 (m, 4H, CH₂CH₂), 1.33 (s, 3H, CH₃) and some baseline impurities in spectra; ¹³C NMR (100 MHz, CDCl₃): δ 169.7 (C=O), 142.2, 131.8 (2C), 128.7, 128.4 (2C), 122.3, 117.3, 109.7, 87.8, 84.5, 64.8 (2C), 48.8, 46.8, 35.8, 28.2, 24.1; IR (thin film): 2983, 1703, 1659, 1491, 1324, 1063 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₁₉H₂₂NO₃, 312.1594; found, 312.1585; TLC R_f = 0.17 (50% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde stain.

(4*R**,5*S**)-4-(2-(2-Methyl-1,3-dioxolan-2-yl)ethyl)-3-methylene-5-(phenylethynyl)pyrrolidin-2-one (**14b**). ¹H NMR (400 MHz, CDCl₃): δ 7.41–7.38 (m, 3H, Ph-H), 7.31–7.28 (m, 2H, Ph-H), 6.31 (br s, 1H, NH), 6.11 (d, *J* = 2.4 Hz, 1H, =CH), 5.39 (d, *J* = 2.4 Hz, 1H, =CH), 4.67 (d, *J* = 8.0 Hz, 1H, CH), 3.93–3.87 (m, 4H, OCH₂CH₂O), 3.12–3.08 (m, 1H, CH), 1.94–1.84 (m, 2H, CH₂), 1.76–1.66 (m, 2H, CH₂), 1.33 (s, 3H, CH₃), spectra were obtained as a mixture of isomers and baseline impurities were present in the ¹H NMR. ¹³C NMR (100 MHz, CDCl₃): δ 170.3 (C=O), 141.8, 131.8 (2C), 129.0, 128.5 (2C), 122.3, 117.1, 109.9, 86.7, 85.0, 64.8, 64.7, 47.7, 42.7, 36.4, 24.7, 24.1; IR (thin film): 2983, 2245, 1704, 1659, 1490, 1321, 1269, 1062 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₁₉H₂₂NO₃, 312.1594; found, 312.1586; TLC R_f = 0.32 (50% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

14b was prepared according to General Procedure A. Run 1: 3-Triisopropylsilyl-2-propynal (**13**) (340 mg, 1.62 mmol), ethanol (6 mL), ammonium hydroxide (28–30% ammonia in water, 2.2 mL, 16.0 mmol), and allylboronate **9** (500 mg, 1.47 mmol, 2:1 Z/E, contained 15% **S4**) were stirred for 5 h. HCl (2 N, 10 mL) was added to the solution, then transferred to a separatory funnel, and extracted with diethyl ether (3 × 20 mL). Column chromatography (gradient elution with 10–40% ethyl acetate in hexanes) provided *trans* lactam **15a** (126 mg, 18% yield), *cis* lactam **15b** (23 mg, 3% yield, contaminated with 25% *trans* isomer), and a 2.3:1 *trans* (**15a**):*cis* (**15b**) mixture (217 mg, 31% yield) as yellow oils in a 52% overall yield and a ratio of 3.5:1 *trans/cis* lactams. The *trans* and *cis* isomers were separated for characterization purposes but were taken on as a mixture to the next step. Run 2: Allylboronate **9** (1.67 g, 4.86 mmol, 2:1 Z/E), 3-triisopropylsilyl-2-propynal (**13**) (1.13 g, 5.35 mmol), ammonium hydroxide (28–30% ammonia in water, 7.2 mL), and ethanol (20 mL) provided lactams **15a,b** (1.72 g, 90% yield, 2:1 *trans/cis*) as a brown oil after column chromatography.

(4*S**,5*S**)-4-(2-(2-Methyl-1,3-dioxolan-2-yl)ethyl)-3-methylene-5-((triisopropylsilyl)-ethynyl)pyrrolidin-2-one (**15a**). ¹H NMR (400 MHz, CDCl₃): δ 6.06 (d, *J* = 2.6 Hz, 1H, =CH), 6.02 (br s, 1H, NH), 5.37 (d, *J* = 2.6 Hz, 1H, =CH), 4.02 (d, *J* = 5.2 Hz, 1H, CH), 3.98–3.87 (m, 4H, OCH₂CH₂O), 3.01–2.97 (m, 1H, CH), 1.91–1.63 (m, 4H, CH₂CH₂), 1.31 (s, 3H, CH₃), 1.05 (br s, 21H, Si(CH(CH₃)₂)₃); ¹³C NMR (100 MHz, CDCl₃): δ 169.3 (C=O), 142.0, 117.0, 109.6, 106.0, 86.2, 64.9 (2C), 48.9, 47.4, 36.2, 27.7, 25.0, 18.7 (6C), 11.2 (3C); IR (thin film): 2943, 2175, 1708, 1660, 1462, 1376, 1325, 1150, 1063; HRMS (TOF MS ES+) *m/z*: [M + H]⁺

calcd for $C_{22}H_{38}NO_3Si$, 392.2616; found, 392.2616; TLC R_f = 0.30 (50% EtOAc/hexanes), visualized with UV and $KMnO_4$.

(4R*,5S*)-4-(2-(2-Methyl-1,3-dioxolan-2-yl)ethyl)-3-methylene-5-((triisopropylsilyl)-ethynyl)pyrrolidin-2-one (15b). 1H NMR (300 MHz, $CDCl_3$): δ 6.23 (br s, 1H, NH), 6.05 (d, J = 2.0 Hz, 1H, =CH), 5.34 (d, J = 2.0 Hz, 1H, =CH), 4.45 (d, J = 7.6 Hz, 1H, CH), 3.95–3.79 (m, 4H, OCH_2CH_2O), 3.00–2.97 (m, 1H, CH), 1.90–1.70 (m, 4H, CH_2CH_2), 1.31 (s, 3H, CH_3), 1.02 (br s, 21H, Si($CH(CH_3)_2$)₃); ^{13}C NMR (100 MHz, $CDCl_3$): δ 170.3 (C=O), 142.1, 116.8, 103.2, 109.9, 88.4, 64.8, 64.7, 47.9, 42.8, 36.4, 24.3, 23.9, 18.7 (6C), 11.2 (3C); IR (thin film): 3213, 2866, 2175, 1660, 1377, 1221, 1143, 1112 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[M + H]^+$ calcd for $C_{22}H_{38}NO_3Si$, 392.2616; found, 392.2616; TLC R_f = 0.21 (50% EtOAc/hexanes), visualized with UV and p -anisaldehyde.

(4S*,5S*)-1-Methyl-4-(2-(2-methyl-1,3-dioxolan-2-yl)ethyl)-3-methylene-5-(phenylethynyl)pyrrolidin-2-one (S8). Run 1: A flame-dried, 10 mL, single-necked, round-bottomed flask equipped with a magnetic stir bar, septum, and nitrogen inlet needle was charged with **14a,b** (337 mg, 1.03 mmol, ~10:1 trans/cis) dissolved in dimethylformamide (5.2 mL) and cooled to 0 °C in an ice bath. Sodium hydride (60% in mineral oil, 62 mg, 1.5 mmol) was added in a single portion, and the reaction was stirred 15 min at 0 °C. Iodomethane (0.13 mL, 2.1 mmol) was added dropwise over 1 min, and the reaction was stirred at 0 °C for 15 min before removing the ice bath and allowing the reaction to warm to rt for 15 min. Upon disappearance of the starting material, the reaction mixture was poured into a separatory funnel containing saturated aqueous ammonium chloride solution (20 mL). The mixture was extracted with diethyl ether (3 $\times$ 25 mL). The organic layers were combined, washed with brine (10 mL), dried over magnesium sulfate, gravity filtered, concentrated by rotary evaporation, and purified by filtration through a plug of silica gel (elution with 50% ethyl acetate and hexanes) to provide the N -methyl lactam **S8** (225 mg, 67% yield, contains 6% of the cis isomer and some other baseline impurities) as a yellow oil. This reaction was repeated three times with the yields ranging from 67 to 74%. Run 2: Ketal **14a,b** (430 mg, 1.3 mmol, 4:1 trans/cis), sodium hydride (60% in mineral oil, 105 mg, 2.6 mmol), iodomethane (0.21 mL, 3.3 mmol), and DMF (7 mL) provided ketal **S8** (321 mg, 71% yield, 4:1 trans/cis) as a yellow oil after column chromatography. 1H NMR (400 MHz, $CDCl_3$): δ 7.42–7.40 (m, 2H, Ph-H), 7.34–7.31 (m, 3H, Ph-H), 6.08 (d, J = 2.4 Hz, 1H, =CH), 5.36 (d, J = 2.4 Hz, 1H, =CH), 4.12 (d, J = 4.0 Hz, 1H, CH), 3.99–3.90 (m, 4H, OCH_2CH_2O), 3.05–3.04 (m, 1H, CH), 3.03 (s, 3H, NCH_3), 1.87–1.69 (m, 4H, CH_2CH_2), 1.33 (s, 3H, CH_3); ^{13}C NMR (100 MHz, $CDCl_3$): δ 167.2, 142.3, 131.8 (2C), 128.9, 128.5 (2C), 122.2, 116.4, 109.7, 86.2, 85.7, 64.9 (2C), 55.4, 44.6, 35.9, 29.8, 28.5, 24.1; IR (thin film): 2925, 1697, 1660, 1427, 1396, 1145, 1082 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[M + H]^+$ calcd for $C_{20}H_{24}NO_3$, 326.1751; found, 326.1752; TLC R_f = 0.39 (75% EtOAc/hexanes), visualized with UV and p -anisaldehyde.

General Procedure B. Methyl Ketone Formation via Hydrolysis of the Ketal-Protecting Group. Run 1: A 10 mL, single-necked, round-bottomed flask equipped with a magnetic stir bar was charged with ketal **14a,b** (100 mg, 0.304 mmol, 5:1 trans/cis), acetone (4 mL), and water (0.3 mL). Pyridinium PPTS (38 mg, 0.152 mmol) was added in a single portion, a reflux condenser capped with a septum and nitrogen inlet needle was attached, and the reaction was refluxed for 16 h (oil bath temperature 70 °C). Upon completion of the reaction as observed by TLC, the reaction was allowed to cool to rt, diluted with ethyl acetate (20 mL), transferred to a separatory funnel, washed consecutively with water (2 $\times$ 5 mL) and brine (1 $\times$ 5 mL), dried over magnesium sulfate, gravity filtered, concentrated by rotary evaporation, and then eluted through a plug of silica gel with 75% ethyl acetate in hexanes to yield ketone **18a,b** (63 mg, 73% yield, 5:1 trans/cis) as a light yellow oil. For larger-scale reactions, the reaction was concentrated by rotary evaporation prior to dilution with ethyl acetate. The ratios of **18a:18b** were based upon integrated values of *trans*-lactam methine (**18a**, CHNH at 4.21 ppm) and *cis*-lactam methine (**18b**, CHNH at 4.68 ppm). This reaction was repeated nine times with yields ranging from 71 to 91% with an

average yield of 83%. The *trans* and *cis* isomers were generally taken on as a mixture but separated for characterization purposes. This reaction was performed successfully on gram scale. Run 2: Ketal **14a,b** (1.66 g, 5.33 mmol, 3:1 *trans/cis*), pyridinium PPTS (670 mg, 2.7 mmol), acetone (72 mL), and water (4 mL) provided ketone **18a,b** (1.3 g, 91% yield, 3:1 *trans/cis*) as a pale yellow oil after column chromatography.

(4S*,5S*)-3-Methylene-4-(3-oxobutyl)-5-(phenylethynyl)-pyrrolidin-2-one (18a). 1H NMR (300 MHz, $CDCl_3$): δ 7.38–7.37 (m, 2H, Ph-H), 7.32–7.29 (m, 3H, Ph-H), 6.77 (br s, 1H, NH), 6.11 (d, J = 2.4 Hz, 1H, =CH), 5.40 (d, J = 2.4 Hz, 1H, =CH), 4.21 (d, J = 4.8 Hz, 1H, CH), 3.16–3.08 (m, 1H, CH), 2.64 (t, J = 7.8 Hz, 2H, CH_2), 2.14 (s, 3H, CH_3), 2.14–2.13 (m, 1H), 1.92–1.80 (m, 1H), 16% *cis*-isomer **18b**, where distinguishable: δ 4.68 (d, J = 7.5 Hz, 1H, CH); ^{13}C NMR (100 MHz, $CDCl_3$): δ 207.5, 169.5, 141.8, 131.8 (2C), 128.9, 128.5 (2C), 122.1, 117.4, 87.4, 84.9, 48.8, 46.1, 39.9, 30.3, 27.0; IR (thin film): 3434, 2088, 1643 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[M + H]^+$ calcd for $C_{17}H_{18}NO_2$, 268.1344; found, 268.1332; TLC R_f = 0.32 (75% EtOAc/hexanes), visualized with UV and p -anisaldehyde.

(4R*,5S*)-3-Methylene-4-(3-oxobutyl)-5-(phenylethynyl)-pyrrolidin-2-one (18b). **18b** was prepared according to General Procedure B. Ketal **14a,b** (426 mg, 1.30 mmol, 3:1 *trans/cis*), pyridinium PPTS (163 mg, 0.648 mmol), acetone (18 mL), and deionized water (1 mL) provided ketone **18a,b** (307 mg, 88% yield, 3:1 *trans/cis*) as a clear oil after filtration through a silica gel plug (elution with 50% ethyl acetate in hexanes). Repurification of the mixture by column chromatography afforded fractions with predominantly the *cis* isomer, which were used for characterization purposes. 1H NMR (300 MHz, $CDCl_3$): δ 7.39–7.37 (m, 2H, Ph-H), 7.34–7.30 (m, 3H, Ph-H), 6.39 (br s, 1H, NH), 6.13 (d, J = 2.3 Hz, 1H, =CH), 5.42 (d, J = 2.3 Hz, 1H, =CH), 4.70 (d, J = 7.5 Hz, 1H, CH), 3.17–3.13 (m, 1H, CH), 2.72–2.58 (m, 2H, CH_2), 2.16 (s, 3H, CH_3), 2.15–2.09 (m, 2H, CH_2), 10% *trans*-isomer **18a**, where distinguishable: δ 4.21 (d, J = 4.8 Hz, 1H, CH); ^{13}C NMR (100 MHz, $CDCl_3$): δ 207.9, 169.9, 141.5, 131.9 (2C), 129.0, 128.6 (2C), 122.0, 117.6, 87.2, 84.6, 47.5, 41.7, 40.5, 30.2, 24.5; IR (thin film): 3245, 2927, 1708, 1657, 1418, 1360, 1273, 1166 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[M + H]^+$ calcd for $C_{17}H_{18}NO_2$, 268.1332; found, 268.1327; TLC R_f = 0.28 (75% EtOAc/hexanes), visualized with UV and p -anisaldehyde.

(4S*,5S*)-1-Methyl-3-methylene-4-(3-oxobutyl)-5-(phenylethynyl)pyrrolidin-2-one (27a). **27a** was prepared according to General Procedure B. Run 1: *Trans*-ketal **S8** (290 mg, 0.882 mmol, contained 6% *cis* isomer), pyridinium PPTS (111 mg, 0.441 mmol), acetone (12 mL), and water (0.6 mL) provided the ketone **27a** (164 mg, 66%) as a yellow oil after column chromatography. Run 2: Ketal **S8** (490 mg, 1.4 mmol, 5:1 *trans/cis*), PPTS (180 mg, 0.72 mmol), acetone (19 mL), and water (1 mL) provided *trans* ketone **27a** (310 mg, 73% yield, 19:1 *trans/cis*) and *cis* ketone (18 mg, 4% yield, 7:1 *cis/trans*) each as clear oils after column chromatography. 1H NMR (500 MHz, $CDCl_3$): δ 7.42–7.40 (m, 2H, Ph-H), 7.35–7.32 (m, 3H, Ph-H), 6.11 (d, J = 2.5 Hz, 1H, =CH), 5.36 (d, J = 2.5 Hz, 1H, =CH), 4.08 (d, J = 4.5 Hz, 1H, CH), 3.08–3.04 (m, 1H, CH), 3.03 (s, 3H, NCH_3), 2.66–2.63 (app t, J = 7.5 Hz, 2H, CH_2), 2.17 (s, 3H, CH_3), 2.12–2.04 (m, 1H), 1.90–1.84 (m, 1H); ^{13}C NMR (125 MHz, $CDCl_3$): δ 207.5, 167.1, 142.0, 131.9 (2C), 129.0, 128.6 (2C), 122.1, 116.5, 86.1, 85.9, 55.4, 44.1, 40.1, 30.3, 28.5, 27.3; IR (thin film): 2924, 1696, 1424, 1396, 1265 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[M + H]^+$ calcd for $C_{18}H_{20}NO_2$, 282.1489; found, 282.1489; TLC R_f = 0.48 (50% EtOAc/hexanes), visualized with UV and p -anisaldehyde.

(4S*,5S*)-3-Methylene-4-(3-oxobutyl)-5-((triisopropylsilyl)-ethynyl)pyrrolidin-2-one (19a). **19a** was prepared according to General Procedure B. Run 1: Ketal **15a** (120 mg, 0.306 mmol, pure *trans*), pyridinium PPTS, 39 mg, 0.15 mmol, acetone (4.1 mL), and water (0.3 mL) provided ketone **19a** (74 mg, 70% yield) as a pale yellow oil after filtration through a plug of silica (elution with 50% ethyl acetate in hexanes). This reaction was repeated six times with the yields ranging from 58 to 85% with an average yield of 73%. The *cis* and *trans* isomers were taken on to the next step as a mixture but

were separated for characterization purposes. Run 2: Ketal **15a,b** (1.74 g, 4.4 mmol, 2:1 trans/cis), PPTS (552 mg, 2.2 mmol), acetone (60 mL), and water (3 mL) provided ketone **19a,b** (1.05 g, 68%, 2:1 trans/cis) as a pale yellow oil after column chromatography. Run 3: Ketal **15a,b** (350 mg, 0.89 mmol, 2:1 trans/cis), PPTS (112 mg, 0.45 mmol), acetone (13 mL), and water (1 mL) provided ketone **19a,b** (265 mg, 85% yield, 1.5:1 trans/cis) as a clear oil after column chromatography. ^1H NMR (400 MHz, CDCl_3): δ 6.26 (br s, 1H, NH), 6.08 (d, J = 2.6 Hz, 1H, =CH), 5.38 (dd, J = 2.6 Hz, 0.8 Hz, 1H, =CH), 4.00 (d, J = 4.8 Hz, 1H, CH), 3.03–2.97 (m, 1H, CH), 2.68–2.58 (m, 2H, CH_2), 2.16 (s, 3H, CH_3), 2.15–2.06 (m, 1H), 1.85–1.75 (m, 1H), 1.05 (br s, 21H, $\text{Si}(\text{CH}(\text{CH}_3)_2)_3$); ^{13}C NMR (100 MHz, CDCl_3): δ 207.3, 169.1, 141.6, 117.2, 105.9, 86.5, 48.9, 46.6, 40.3, 30.1, 26.9, 18.7 (6C), 11.2 (3C); IR (thin film): 2943, 2865, 2175, 1709, 1659, 1463, 1366, 1323 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{20}\text{H}_{34}\text{O}_2\text{NSi}$, 348.2359; found, 348.2373; TLC R_f = 0.48 (75% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

(4*R,5*S**)-3-Methylene-4-(3-oxobutyl)-5-((triisopropylsilyl)ethynyl)pyrrolidin-2-one (19b).** **19b** was prepared according to General Procedure B. Ketal **15a,b** (1.72 g, 4.39 mmol, 4:1 trans/cis), PPTS (552 mg, 2.20 mmol), acetone (60 mL), and deionized water (3 mL) provided a first fraction of ketone **19a,b** (1.22 g, 80% yield, 4:1 trans/cis) and a second fraction of ketone **19b,a** (30 mg, 2% yield, 5:1 cis/trans) after column chromatography (gradient elution with 25–75% ethyl acetate in hexanes) as pale yellow oils. ^1H NMR (400 MHz, CDCl_3): δ 6.48 (br s, 1H, NH), 6.07 (d, J = 2.4 Hz, 1H, =CH), 5.37 (d, J = 2.4 Hz, 1H, =CH), 4.46 (d, J = 8.0 Hz, 1H, CH), 3.04–2.99 (m, 1H, CH), 2.69–2.51 (m, 2H, CH_2), 2.14 (s, 3H, CH_3), 2.13–1.97 (m, 2H, CH_2), 1.0 (br s, 21H, $\text{Si}(\text{CH}(\text{CH}_3)_2)_3$), NMR contains ~20% of trans isomer; ^{13}C NMR (100 MHz, CDCl_3): δ 207.5, 170.0, 141.5, 117.3, 102.8, 88.8, 47.5, 41.3, 40.5, 30.0, 24.3, 18.6 (6C), 11.2 (3C); IR (thin film): 3175, 2909, 2832, 2145, 1690, 1639, 1446, 1348, 1308, 1257, 1151 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{20}\text{H}_{34}\text{O}_2\text{NSi}$, 348.2353; found, 348.2363; TLC R_f = 0.41 (75% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

General Procedure C. Propargyl Alcohol Formation via Addition of Ethynyl Magnesium Bromide to Methyl Ketone. Run 1: A flame-dried, 100 mL, single-necked, round-bottomed flask equipped with a magnetic stir bar, septum, and nitrogen inlet needle was charged with ketone **18a,b** (558 mg, 2.08 mmol, ~3:1 trans/cis) dissolved in THF (25 mL). The solution was cooled to 0 °C in an ice bath, and then ethynyl magnesium bromide (0.5 M solution in THF, 17 mL, 8.5 mmol) was added dropwise via a syringe over 15 min. The reaction was maintained at 0 °C for 3 h. Next, 1 N HCl (25 mL) was added dropwise at 0 °C. The resulting solution was transferred to a separatory funnel and extracted with diethyl ether (4 × 40 mL). The organic layers were combined, washed with brine (20 mL), dried over magnesium sulfate, gravity filtered, concentrated by rotary evaporation, and purified by silica gel flash column chromatography eluting with 40–60% ethyl acetate in hexanes to provide trans propargyl alcohol **20a** (41 mg, 7% yield), cis propargyl alcohol **20b** (1 mg, 1% yield, cis), and a 4:1 trans/cis mixture of propargyl alcohols **20a** and **20b** (385 mg, 63% yield) each as clear oils and as a 1:1 ratio of diastereomers as determined by ^{13}C NMR. The cis and trans isomers were taken on as a mixture but were separated for characterization purposes. This reaction was repeated six times with the yields ranging from 65 to 81%. The trans/cis ratio of propargyl alcohol products reflected the ratio of the ketone starting material. Run 2: Ketone **18a,b** (1.30 g, 4.9 mmol, 3:1 trans/cis), ethynyl magnesium bromide (0.5 M in THF, 38 mL, 19 mmol), and THF (60 mL) provided propargyl alcohols **20a** and **20b** (1.13 g, 78% yield, 3:1 trans/cis) as a clear oil after column chromatography. Run 3: Ketone **18a,b** (640 mg, 2.4 mmol, 4:1 trans/cis), ethynyl magnesium bromide (0.5 M in THF, 19 mL, 9.5 mmol), and THF (29 mL) provided propargyl alcohols **20a** and **20b** in two fractions (528 mg, 75% yield, 6:1 trans/cis, and 39 mg, 6% yield, 2:1 cis/trans) each as clear oils after column chromatography.

(4*S,5*S**)-4-(3-Hydroxy-3-methylpent-4-ynyl)-3-methylene-5-(phenylethynyl)pyrrolidin-2-one (20a).** ^1H NMR (500 MHz, CDCl_3): δ 7.40–7.38 (m, 2H, Ph-H), 7.31–7.27 (m, 3H, Ph-H), 6.98 (br s, 1H, NH), 6.12 (d, J = 2.0 Hz, 1H, =CH), 5.43 (d, J = 2.0 Hz, 1H, =CH), 4.26 (d, J = 4.5 Hz, 1H, CH), 3.14–3.11 (m, 1H, CH), 2.62 (br s, 1H, OH), 2.46 (s, 1H, =CH), 2.04–1.96 (m, 1H), 1.90–1.79 (m, 3H), 1.52 (s, 3H, CH_3); diethyl ether at δ 3.48 and 1.21; ^{13}C NMR (125 MHz, CDCl_3): δ 169.8, 142.0, 131.8 (2C), 128.8, 128.5 (2C), 122.2, 117.6, 117.5*, 87.7, 87.4, 84.7, 72.01, 71.97*, 67.67, 67.65*, 48.9, 48.8*, 46.80, 46.78*, 39.9, 39.8*, 30.3, 30.2*, 28.9, 28.8*, *Discernible signals for one of two diastereomers at δ 65.9; IR (thin film): 3448, 1652, 1156 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{19}\text{H}_{20}\text{NO}_2$, 294.1489; found, 294.1477; TLC R_f = 0.36 (75% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

(4*R,5*S**)-4-(3-Hydroxy-3-methylpent-4-yn-1-yl)-3-methylene-5-(phenylethynyl)pyrrolidin-2-one (20b).** ^1H NMR (500 MHz, CDCl_3): δ 7.41–7.39 (m, 2H, Ph-H), 7.33–7.27 (m, 3H, Ph-H), 6.19 (br s, 1H, NH), 6.13 (d, J = 2.3 Hz, 1H, =CH), 5.44 (d, J = 2.3 Hz, 1H, =CH), 5.42 (d, J = 2.0 Hz, 1H, =CH)*, 4.70 (d, J = 7.5 Hz, 1H, CH), 4.69 (d, J = 8.0 Hz, 1H, CH)*, 3.16–3.13 (m, 1H, CH), 2.36 (s, 1H, =CH), 2.16 (s, 1H, =CH)*, 2.14–2.06 (m, 2H), 1.95–1.90 (m, 1H), 1.84–1.78 (m, 1H), 1.53 (s, 3H, CH_3); ^{13}C NMR (125 MHz, CDCl_3): δ 170.2, 141.7, 131.9 (2C), 131.85*(2C), 129.0, 128.9*, 128.6 (2C), 128.5*, 122.2*, 117.3, 117.2*, 87.3, 87.2*, 85.0, 84.9*, 72.1, 71.9*, 68.0, 67.8*, 47.74, 47.71*, 47.4, 46.2*, 42.74, 42.71*, 40.55, 40.51*, 30.4, 30.3*, 30.2, 30.1*, *Discernible signals for one of two diastereomers; IR: (thin film): 2929, 2231, 2145, 1686, 1660, 1490, 1435, 1400, 1276, 1082 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{19}\text{H}_{20}\text{NO}_2$, 294.1489; found, 294.1477; TLC R_f = 0.29 (75% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

(4*S,5*S**)-4-(3-Hydroxy-3-methylpent-4-ynyl)-1-methyl-3-methylene-5-(phenylethynyl)pyrrolidin-2-one (28a).** **28a** was prepared according to General Procedure C. Ketone **27a** (310 mg, 1.04 mmol, trans/cis 7:1), THF (14 mL), and ethynyl magnesium bromide (0.5 M solution in THF, 6.3 mL, 3.1 mmol) provided propargyl alcohol **28a** (198 mg, 62% yield, trans/cis 7:1, 1:1 ratio of diastereomers) as a white, sticky solid after column chromatography (gradient elution with 20–60% ethyl acetate in hexanes). The diastereomeric ratio was based on the ^{13}C NMR, as the diastereomers were indistinguishable by ^1H NMR. ^1H NMR (400 MHz, CDCl_3): δ 7.42–7.39 (m, 2H, Ph-H), 7.34–7.31 (m, 3H, Ph-H), 6.09 (d, J = 2.6 Hz, 1H, =CH), 5.38 (d, J = 2.6 Hz, 1H, =CH), 4.13 (d, J = 4.0 Hz, 1H, CH), 3.07–3.05 (m, 1H, CH), 3.03 (s, 3H, NCH_3), 2.47 (s, 1H, =CH), 2.28 (br s, 1H, OH), 2.06–1.95 (m, 1H), 1.87–1.78 (m, 3H), 1.53 (s, 3H, CH_3); NMR contains ~7% of the ketone starting material; ^{13}C NMR (100 MHz, CDCl_3): δ 167.2, 142.14, 142.12*, 131.9 (2C), 128.9, 128.5 (2C), 122.1, 116.64, 116.66*, 87.2, 86.1, 85.9, 72.13, 72.09*, 67.8, 67.7*, 55.4, 55.3*, 44.6, 44.5*, 39.9, 39.8*, 30.4, 30.3*, 29.2, 29.0*, 28.6, *Discernible signals for one of two diastereomers; IR (thin film): 3298, 2979, 2929, 2231, 2108, 1680, 1659, 1490, 1432, 1292, 1159, 1084 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{20}\text{H}_{22}\text{NO}_2$, 308.1651; found, 308.1658; TLC R_f = 0.60 (50% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

(4*S,5*S**)-4-(3-Hydroxy-3-methylpent-4-ynyl)-3-methylene-5-((triisopropylsilyl) ethynyl)pyrrolidin-2-one (21a).** **21a** was prepared according to General Procedure C. Run 1: Ketone **19a** (270 mg, 0.78 mmol, pure trans), ethynyl magnesium bromide (0.5 M solution in THF, 6.2 mL, 3.1 mmol), and THF (9.3 mL) yielded the propargyl alcohol **21a** (265 mg, 91% yield) as a pale yellow oil and a 1:1 mixture of diastereomers based on the ^{13}C NMR. This reaction was repeated five times with the yields ranging from 64 to 91% with an average yield of 82%. The trans/cis ratio remained the same as the starting material. ^1H NMR (500 MHz, CDCl_3): δ 6.09 (d, J = 3.0 Hz, 1H, =CH), 5.90 (br s, 1H, NH), 5.42–5.40 (m, 1H, =CH), 4.05 (d, J = 5.0 Hz, 1H, CH), 3.05–3.04 (m, 1H), 2.47 (s, 1H, =CH)*, 2.46 (s, 1H, =CH), 2.16–2.03 (m, 1H), 1.93–1.77 (m, 4H), 1.24 (s, 3H, CH_3), 1.05 (s, 21H, $\text{Si}(\text{CH}(\text{CH}_3)_2)_3$); ^{13}C NMR (125 MHz, CDCl_3): δ 169.1, 141.8, 117.13, 117.09*, 105.9, 87.2, 87.1*, 86.4, 72.1, 67.80,

67.77*, 49.0, 48.9*, 47.4, 47.3*, 30.5, 30.3*, 28.45, 28.40*, 25.0, 18.7 (6C), 11.2 (3C), *Discernible signals for one of two diastereomers; IR (thin film): 2943, 2865, 1703, 1660, 1462, 1323, 1260, 1093, 1019 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[M + H]^+$ calcd for $\text{C}_{22}\text{H}_{36}\text{NO}_2\text{Si}$, 374.2515; found, 374.2508; TLC R_f = 0.33 (50% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

(4*R*,5*S*)-4-(3-Hydroxy-3-methylpent-4-yn-1-yl)-3-methylene-5-((triisopropyl silyl)ethynyl)pyrrolidin-2-one (21b). 21b was prepared according to General Procedure C. Ketone 19a,b (1.22 g, 3.51 mmol, 4:1 trans/cis), ethynyl magnesium bromide (0.5 M solution in THF, 28 mL, 14 mmol), and THF (42 mL) provided one fraction of predominantly trans propargyl alcohol 21a (1.07 g, 82% yield, 4:1 trans/cis) and one fraction of predominantly cis propargyl alcohol 21b (52 mg, 4% yield, 2:1 cis/trans) each as a white sticky solid after column chromatography (gradient elution with 50–75% ethyl acetate in hexanes). The ratio of diastereomers was estimated to be about 1.5:1 based on the ratio of ^{13}C NMR peaks, but the peaks could not be integrated separately in the ^1H NMR. ^1H NMR (400 MHz, CDCl_3): δ 6.52 (br s, 1H, NH), 6.06 (d, J = 1.8 Hz, 1H, =CH), 5.38 (d, J = 1.8 Hz, 1H, =CH), 4.50–4.45 (m, 1H, CH), 3.04–3.00 (m, 1H, CH), 2.42 (s, 1H, $\equiv\text{CH}$), 2.41* (s, 1H, $\equiv\text{CH}$), 2.04–1.95 (m, 2H), 1.89–1.75 (m, 2H), 1.50 (s, 3H, CH_3), 1.49* (s, 3H, CH_3), 1.02 (s, 21H, Si($\text{CH}(\text{CH}_3)_2$)), NMR contains ~56% of trans isomer; ^{13}C NMR (100 MHz, CDCl_3): δ 170.3, 170.2*, 141.8, 117.2, 116.9*, 103.2, 103.0*, 88.7, 88.5*, 87.7, 87.3*, 72.0, 71.7*, 67.9, 67.7*, 48.0, 47.9*, 42.8, 42.7*, 40.4, 40.2*, 30.2, 30.0*, 25.2, 25.0*, 18.70 (6C), 18.67*, 11.2 (3C), *Discernible signals for one of two diastereomers; IR (thin film): 2943, 2246, 2172, 1703, 1658, 1544, 1462, 1321, 1172, 1072 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[M + H]^+$ calcd for $\text{C}_{22}\text{H}_{36}\text{NO}_2\text{Si}$, 308.1651; found, 308.1658; TLC R_f = 0.22 (50% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

3-Methyl-5-((2*S*,3*S*)-1-methyl-4-methylene-5-oxo-2-(phenylethynyl)pyrrolidin-3-yl)pent-1-yn-3-yl Acetate (29a). A flame-dried, 5 mL, single-necked, round-bottomed flask equipped with a stir bar, septum, and nitrogen inlet needle was charged with 4-dimethylaminopyridine (DMAP) (4 mg, 0.03 mmol) and propargyl alcohol 28a (86 mg, 0.27 mmol) dissolved in CH_2Cl_2 (1.3 mL) and cooled to 0 °C. Triethylamine (0.37 mL, 2.7 mmol) was added dropwise via a syringe over 2 min, followed by dropwise addition of acetic anhydride (0.12 mL, 1.3 mmol) over 1 min via a syringe. The ice bath was removed, and the mixture was allowed to warm to rt for 3 h, at which time the TLC showed consumption of the starting material. The mixture was diluted with CH_2Cl_2 (25 mL) and transferred to a separatory funnel. The organic layer was washed with a saturated solution of aqueous ammonium chloride (5 mL) and then brine (5 mL), dried over magnesium sulfate, gravity filtered, concentrated by rotary evaporation, and purified by flash column chromatography on a silica gel (gradient elution with 40–50% ethyl acetate in hexanes) to provide lactam 29a (50 mg, 54% yield, dr 1:1) as a clear oil. Following General Procedure D afforded a 30% yield of 29a after column chromatography.

^1H NMR (400 MHz, CDCl_3): δ 7.42–7.40 (m, 2H, Ph-H), 7.34–7.29 (m, 3H, Ph-H), 6.13 (d, J = 2.4 Hz, 1H, =CH), 5.37 (d, J = 2.4 Hz, 1H, =CH), 4.11 (d, J = 4.0 Hz, 1H, CH), 3.08–3.05 (m, 1H, CH), 3.03 (s, 3H, NCH_3), 2.58 (s, 1H, $\equiv\text{CH}$), 2.16–2.07 (m, 1H), 2.03 (s, 3H, CH_3), 1.98–1.81 (m, 3H), 1.70 (s, 3H, CH_3); ^{13}C NMR (100 MHz, CDCl_3): δ 169.42, 169.39*, 167.1, 142.0, 131.9 (2C), 129.0, 128.5 (2C), 122.1, 116.6, 116.5*, 86.0, 83.42, 83.39*, 74.51, 74.48*, 74.02, 74.00*, 55.3, 55.2*, 44.5, 44.4*, 38.23, 38.19*, 28.63, 28.57, 28.5, 26.71, 26.66*, 22.01, *Discernible signals for one of two diastereomers; IR (thin film): 3292, 2937, 2243, 2120, 1742, 1697, 1661, 1427, 1243, 1173, 1082 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[M + H]^+$ calcd for $\text{C}_{22}\text{H}_{24}\text{NO}_3$, 350.1751; found, 350.1751; TLC R_f = 0.26 (50% EtOAc/hexanes), visualized with UV and KMnO_4 .

General Procedure D. Propargyl Pivalate Formation from Propargyl Alcohol Using Scandium(III) Trifluoromethanesulfonate and Trimethyl Acetic Anhydride. Run 1: This reaction was performed on a mixture of cis and trans propargyl alcohols 20a,b or 21a,b. A flame-dried, 10 mL test tube equipped with a magnetic stir bar, septum, and nitrogen inlet needle was charged with propargyl

alcohol 20a (56 mg, 0.18 mmol, pure trans) dissolved in acetonitrile (0.72 mL). Trimethyl acetic anhydride (0.05 mL, 0.2 mmol) was added via a syringe, followed by scandium(III) trifluoromethanesulfonate (36 mg, 0.073 mmol). The reaction was stirred at rt for 16 h at which time the TLC showed consumption of the starting material. The reaction was diluted with diethyl ether (20 mL), transferred to a separatory funnel, and then washed with saturated sodium bicarbonate solution (10 mL). The aqueous layer was extracted with diethyl ether (2 $\times$ 10 mL), the organic layers were combined, washed with brine, dried over magnesium sulfate, gravity filtered, concentrated by rotary evaporation, and purified by silica gel flash column chromatography (gradient elution with 15–30% ethyl acetate in hexanes) to provide propargyl pivalate 22a (54 mg, 76%, dr 1:1) as a clear oil. Run 2: Propargyl alcohol 20a,b (368 mg, 1.25 mmol, 3.5:1 trans/cis), pivalic anhydride (0.36 mL, 1.8 mmol), scandium(III) trifluoromethanesulfonate (246 mg, 0.50 mmol), and acetonitrile (5 mL) provided trans pivalate 22a (299 mg, 61% yield) and cis pivalate 22b (79 mg, 16% yield) after column chromatography. Run 3: Propargyl alcohol 20a,b (1.13 g, 3.85 mmol, 3:1 trans/cis), pivalic anhydride (0.94 mL, 4.62 mmol), scandium(III) trifluoromethanesulfonate (758 mg, 1.54 mmol), and acetonitrile (15 mL) provided trans pivalate 22a (490 mg, 34% yield), cis pivalate 22b (163 mg, 11% yield), and a 4:1 trans/cis mixture of 22a,b (176 mg, 12% yield) each as a clear oil after column chromatography.

3-Methyl-5-((2*S*,3*S*)-4-methylene-5-oxo-2-(phenylethynyl)pyrrolidin-3-yl)pent-1-yn-3-yl Pivalate (22a). ^1H NMR (500 MHz, CDCl_3): δ 7.35–7.34 (m, 2H, Ph-H), 7.27–7.21 (m, 3H, Ph-H), 6.98 (s, 1H, NH), 6.08 (d, J = 2 Hz, 1H, =CH), 5.36 (s, 1H, =CH), 4.21 (d, J = 4.5 Hz, 1H, CH), 3.10–3.09 (m, 1H, CH), 2.50 (s, 1H, $\equiv\text{CH}$), 2.12–1.77 (m, 4H, CH_2CH_2), 1.64 (s, 3H, CH_3), 1.13 (s, 9H, $\text{C}(\text{CH}_3)_3$); ^{13}C NMR (125 MHz, CDCl_3): δ 176.6, 169.6, 142.0, 131.8 (2C), 128.8, 128.4 (2C), 122.2, 117.3, 117.2*, 87.5, 84.8, 83.52, 83.47*, 74.0, 73.9*, 73.7, 73.6*, 48.8, 48.7*, 46.7, 39.3, 38.34, 38.27*, 28.2, 27.1 (3C), 26.6, *Discernible signals for one of two diastereomers; IR (thin film): 2926, 1702, 1711, 1456, 1153 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[M + H]^+$ calcd for $\text{C}_{24}\text{H}_{28}\text{NO}_3$, 378.2064; found, 378.2049; TLC R_f = 0.47 (50% EtOAc/hexanes), visualized with UV and KMnO_4 .

3-Methyl-5-((2*S*,3*R*)-4-methylene-5-oxo-2-(phenylethynyl)pyrrolidin-3-yl)pent-1-yn-3-yl Pivalate (22b). Propargyl pivalate 22b was prepared according to General Procedure D. A mixture of propargyl alcohols 20a and 20b (140 mg, 0.48 mmol, 5:1 trans/cis), trimethyl acetic anhydride (0.12 mL, 0.57 mmol), scandium(III) trifluoromethanesulfonate (94 mg, 0.19 mmol), and acetonitrile (1.9 mL) provided trans isomer 22a (99 mg, 55% yield) as a clear oil, cis isomer 22b (20 mg, 11% yield) as a clear oil, and a 2:1 trans/cis mixture of 22a,b (7 mg, 4% yield) each as a clear oil after column chromatography (gradient elution with 15–30% ethyl acetate in hexanes). A 1.5:1 dr was determined by integration of terminal alkyne resonances in the ^1H NMR at 2.48 (major) and 2.43 ppm (minor). ^1H NMR (500 MHz, CDCl_3): δ 7.39–7.38 (m, 2H, Ph-H), 7.31–7.28 (m, 3H, Ph-H), 6.58 (br s, 1H, NH), 6.57* (br s, 1H, NH), 6.13–6.12 (m, 1H, =CH), 5.41–5.40 (m, 1H, =CH), 4.71 (d, J = 7.5 Hz, 1H, CH), 4.70* (d, J = 7.5 Hz, 1H), 3.12–3.08 (m, 1H, CH), 2.48 (s, 1H, $\equiv\text{CH}$), 2.43* (s, 1H, $\equiv\text{CH}$), 2.17–2.01 (m, 4H, CH_2CH_2), 1.68 (s, 3H, CH_3), 1.45* (s, 9H, $\text{C}(\text{CH}_3)_3$), 1.11 (s, 9H, $\text{C}(\text{CH}_3)_3$); ^{13}C NMR (125 MHz, CDCl_3): δ 176.7, 170.20, 170.15*, 141.8, 141.7*, 131.9 (2C), 128.9, 128.4 (2C), 122.2, 122.18*, 117.1, 87.03, 86.99*, 84.9, 84.8*, 83.8, 83.5*, 74.2, 73.9*, 73.7, 73.5*, 47.8, 42.70, 42.68*, 30.32, 39.30*, 39.1, 39.0*, 27.1 (3C), 26.7, 26.6*, 25.0, 24.8*, *Discernible signals for one of two diastereomers; IR (thin film): 2974, 2934, 1705, 1660, 1491, 1479, 1444, 1322, 1285, 1150, 1103 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[M + H]^+$ calcd for $\text{C}_{24}\text{H}_{28}\text{NO}_3$, 378.2064; found, 378.2068; TLC R_f = 0.37 (50% EtOAc/hexanes), visualized with UV and KMnO_4 .

3-Methyl-5-((2*S*,3*S*)-4-methylene-5-oxo-2-((triisopropylsilyl)ethynyl)pyrrolidin-3-yl)pent-1-yn-3-yl Pivalate (23a). 23a was prepared according to General Procedure D. Run 1: Propargyl alcohol 21a (100 mg, 0.27 mmol, pure trans), scandium(III) trifluoromethanesulfonate (53 mg, 0.11 mmol), pivalic anhydride

(0.07 mL, 0.35 mmol), and acetonitrile (1.1 mL) provided pivalate **23a** (118 mg, 96% yield) as a clear oil after column chromatography (gradient elution with 20–50% ethyl acetate in hexanes). ^{13}C NMR showed a 1:1 mixture of diastereomers. Run 2: Propargyl alcohol **21a,b** (230 mg, 0.62 mmol, 2.5:1 trans/cis), scandium(III) trifluoromethanesulfonate (121 mg, 0.25 mmol), pivalic anhydride (0.16 mL, 0.80 mmol), and acetonitrile (2.5 mL) provided trans pivalate **23a** (110 mg, 39% yield) and cis pivalate **23b** (44 mg, 16% yield) each as clear oils after column chromatography in a total yield of 55%. ^1H NMR (500 MHz, CDCl_3): δ 6.49 (s, 1H, NH), 6.08 (d, J = 2.6 Hz, 1H, =CH), 5.38 (d, J = 2.6 Hz, 1H, =CH), 5.37* (d, J = 2.4 Hz, 1H, =CH), 4.08–4.00 (m, 1H, CH), 3.05–3.01 (m, 1H, CH), 2.53 (s, 1H, $\equiv\text{CH}$), 2.53* (s, 1H, $\equiv\text{CH}$), 2.15–1.76 (m, 4H, CH_2CH_2), 1.67 (s, 3H, CH_3), 1.18 (s, 9H, $\text{C}(\text{CH}_3)_3$), 1.02 (s, 21H, $\text{Si}(\text{CH}(\text{CH}_3)_2)_3$); ^{13}C NMR (125 MHz, CDCl_3): δ 176.6, 169.4, 141.83, 141.81*, 117.0, 116.9*, 105.75, 105.74*, 86.4, 83.43, 83.40*, 73.9, 73.8*, 73.65, 73.60*, 49.0, 48.9*, 47.0, 39.3, 38.73*, 38.69, 38.57*, 27.9, 27.8*, 27.3 (3C), 27.2* (3C), 26.51, 26.49*, 18.7 (6C), 11.2 (3C), *Discernible signals for one of two diastereomers; IR (thin film): 2943, 2866, 1709, 1660, 1462, 1367, 1325, 1285, 1101 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{27}\text{H}_{44}\text{NO}_3\text{Si}$, 458.3090; found, 458.3059; TLC R_f = 0.37 (25% EtOAc/hexanes), visualized with UV and KMnO_4 .

3-Methyl-5-((2S*,3R*)-4-methylene-5-oxo-2-((triisopropylsilyl)ethynyl)pyrrolidin-3-yl)pent-1-yn-3-yl Pivalate (23b). Propargyl pivalate **23b** was prepared according to General Procedure D. Run 3: Propargyl alcohol **21a,b** (1.07 g, 2.87 mmol, 4:1 trans/cis), pivalic anhydride (0.70 mL, 3.5 mmol), scandium(III) trifluoromethanesulfonate (565 mg, 1.15 mmol), and acetonitrile (12 mL) provided trans lactam **23a** (565 mg, 43% yield), cis lactam **23b** (139 mg, 11% yield), and a 2:1 trans/cis mixture (72 mg, 5% yield) each as a clear oil after column chromatography (gradient elution with 5–25% ethyl acetate in hexanes). The ratios of **23a:23b** was based upon integrated values of trans-lactam methylene (**23a**, =CH at 5.38 ppm) and cis-lactam methylene (**23b**, =CH at 5.40 ppm). ^1H NMR (500 MHz, CDCl_3): δ 6.11–6.07 (m, 1H, =CH), 5.84 (br s, 1H, NH), 5.40 (d, J = 2.0 Hz, 1H, =CH), 5.38* (d, J = 1.5 Hz, 1H, =CH), 4.53–4.49 (m, 1H, CH), 3.05–3.02 (m, 1H, CH), 2.52 (s, 1H, $\equiv\text{CH}$), 2.51* (s, 1H, $\equiv\text{CH}$), 2.08–1.88 (m, 4H, CH_2CH_2), 1.68 (s, 3H, CH_3), 1.66* (s, 3H, CH_3), 1.183 (s, 9H, $\text{C}(\text{CH}_3)_3$), 1.177* (s, 9H, $\text{C}(\text{CH}_3)_3$), 1.04 (s, 21H, $\text{Si}(\text{CH}(\text{CH}_3)_2)_3$), 1.03* (s, 21H, $\text{Si}(\text{CH}(\text{CH}_3)_2)_3$); ^{13}C NMR (125 MHz, CDCl_3): δ 176.7, 169.85, 169.77*, 141.65, 141.61*, 117.3, 117.2*, 103.04, 102.99*, 89.0, 88.9*, 83.9, 83.4*, 74.3, 73.9*, 73.8, 73.4*, 47.9, 42.8, 42.6*, 39.4, 39.3*, 39.0, 38.8*, 27.20 (3C), 27.19*, 26.6, 26.3*, 25.0, 24.8*, 18.7 (6C), 11.2 (3C), *Discernible signals for one of two diastereomers; IR (thin film): 3270, 2909, 2832, 2219, 2150, 1684, 1640, 1445, 1308 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{27}\text{H}_{44}\text{NO}_3\text{Si}$, 458.3085; found, 458.3095. TLC R_f = 0.29 (25% EtOAc/hexanes), visualized with UV and KMnO_4 .

General Procedure E. 3,3-Disubstituted Allene Formation from Propargyl Pivalate Using (Triphenylphosphine)copper Hydride Hexamer. A flame-dried, 15 mL, round-bottomed flask equipped with a stir bar, septum, and nitrogen inlet needle was charged with propargyl pivalate **22a** (107 mg, 0.272 mmol), toluene (7 mL), and deionized water (0.01 mL). This solution was degassed by bubbling nitrogen through the solution for 10 min, and then the solution was cooled to -10°C with a slurry of ice and sodium chloride. (Triphenylphosphine)copper hydride hexamer (533 mg, 0.272 mmol) was weighed in a glovebox under a N_2 atmosphere into a weighing boat, removed from the glovebox, and exposed to air for ~ 1 min. The septum was removed from the flask, and the (triphenylphosphine)copper hydride hexamer was added in a single portion. The flask was evacuated and filled with nitrogen (3 $\times$) and stirred under nitrogen for 2 h. The reaction progress was monitored by removing aliquots from the reaction and measuring the disappearance of the terminal alkyne proton at 2.5 ppm by ^1H NMR. Upon completion, the mixture was poured into a cooled (0°C) solution of saturated ammonium chloride. The mixture was diluted with diethyl ether (10 mL) and stirred open to air for 30 min, then poured into a separatory funnel and extracted with diethyl ether (3 $\times$ 10 mL). The organic layers were

dried over magnesium sulfate, filtered through a plug of silica gel eluting with diethyl ether, concentrated by rotary evaporation, and then purified by silica gel flash column chromatography eluting with 15–30% ethyl acetate in hexanes to give allene **24a** (24 mg, 33%) as a clear oil.

When performing this reaction with a previously opened container of Stryker's reagent purchased from Sigma-Aldrich (90%) that was stored in the glovebox, reduction of the α -methylene group was not observed. When a newly opened bottle of Stryker's reagent purchased from Acros (97%) was used, significant amounts of the over-reduction product (**24a/31a**; 1:1) were obtained when performing the reaction at rt; however, lowering the temperature to -10°C gave better ratios (**24a/31a**; 10:1). Temperature was a major contributing factor to this product ratio as all other commercial sources and batches of Stryker's reagent (Sigma-Aldrich 90%, Acros 97%), or reagent that was freshly synthesized from copper(II) acetate, triphenylphosphine, and diphenylsilane⁶⁸ afforded little over-reduction product when the reaction was conducted at -10°C . The sources of Stryker's reagent varied in appearance: Acros (97%) was bright to dark orange depending on batch, Sigma-Aldrich (90%) was brick red to light brown depending on batch, and freshly prepared Stryker's reagent⁶⁸ was a reddish orange color similar to that purchased from Acros.

(4S*,5S*)-3-Methylene-4-(3-methylpenta-3,4-dienyl)-5-(phenylethynyl)pyrrolidin-2-one (24a). ^1H NMR (400 MHz, CDCl_3): δ 7.41–7.38 (m, 2H, Ph-H), 7.33–7.30 (m, 3H, Ph-H), 6.77 (br s, 1H, NH), 6.11 (d, J = 2.2 Hz, 1H, =CH), 5.41 (d, J = 2.2 Hz, 1H, =CH), 4.66–4.64 (m, 2H, = CH_2), 4.26 (d, J = 4.0 Hz, 1H, CH), 3.17–3.16 (m, 1H, CH), 2.15–2.08 (m, 2H), 1.92–1.86 (m, 1H), 1.81–1.74 (m, 1H), 1.71 (t, J = 3.2 Hz, 3H, CH_3); ^{13}C NMR (100 MHz, CDCl_3): δ 206.1, 169.7, 142.2, 131.8 (2C), 128.8, 128.5 (2C), 122.3, 117.3, 97.7, 87.8, 84.5, 75.3, 49.0, 46.4, 31.9, 29.9, 19.1; IR (thin film): 2922, 1741, 1736, 1445, 1372, 1270, 1216, 1110, 1042 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{19}\text{H}_{20}\text{NO}$, 278.1539; found, 278.1533; TLC R_f = 0.44 (50% EtOAc/hexanes), visualized with UV and vanillin.

(4S*,5S*)-1-Methyl-3-methylene-4-(3-methylpenta-3,4-dien-1-yl)-5-(phenylethynyl)pyrrolidin-2-one (30a). **30a** was prepared according to General Procedure E. Propargyl acetate **29a** (69 mg, 0.20 mmol), (triphenylphosphine)copper hydride hexamer (385 mg, 0.196 mmol), toluene (5.0 mL), and deionized water (0.01 mL) provided allene **30a** (30 mg, 53% yield) as a clear oil after column chromatography (gradient elution with 10–20% ethyl acetate in hexanes). Some baseline impurities from triphenylphosphine by-products of Stryker's reagent were observed in the aromatic regions of the ^1H and ^{13}C NMR spectra. ^1H NMR (500 MHz, CDCl_3): δ 7.41–7.40 (m, 2H, Ph-H), 7.34–7.30 (m, 3H, Ph-H), 6.07 (d, J = 1.5 Hz, 1H, =CH), 5.35 (s, 1H, =CH), 4.66–4.64 (m, 2H, = CH_2), 4.11 (d, J = 4.0 Hz, 1H, CH), 3.10–3.01 (m, 1H, CH), 3.03 (s, 3H, CH_3), 2.14–2.08 (m, 2H), 1.90–1.84 (m, 1H), 1.77–1.73 (m, 1H), 1.71 (t, J = 3.0 Hz, 3H, CH_3); ^{13}C NMR (125 MHz, CDCl_3): δ 201.2, 167.3, 142.4, 131.9 (2C), 128.9, 128.5 (2C), 122.2, 116.3, 97.7, 86.2, 85.7, 75.2, 55.5, 44.2, 32.2, 30.0, 28.5, 19.1; IR (thin film): 2078, 1633, 1414, 1381, 1260, 1058 cm^{-1} ; HRMS (TOF MS ES+) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{20}\text{H}_{22}\text{NO}$, 292.1701; found, 292.1712; TLC R_f = 0.66 (50% EtOAc/hexanes), visualized with UV and KMnO_4 .

(4R*,5R*)-3-Methyl-4-(3-methylpenta-3,4-dien-1-yl)-5-(phenylethynyl)pyrrolidin-2-one (31a). The structure of the reduced product **31a** was not confirmed because of our inability to separate this compound from **24a** during the column chromatography; however, two resonances in the ^1H NMR at 1.03 ppm (d, J = 7.0 Hz, 3H, CH_3) and 4.23 ppm (d, J = 4.5 Hz, 1H, CHNH) closely match that of lactams **32a** and **56**.

(4R*,5S*)-3-Methyl-4-(3-methylpenta-3,4-dien-1-yl)-5-(phenylethynyl)pyrrolidin-2-one (57). **57** was prepared according to General Procedure E. Cis lactam **22b** (75 mg, 0.19 mmol), (triphenylphosphine)copper hydride hexamer (373 mg, 0.19 mmol), toluene (5 mL), and deionized water (0.01 mL) provided allene **57** (19 mg, 36% yield) after column chromatography (gradient elution with 15–25% ethyl acetate in hexanes) as a clear oil and as the only product. ^1H NMR (500 MHz, CDCl_3): δ 7.40–7.38 (m, 2H, Ph-H),

7.33–7.31 (m, 3H, Ph-H), 5.77 (br s, 1H, NH), 4.64–4.61 (m, 2H, =CH₂), 4.51 (d, *J* = 7.0 Hz, 1H, CH), 2.27–2.14 (m, 3H), 2.03–1.98 (m, 1H), 1.92–1.84 (m, 2H), 1.71 (t, *J* = 3.0 Hz, 3H, CH₃), 1.20 (d, *J* = 7.0 Hz, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃): δ 206.3, 179.8, 131.8 (2C), 128.8, 128.5 (2C), 122.5, 97.8, 86.4, 85.3, 74.7, 47.6, 46.8, 40.3, 31.2, 28.0, 18.8, 13.8; IR (thin film): 3245, 2895, 1937, 1681, 1426, 1361 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₁₉H₂₂NO, 280.1696; found, 280.1698; TLC *R*_f = 0.33 (50% EtOAc/hexanes), visualized using UV and vanillin.

(4*S,5*S**)-3-Methylene-4-(3-methylpenta-3,4-dienyl)-5-((triisopropylsilyl)ethynyl)pyrrolidin-2-one (25a).** 25a was prepared according to General Procedure E using a reaction temperature of –20 °C. Propargyl pivalate 23a (110 mg, 0.24 mmol), (triphenylphosphine)copper hydride hexamer (424 mg, 0.22 mmol), and toluene (6 mL) yielded allenes 25a (39 mg, 46% yield) and 32a (9 mg, 11% yield) in a 4:1 ratio each as a clear oil after column chromatography (gradient elution with 5–25% ethyl acetate in hexanes). Other runs gave allene 25a in 31–53% yield along with 32a 0–31% yield. Baseline impurities in the alkenyl and aryl region were present in ¹H and ¹³C NMR. ¹H NMR (400 MHz, CDCl₃): δ 6.24 (br s, 1H, NH), 6.05 (d, *J* = 2.4 Hz, 1H, =CH), 5.36 (d, *J* = 2.4 Hz, 1H, =CH), 4.66–4.63 (m, 2H, =CH₂), 4.03 (d, *J* = 4.8 Hz, 1H, CH), 3.08–3.03 (m, 1H, CH), 2.16–2.09 (m, 1H), 2.06–1.99 (m, 1H), 1.93–1.84 (m, 1H), 1.72–1.61 (m, 1H), 1.69 (t, *J* = 3.0 Hz, 3H, CH₃), 1.04 (s, 21H, Si(CH(CH₃)₂)₃); ¹³C NMR (100 MHz, CDCl₃): δ 206.0, 169.5, 142.1, 117.0, 106.2, 97.9, 86.1, 75.4, 49.1, 47.0, 31.7, 30.1, 19.2, 18.7 (6C), 11.2 (3C); IR (thin film): 2926, 2865, 1704, 1658, 1462, 1324 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₂₂H₃₆NO_{Si}, 358.2561; found, 358.2566; TLC *R*_f = 0.32 (25% EtOAc/hexanes), visualized with UV and KMnO₄.

(4*S,5*S**)-3-Methyl-4-(3-methylpenta-3,4-dien-1-yl)-5-((triisopropylsilyl)ethynyl)pyrrolidin-2-one (32a).** The diastereomeric ratio was based upon integrated values of *trans*-lactam methine (32a, CHNH at 4.02 ppm) and *cis*-lactam methine (CHNH at 3.95 ppm). Peaks were reported for major diastereomer only. ¹H NMR (500 MHz, CDCl₃): δ 5.77 (br s, 1H, NH), 4.64–4.62 (m, 2H, =CH₂), 3.95 (d, *J* = 8.0 Hz, 1H, CH), 2.18–2.01 (m, 4H), 1.80–1.74 (m, 1H), 1.69 (t, *J* = 2.7 Hz, 3H, CH₃), 1.69–1.60 (m, 1H), 1.24 (d, *J* = 10.5 Hz, 3H, CH₃), 1.05 (s, 21H, Si(CH(CH₃)₂)₃); ¹³C NMR (125 MHz, CDCl₃): δ 206.0, 178.5, 106.3, 98.2, 85.9, 75.3, 51.5, 49.7, 42.4, 31.1, 31.0, 19.1, 18.7 (6C), 15.1, 11.3 (3C); IR (thin film): 2908, 2831, 2156, 1961, 1685, 1443 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₂₂H₃₈NO_{Si}, 360.2717; found, 360.2708; TLC *R*_f = 0.38 (35% EtOAc/hexanes), visualized with UV and vanillin.

General Procedure F. Terminal Alkyne Formation from Triisopropylsilyl-Substituted Alkyne Using TBAF. Run 1: A flame-dried, 5 mL, round-bottomed flask equipped with a stir bar, septum, and nitrogen inlet needle was charged with TIPS allene 25a (38 mg, 0.11 mmol, pure *trans* isomer) dissolved in THF (1.1 mL) and cooled to 0 °C in an ice–water bath. TBAF (1 M solution in THF, 0.15 mL, 0.15 mmol) was added dropwise *via* a syringe and the reaction was stirred for 45 min when TLC showed consumption of the starting material. Saturated aqueous ammonium chloride (2 mL) was added to the cooled reaction mixture, and the biphasic mixture was transferred to a separatory funnel. The aqueous layer was extracted with ethyl acetate (4 × 5 mL). The combined organic layers were washed with brine (1 × 5 mL), dried over magnesium sulfate, gravity filtered, concentrated by rotary evaporation, and purified by silica gel flash column chromatography (gradient elution with 25–75% ethyl acetate in hexanes) to provide allene 26 (12 mg, 57% yield) as a clear oil.

This reaction was repeated four times with yields ranging from 57 to 93% with an average yield of 79%. Run 2: TIPS allene 25a (40 mg, 0.11 mmol, *trans* isomer), TBAF (1 M solution in THF, 0.17 mL, 0.17 mmol), and THF (1.1 mL) provided allene 26 (18 mg, 82% yield) as a clear oil after column chromatography.

(4*S,5*S**)-5-Ethynyl-3-methylene-4-(3-methylpenta-3,4-dien-1-yl)pyrrolidin-2-one (26).** ¹H NMR (400 MHz, CDCl₃): δ 6.79 (br s, 1H, NH), 6.07 (d, *J* = 2.2 Hz, 1H, =CH), 5.38 (d, *J* = 2.2 Hz, 1H, =CH), 4.65–4.63 (m, 2H, =CH₂), 4.02 (dd, *J* = 2.0, 4.0 Hz, 1H, CH),

3.09–3.05 (m, 1H, CH), 2.39 (d, *J* = 2.0 Hz, 1H, =CH), 2.10–2.02 (m, 2H), 1.88–1.80 (m, 1H), 1.73–1.71 (m, 1H), 1.70 (t, *J* = 3.0 Hz, 3H, CH₃); ¹³C NMR (100 MHz, CDCl₃): δ 206.1, 169.7, 141.9, 117.4, 97.6, 82.7, 75.3, 72.9, 48.1, 46.1, 31.9, 29.8, 19.0; IR (thin film): 2094, 1936, 1633, 1411, 1308, 1061 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₁₃H₁₆NO, 202.1226; found, 202.1227; TLC *R*_f = 0.36 (50% EtOAc/hexanes), visualized using UV and *p*-anisaldehyde.

(4*S,5*S**)-5-Ethynyl-3-methyl-4-(3-methylpenta-3,4-dien-1-yl)pyrrolidin-2-one (56).** 56 was prepared according to General Procedure F. Lactam 32a (26 mg, 0.072 mmol, 9:1 dr), TBAF (1 M in THF, 0.11 mL, 0.10 mmol), and THF (0.8 mL) yielded terminal alkyne 56 (14 mg, 86% yield, 9:1 dr) as a clear oil after column chromatography (gradient elution with 25–50% ethyl acetate in hexanes). The diastereomeric ratio was based upon integrated values of *trans*-lactam methine (56, CHNH at 3.92 ppm) and *cis*-lactam methine (CHNH at 3.93 ppm). ¹H NMR (500 MHz, CDCl₃): δ 6.04 (br s, 1H, NH), 4.65–4.63 (m, 2H, =CH₂), 3.92 (dd, *J* = 7.5, 1.7 Hz, 1H, CH), 2.39 (d, *J* = 1.7 Hz, 1H, =CH), 2.17–2.01 (m, 4H), 1.79–1.68 (m, 2H), 1.70 (t, *J* = 3.2 Hz, 3H, CH₃), 1.24 (d, *J* = 7.0 Hz, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃): δ 206.1, 178.7, 98.0, 82.8, 75.3, 72.8, 50.6, 48.8, 42.2, 31.0, 30.7, 19.0, 15.3; IR (thin film): 3190, 2889, 2060, 1936, 1679, 1440, 1364 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₁₃H₁₈NO, 204.1383; found, 204.1378; TLC *R*_f = 0.24 (50% EtOAc/hexanes), visualized using UV and vanillin.

General Procedure G. *N*-Arylated Lactam Formation Using Buchwald–Hartwig Cross-Coupling Conditions. This procedure is modified from the method reported for the amidation of aryl halides.³⁸ A flame-dried, 10 mL, test tube equipped with a magnetic stir bar was charged with copper iodide (4 mg, 0.02 mmol) and cesium carbonate (50 mg, 0.15 mmol). The test tube was capped with a septum, then evacuated, and filled with nitrogen (3×). Iodobenzene (13 μL, 0.11 mmol) and *trans*-*N,N'*-dimethylcyclohexane-1,2-diamine (8 μL, 0.05 mmol) were added *via* a syringe. Lactam 24a (21 mg, 0.076 mmol) was dissolved in toluene (0.8 mL) and added to this test tube *via* a syringe. The test tube was evacuated and filled with nitrogen (3×), and the nitrogen inlet needle was removed. The test tube was then placed in an oil bath (preheated 80 °C) for 20 h at which point TLC showed a new spot. The reaction was allowed to cool to rt, filtered through a plug of silica gel eluting with 50% ethyl acetate in hexanes, concentrated, and then purified by silica gel flash column chromatography eluting with 5–15% ethyl acetate in hexanes to give lactam 37 as a clear oil (12 mg, 44% yield).

Changing the ligand to *N,N'*-dimethylethylenediamine and the base to potassium carbonate afforded a lower yield with recovery of the starting material. Lactam 24a (10 mg, 0.033 mmol), iodobenzene (5 μL, 0.04 mmol), copper iodide (5 mg, 0.03 mmol), *trans*-*N,N'*-dimethylcyclohexane-1,2-diamine (4 μL, 0.03 mmol), potassium carbonate (10 mg, 0.065 mmol), and toluene (0.4 mL) provided lactam 37 (3 mg, 27% yield, PAJ 8–6) as a clear oil after column chromatography.

(4*S,5*S**)-3-Methylene-4-(3-methylpenta-3,4-dien-1-yl)-1-phenyl-5-(phenylethynyl)pyrrolidin-2-one (37).** ¹H NMR (400 MHz, CDCl₃): δ 7.77–7.75 (m, 2H, Ph-H), 7.44–7.40 (m, 2H, Ph-H), 7.30–7.20 (m, 6H, Ph-H), 6.25 (d, *J* = 2.2 Hz, 1H, =CH), 5.51 (d, *J* = 2.2 Hz, 1H, =CH), 4.69–4.66 (m, 3H), 3.26–3.23 (m, 1H, CH), 2.18–2.13 (m, 2H), 1.91–1.83 (m, 2H), 1.73 (t, *J* = 3.2 Hz, 3H, CH₃); ¹³C NMR (100 MHz, CDCl₃): δ 206.2, 166.3, 143.0, 138.3, 131.8 (2C), 129.0 (2C), 128.8 (2C), 128.4, 125.7, 122.2 (2C), 118.4, 97.6, 87.1, 85.8, 76.8, 75.4, 55.1, 44.6, 32.5, 30.0, 19.1; IR (thin film): 2924, 2227, 1959, 1701, 1660, 1498, 1372 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₂₅H₂₄NO, 354.1852; found, 354.1837; TLC *R*_f = 0.86 (50% EtOAc/hexanes), visualized with UV and vanillin.

4-((4*S,5*S**)-3-Methylene-4-(3-methylpenta-3,4-dien-1-yl)-2-oxo-5-(phenylethynyl)pyrrolidin-1-yl)benzonitrile (38).** 38 was prepared according to General Procedure G. Lactam 24a (33 mg, 0.12 mmol), 4-iodobenzonitrile (38 mg, 0.17 mmol), copper iodide (5 mg, 0.02 mmol), *N,N'*-dimethylethylenediamine (5 μL, 0.05

mmol), cesium carbonate (78 mg, 0.24 mmol), and toluene (1.2 mL) provided lactam **38** (19 mg, 42% yield) as a pale yellow oil after column chromatography (gradient elution with 5–15% ethyl acetate in hexanes). ¹H NMR (400 MHz, CDCl₃): δ 7.99 (app dd, *J* = 2.0, 6.8 Hz, 2H, Ph-H), 7.70 (app dd, *J* = 2.0, 6.8 Hz, 2H, Ph-H), 7.33–7.28 (m, 5H, Ph-H), 6.30 (d, *J* = 2.2 Hz, 1H, =CH), 5.59 (d, *J* = 2.2 Hz, 1H, =CH), 4.71 (d, *J* = 3.2 Hz, 1H, CH), 4.68–4.66 (m, 2H, =CH₂), 3.28–3.27 (m, 1H, CH), 2.17–2.13 (m, 2H), 1.89–1.83 (m, 2H), 1.73 (t, *J* = 3.2 Hz, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃): δ 206.2, 166.5, 142.2, 133.0 (2C), 131.8 (2C), 129.2, 128.5 (2C), 121.6, 121.0 (2C), 120.0, 118.9, 108.11, 97.4, 86.5, 85.9, 75.5, 54.4, 44.3, 32.4, 29.9, 19.1; IR (thin film): 3432, 2101, 1642, 1511 cm^{−1}; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₂₆H₂₃N₂O, 379.1732; found, 379.1735; TLC *R*_f = 0.43 (25% EtOAc/hexanes), visualized with UV and vanillin.

(4*S**,5*S**)-3-Methylene-4-(3-methylpenta-3,4-dien-1-yl)-5-(phenylethynyl)-1-(4-(trifluoromethyl)phenyl)pyrrolidin-2-one (**39**). **39** was prepared according to General Procedure G. Lactam **24a** (24 mg, 0.087 mmol), 4-benzotrifluoromethyl iodide (0.02 mL, 0.1 mmol), copper iodide (2 mg, 0.02 mmol), *N,N'*-dimethylethylenediamine (4 μL, 0.04 mmol), cesium carbonate (57 mg, 0.17 mmol), and toluene (0.9 mL) provided lactam **39** (17 mg, 47%) as a clear oil after column chromatography (gradient elution with 5–10% ethyl acetate in hexanes). When potassium carbonate was used, only 14–16% of lactam **39** was isolated and 20–25% of the starting material was recovered. ¹H NMR (400 MHz, CDCl₃): δ 7.96 (d, *J* = 8.6 Hz, 2H, Ph-H), 7.67 (d, *J* = 8.6 Hz, 2H, Ph-H), 7.32–7.28 (m, 5H, Ph-H), 6.29 (d, *J* = 2.0 Hz, 1H, =CH), 5.56 (d, *J* = 2.0 Hz, 1H, =CH), 4.72 (d, *J* = 3.2 Hz, 1H, CH), 4.68–4.66 (m, 2H, =CH₂), 3.29–3.26 (m, 1H, CH), 2.19–2.13 (m, 2H), 1.90–1.83 (m, 2H), 1.73 (t, *J* = 3.0 Hz, 3H, CH₃); ¹³C NMR (100 MHz, CDCl₃): δ 206.2, 166.5, 142.5, 141.3, 131.8 (2C), 129.1 (2C), 128.5 (2C), 127.2, 126.08 (q, *J* = 4.0 Hz, 1C), 125.6, 121.9, 121.2 (2C), 119.5, 97.5, 86.3, 75.5, 54.7, 44.5, 32.5, 29.9, 19.1; IR (thin film): 2942, 2381, 1715, 1325, 1213, 1122, 1068 cm^{−1}; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₂₆H₂₃F₃NO, 422.1726; found, 422.1710; TLC *R*_f = 0.86 (50% EtOAc/hexanes), visualized with UV and vanillin.

(4*S**,5*S**)-1-(4-Methoxyphenyl)-3-methylene-4-(3-methylpenta-3,4-dien-1-yl)-5-(phenylethynyl)pyrrolidin-2-one (**40**). **40** was prepared according to General Procedure G. Lactam **24a** (29 mg, 0.11 mmol), 4-iodoanisole (37 mg, 0.16 mmol), copper iodide (4 mg, 0.02 mmol), *N,N'*-dimethylethylenediamine (5 μL, 0.04 mmol), cesium carbonate (68 mg, 0.21 mmol), and toluene (1 mL) provided lactam **40** (25 mg, 63% yield) as a pale yellow oil after column chromatography (gradient elution with 5–20% ethyl acetate in hexanes).

Performing this reaction with potassium carbonate as a base, lactam **24a** (20 mg, 0.072 mmol), 4-iodoanisole (20 mg, 0.087 mmol), copper iodide (3 mg, 0.01 mmol), *N,N'*-dimethylethylenediamine (4 μL, 0.03 mmol), potassium carbonate (20 mg, 0.14 mmol), and toluene (0.72 mL) provided lactam **40** (5 mg, 19% yield) as a pale yellow oil and starting material **24a** (5 mg, 25%) after column chromatography. ¹H NMR (500 MHz, CDCl₃): δ 7.60 (d, *J* = 8.7 Hz, 2H, Ph-H), 7.32–7.27 (m, 5H, Ph-H), 6.94 (d, *J* = 8.7 Hz, 2H, Ph-H), 6.22 (d, *J* = 1.5 Hz, 1H, =CH), 5.48 (d, *J* = 1.5 Hz, 1H, =CH), 4.68–4.66 (m, 2H, =CH₂), 4.62 (d, *J* = 3.0 Hz, 1H, CH), 3.82 (s, 3H, OCH₃), 3.27–3.21 (m, 1H, CH), 2.18–2.13 (m, 2H), 1.93–1.82 (m, 2H), 1.73 (t, *J* = 2.7 Hz, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃): δ 206.3, 166.3, 157.7, 143.1, 131.8 (2C), 131.3, 128.8, 128.4 (2C), 124.5 (2C), 122.3, 117.8, 114.3 (2C), 97.7, 87.3, 85.8, 75.3, 55.7, 55.6, 44.7, 32.5, 30.0, 19.1; IR (thin film): 2924, 1995, 1697, 1512, 1377, 1300, 1249 cm^{−1}; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₂₆H₂₆NO₂, 384.1958; found, 384.1968; TLC *R*_f = 0.59 (50% EtOAc/hexanes), visualized with UV and vanillin.

(4*S**,5*S**)-3-Methylene-4-(3-methylpenta-3,4-dien-1-yl)-5-(phenylethynyl)-1-(thiophen-2-yl)pyrrolidin-2-one (**42**). **42** was prepared according to General Procedure G. Run 1: Lactam **24a** (21 mg, 0.076 mmol), copper iodide (3 mg, 0.02 mmol), *trans*-*N,N'*-dimethylcyclohexane-1,2-diamine (5 μL, 0.03 mmol), cesium carbonate (50 mg, 0.15 mmol), 2-iodothiophene (12 μL, 0.10 mmol), and toluene (0.8

mL) provided lactam **42** (15 mg, 56% yield) as a pale yellow oil after column chromatography (5–10% ethyl acetate in hexanes gradient elution). Run 2: Lactam **24a** (23 mg, 0.083 mmol), copper iodide (3 mg, 0.02 mmol), *trans*-*N,N'*-dimethylcyclohexane-1,2-diamine (4 μL, 0.03 mmol), cesium carbonate (52 mg, 0.16 mmol), iodothiophene (10 μL, 0.11 mmol), and toluene (0.8 mL) provided lactam **42** (19 mg, 63% yield) as a pale yellow oil after column chromatography. ¹H NMR (400 MHz, CDCl₃): δ 7.38–7.36 (m, 2H, Ph-H), 7.32–7.28 (m, 3H, Ph-H), 7.03–7.02 (m, 1H, HetAr-H), 6.97–6.94 (m, 2H, HetAr-H), 6.27 (d, *J* = 2.0 Hz, 1H, =CH), 5.54 (d, *J* = 2.0 Hz, 1H, =CH), 4.70 (d, *J* = 2.8 Hz, 1H, CH), 4.68–4.66 (m, 2H, =CH₂), 3.34–3.29 (m, 1H, CH), 2.15–2.11 (m, 2H), 1.84 (m, 2H), 1.73 (t, *J* = 3.2 Hz, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃): δ 206.2, 164.3, 141.5, 139.4, 131.9 (2C), 129.0, 128.5 (2C), 124.2, 122.0, 119.3, 119.2, 113.0, 97.4, 86.1, 86.0, 75.4, 55.4, 44.9, 33.2, 29.8, 19.1; IR (thin film): 2924, 2227, 1959, 1695, 1534, 1451, 1399 cm^{−1}; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₂₃H₂₂NOS, 360.1417; found, 360.1401; TLC *R*_f = 0.58 (25% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

(4*S**,5*S**)-3-Methylene-4-(3-methylpenta-3,4-dien-1-yl)-5-(phenylethynyl)-1-tosylpyrrolidin-2-one (**43**). Run 1: A flame-dried test tube equipped with a stir bar, septum, and nitrogen inlet needle was charged with lactam **24a** (20 mg, 0.07 mmol) and DMF (0.75 mL) and cooled to 0 °C under a nitrogen atmosphere. Sodium hydride (60% dispersion in mineral oil, 6 mg, 0.14 mmol) was added in a single portion, and the reaction mixture was stirred for 15 min at 0 °C. *para*-Toluene sulfonyl chloride (28 mg, 0.14 mmol) was added in a single portion, the ice bath was removed, and the solution was allowed to warm to rt for 2.5 h. TLC analysis revealed consumption of the starting material. The reaction mixture was poured into a separatory funnel containing saturated aqueous ammonium chloride (5 mL). The aqueous layer was extracted with ethyl acetate (3 × 10 mL). The organic layers were combined, dried over magnesium sulfate, gravity filtered, concentrated by rotary evaporation, and purified by silica gel flash column chromatography eluting with 10–25% ethyl acetate in hexanes. Lactam **43** was obtained as a pale yellow oil (9 mg, 29% yield). Run 2: This reaction was repeated, but the temperature was maintained at 0 °C for 1 h after addition of *para*-toluene sulfonyl chloride and then allowed to warm to rt for only 1 h before workup. Lactam **24a** (18 mg, 0.069 mmol), sodium hydride (60% dispersion in mineral oil, 5 mg, 0.1 mmol), *para*-toluene sulfonyl chloride (25 mg, 0.13 mmol), and DMF (0.7 mL) provided lactam **43** (14 mg, 48% yield) as a yellow oil after column chromatography; unidentified contaminants are observed in the ¹H NMR. Additional purification by column chromatography provided allene-yne **43** (4 mg, 18% yield) as a clear oil. ¹H NMR (500 MHz, CDCl₃): δ 8.08 (d, *J* = 8.0 Hz, 2H, Ph-H), 7.25–7.28 (m, 5H, Ph-H), 7.22 (d, *J* = 8.0 Hz, 2H, Ph-H), 6.23 (d, *J* = 1.7 Hz, 1H, =CH), 5.54 (d, *J* = 1.7 Hz, 1H, =CH), 4.98 (d, *J* = 1.5 Hz, 1H, CH), 4.68–4.67 (m, 2H, =CH₂), 3.11–3.08 (m, 1H, CH), 2.38 (s, 3H, CH₃), 2.08–2.05 (m, 2H), 1.88–1.84 (m, 1H), 1.74–1.69 (m, 1H), 1.71 (t, *J* = 3.3 Hz, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃): δ 206.3, 164.9, 145.3, 141.1, 135.6, 131.9 (2C), 131.7, 129.8 (2C), 129.5 (2C), 128.5 (2C), 122.6, 121.9, 97.2, 86.3, 85.9, 75.4, 53.6, 45.5, 32.8, 29.8, 21.8, 19.0; IR (thin film): 2891, 2218, 1937, 1697, 1579, 1429, 1356, 1160, 1078, 804, 750 cm^{−1}; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₂₆H₂₆NO₃S, 432.1628; found, 432.1641; TLC *R*_f = 0.66 (25% EtOAc/hexanes), visualized with UV and vanillin.

tert-Butyl (4*S**,5*S**)-3-methylene-4-(3-methylpenta-3,4-dien-1-yl)-2-oxo-5-(phenylethynyl)pyrrolidine-1-carboxylate (**44**). Run 1: A 5 mL, flame-dried, test tube equipped with a stir bar, septum, and nitrogen inlet needle was charged with allene-yne **24a** (15 mg, 0.054 mmol) and CH₂Cl₂ (0.3 mL). 4-Dimethylaminopyridine (1 mg, 0.008 mmol) was added, and the solution was cooled to 0 °C. Triethylamine (0.08 mL, 0.5 mmol) was added dropwise, followed by addition of di-*tert*-butyl dicarbonate (59 mg, 0.27 mmol) in a single portion. The ice bath was removed, and the solution was allowed to warm to rt and stirred for 2 h. After consumption of the starting material was observed by TLC, the solution was diluted with CH₂Cl₂ (15 mL), transferred to a separatory funnel, and washed sequentially with

saturated aqueous ammonium chloride (5 mL) and brine (5 mL). The organic layer was dried over magnesium sulfate, gravity filtered, concentrated by rotary evaporation, and purified by silica gel flash column chromatography (gradient elution with 5–25% ethyl acetate in hexanes) to provide allene-yne **44** (8 mg, 40% yield) as a clear oil. Run 2: Lactam **24a** (23 mg, 0.083 mmol), dimethylaminopyridine (1 mg, 0.008 mmol), triethylamine (0.12 mL, 0.83 mmol), di-*tert*-butyl dicarbonate (90 mg, 0.41 mmol), and CH₂Cl₂ (0.4 mL) provided lactam **44** (17 mg, 54% yield) as a pale yellow oil after column chromatography. ¹H NMR (500 MHz, CDCl₃): δ 7.38–7.36 (m, 2H, Ph-H), 7.31–7.29 (m, 3H, Ph-H), 6.30 (d, *J* = 1.7 Hz, 1H, =CH), 5.54 (d, *J* = 1.7 Hz, 1H, =CH), 4.67–4.65 (m, 3H), 3.08–3.05 (m, 1H, CH), 2.07–2.05 (m, 2H), 1.80–1.71 (m, 2H), 1.71 (t, *J* = 3.2 Hz, 3H, CH₃), 1.57 (s, 9H, C(CH₃)₃); ¹³C NMR (125 MHz, CDCl₃): δ 206.3, 165.6, 149.9, 142.1, 131.8 (2C), 128.7, 128.5 (2C), 122.4, 121.6, 97.4, 87.5, 84.0, 83.7, 75.3, 52.8, 43.6, 32.8, 29.8, 28.2 (3C), 19.0; IR (thin film): 2969, 2915, 1793, 1759, 1717, 1401, 1302, 1153 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₂₄H₂₈NO₃, 378.2069; found, 378.2082; TLC *R*_f = 0.69 (50% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

General Procedure H. Fused 5,7,5-Ring Structure Formation Using Rh(I)-Catalyzed APKR Conditions (Slow Addition of Allene-yne to the Rhodium Catalyst). Run 1: A flame-dried, 10 mL, two-necked, round-bottomed flask equipped with a magnetic stir bar, condenser capped with a septum, and septum was charged with rhodium biscarbonyl chloride dimer ([Rh(CO)₂Cl]₂, 2 mg, 0.004 mmol) and toluene (3.4 mL). The apparatus was placed under vacuum for 2–3 s by piercing the septum with a needle attached to a Schlenk line and then filled with carbon monoxide *via* a needle attached to a balloon filled with carbon monoxide. This process was repeated three times. The reaction flask was lowered into an oil bath (preheated to 110 °C). Allene-yne **24a** (13 mg, 0.044 mmol) was dissolved in toluene (1.1 mL) and added dropwise to the stirring solution of rhodium biscarbonyl chloride dimer over 1 h using a syringe pump. After the addition was complete, the reaction was stirred for an additional 30 min at 110 °C at which time TLC showed consumption of the starting material. The flask was removed from the oil bath and allowed to cool to rt, then polymer-bound triphenylphosphine (3 mmol/g, 60 mg) was added, and the reaction was stirred (4 h). The polymer was removed by vacuum filtration, the solution was concentrated, and the residue was purified by silica gel flash column chromatography with gradient elution using 25–75% ethyl acetate in hexanes to yield lactam **45** as a white solid (9 mg, 64% yield). Run 2: Allene-yne **24a** (21 mg, 0.076 mmol), rhodium biscarbonyl chloride dimer (3 mg, 0.008 mmol), and toluene (7.6 mL) provided lactam **45** (19 mg, 79% yield) as a white solid after column chromatography. Compound **45** was crystallized by vapor diffusion using ethyl acetate and pentanes, and X-ray crystallography confirmed compound structure **45**.

General Procedure I. Fused 5,7,5-Ring Structure Formation Using Rh(I)-Catalyzed APKR Conditions. Run 3: To a flame-dried, two-necked, 25 mL, round-bottomed flask equipped with a septum, condenser capped with a septum, and magnetic stir bar were added rhodium biscarbonyl chloride dimer (4 mg, 0.01 mmol) and toluene (7 mL). The apparatus was placed under vacuum for 2–3 s by piercing the septum with a needle attached to a Schlenk line then filled with carbon monoxide *via* a needle attached to a balloon filled with carbon monoxide. This process was repeated three times. The reaction flask was lowered into an oil bath (preheated to 110 °C). Allene **24a** (30 mg, 0.11 mmol) was dissolved in toluene (4 mL) and added to the stirring solution of rhodium catalyst *via* a syringe in a single portion. Consumption of the starting material was evident by TLC after 1 h, and the reaction mixture was concentrated by rotary evaporation. The crude residue was purified by silica gel flash column chromatography eluting with a gradient of 30–75% ethyl acetate in hexanes to give lactam **45** as a white solid (25 mg, 75% yield).

(3a*S**,9b*S**)-6-Methyl-3-methylene-9-phenyl-3,3a,4,5-tetrahydro-1*H*-azuleno[4,5-*b*]pyrrole-2,8(7*H*,9*bH*)-dione (**45**). ¹H NMR (400 MHz, CDCl₃): δ 7.46–7.40 (m, 3H, Ph-H), 7.25–7.24 (m, 2H, Ph-H), 5.98 (d, *J* = 3.2 Hz, 1H, =CH), 5.58 (br s, 1H, NH),

5.27 (d, *J* = 3.2 Hz, 1H, =CH), 4.82 (d, *J* = 8.4 Hz, 1H, CH), 3.18, 3.11 (AB q, *J* = 20.0 Hz, 2H, CH₂), 3.05–3.02 (m, 1H, CH), 2.77–2.72 (m, 1H), 2.42–2.35 (m, 2H), 1.99 (s, 3H, CH₃), 1.91–1.83 (m, 1H); ¹³C NMR (100 MHz, CDCl₃): δ 202.0, 168.6, 165.5, 143.1, 140.8, 135.7, 131.8, 131.3, 129.3 (4C), 128.4, 115.3, 56.2, 44.3, 40.9, 31.8, 26.6, 24.8; IR (thin film): 3419, 2950, 2865, 1697, 1675, 1460, 1380, 1305, 1201 cm⁻¹; HRMS (TOF MS ESI+) *m/z*: [M + H]⁺ calcd for C₂₀H₂₀NO₂, 306.1489; found, 306.1479; TLC *R*_f = 0.22 (75% EtOAc/hexanes) visualized with UV and *p*-anisaldehyde.

(3a*S**,9b*S**)-6-Methyl-3-methylene-9-(triisopropylsilyl)-3,3a,4,5-tetrahydro-1*H*-azuleno[4,5-*b*]pyrrole-2,8(7*H*,9*bH*)-dione (**46**). **46** was prepared according to General Procedure H. Run 1: Allene-yne **25a** (20 mg, 0.056 mmol), rhodium dicarbonyl chloride dimer (2 mg, 0.006 mmol), and toluene (5.7 mL) provided lactam **46** (11 mg, 50% yield) as a clear oil after column chromatography (gradient elution with 15–30% ethyl acetate in hexanes). Consumption of the starting material was evident by TLC after 3 h. Run 2: Allene-yne **25a** (25 mg, 0.078 mmol), rhodium biscarbonyl chloride dimer (3 mg, 0.008 mmol), and toluene (8 mL) provided lactam **46** (11 mg, 41% yield) as a white solid after column chromatography. ¹H NMR (300 MHz, CDCl₃): δ 6.19 (br s, 1H, NH), 6.07 (d, *J* = 3.1 Hz, 1H, =CH), 5.31 (d, *J* = 3.1 Hz, 1H, =CH), 4.65 (d, *J* = 8.1 Hz, 1H, CH), 3.05–3.00 (m, 1H, CH), 3.00 (s, 2H, CH₂), 2.99–2.65 (m, 1H), 2.45–2.27 (m, 2H), 1.95 (s, 3H, CH₃), 1.90–1.80 (m, 1H), 1.63–1.53 (m, 3H), 1.13 (d, *J* = 2.0 Hz, 9H), 1.09 (d, *J* = 2.0 Hz, 9H); ¹³C NMR (100 MHz, CDCl₃): δ 208.2, 179.9, 169.3, 143.1, 137.9, 134.1, 133.9, 115.7, 56.8, 44.2, 41.4, 31.9, 27.1, 25.4, 19.5 (3C), 19.4 (3C), 13.0 (3C); IR (thin film): 2943, 2881, 1719, 1688, 1518, 1504 cm⁻¹; HRMS (TOF MS ESI+) *m/z*: [M + H]⁺ calcd for C₂₃H₃₆NO₂Si, 386.2510; found, 386.2515; TLC *R*_f = 0.37 (50% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

(3a*S**,9b*S**)-6-Methyl-3-methylene-1,3a,4,5,7,9b-hexahydro-2*H*-azuleno[4,5-*b*]pyrrole-2,8(3*H*)-dione (**47**). **47** was prepared according to General Procedure H. Run 1: Allene-yne **26** (12 mg, 0.060 mmol), rhodium biscarbonyl chloride dimer (3 mg, 0.006 mmol), and toluene (6 mL) provided **47** (9 mg, 64% yield) as a white solid after column chromatography (gradient elution with 50–100% ethyl acetate in hexanes). Run 2: Allene-yne **26** (17 mg, 0.085 mmol), rhodium dicarbonyl chloride dimer (4 mg, 0.008 mmol), and toluene (8.6 mL) provided lactam **47** (8 mg, 42% yield) as a white solid after column chromatography. ¹H NMR (500 MHz, CDCl₃): δ 7.86 (br s, 1H, NH), 6.14 (s, 1H, =CH-CO), 6.06 (d, *J* = 3.0 Hz, 1H, =CH), 5.32 (d, *J* = 3.0 Hz, 1H, =CH), 4.58 (d, *J* = 8.5 Hz, 1H, CH), 3.00 (s, 2H, CH), 2.86–2.82 (m, 1H), 2.76–2.71 (m, 1H), 2.42–2.31 (m, 2H), 1.94 (s, 3H, CH₃), 1.93–1.87 (m, 1H); ¹³C NMR (126 MHz, CDCl₃): δ 204.0, 173.0, 170.7, 143.9, 137.1, 131.6, 126.6, 115.8, 56.1, 43.4, 41.3, 32.0, 26.8, 24.9; IR (thin film): 2895, 1685, 1556, 1420, 1327, 1231, 1149 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₁₄H₁₆NO₂, 230.1179; found, 230.1176; TLC *R*_f = 0.15 (100% EtOAc), visualized with UV and *p*-anisaldehyde.

(3a*S**,9b*S**)-1,6-Dimethyl-3-methylene-9-phenyl-1,3a,4,5,7,9b-hexahydro-2*H*-azuleno[4,5-*b*]pyrrole-2,8(3*H*)-dione (**48**). **48** was prepared according to General Procedure H. Allene **30a** (30 mg, 0.1 mmol), rhodium biscarbonyl chloride dimer (4 mg, 0.009 mmol), and toluene (8.6 mL) provided lactam **48** (24 mg, 75% yield) as a white solid after column chromatography (gradient elution with 25–75% ethyl acetate in hexanes). ¹H NMR (400 MHz, CDCl₃): δ 7.33–7.31 (m, 3H, Ph-H), 7.11–7.10 (m, 2H, Ph-H), 6.10 (d, *J* = 3.2 Hz, 1H, =CH), 5.34 (d, *J* = 3.2 Hz, 1H, =CH), 4.52 (d, *J* = 8.0 Hz, 1H, CH), 3.17, 3.10 (AB q, *J* = 21.2 Hz, 2H, CH₂), 3.20–3.07 (m, 1H), 2.93–2.84 (m, 1H), 2.57–2.46 (m, 1H), 2.43–2.38 (m, 1H), 2.01 (s, 3H, CH₃), 1.99 (s, 3H, CH₃), 1.96–1.88 (m, 1H); ¹³C NMR (100 MHz, CDCl₃): δ 202.8, 168.8, 165.5, 143.2, 137.6, 136.6, 130.6, 130.3 (2C), 128.4, 127.7 (2C), 115.6, 62.1, 41.7, 40.5, 32.3, 30.4, 29.8, 28.5, 25.8; IR (thin film): 2885, 1670, 1638, 1422, 1378, 1296, 1257, 1141, 1080 cm⁻¹; HRMS (TOF MS ESI+) *m/z*: [M + H]⁺ calcd for C₂₁H₂₂NO₂, 320.1645; found, 320.1651; TLC *R*_f = 0.36 (75% EtOAc/hexanes), visualized with silica gel, UV, *p*-anisaldehyde.

(3a*S**,9b*S**)-6-Methyl-3-methylene-1,9-diphenyl-1,3a,4,5,7,9b-hexahydro-2*H*-azuleno[4,5-*b*]pyrrole-2,8(3*H*)-dione (**49**). **49** was

prepared according to General Procedure I. Run 1: Allene-yne **37** (11 mg, 0.031 mmol), rhodium biscarbonyl chloride dimer (1.2 mL of a solution of Rh(I)-catalyst in toluene, 1 mg Rh/1 mL toluene, 0.003 mmol), and toluene (3.1 mL) provided lactam **49** (8 mg, 67% yield) as a white solid after column chromatography (gradient elution with 25–50% ethyl acetate in hexanes). Run 2: Allene-yne **37** (3 mg, 0.008 mmol), rhodium biscarbonyl chloride dimer (0.5 mL of a 1 mg/mL solution in toluene), and toluene (0.5 mL) provided lactam **49** (2 mg, 66% yield) as a white solid after column chromatography. ¹H NMR (500 MHz, CDCl₃): δ 7.10–6.86 (m, 8H, Ph-H), 6.53–6.52 (m, 2H, Ph-H), 6.24 (d, *J* = 3.3 Hz, 1H, =CH), 5.46 (d, *J* = 3.3 Hz, 1H, =CH), 5.34 (d, *J* = 8.0 Hz, 1H, CH), 3.31–3.29 (m, 1H, CH), 3.17, 3.08 (AB q, *J* = 21.0 Hz, 2H, CH₂), 3.08–3.02 (m, 1H), 2.63–2.60 (m, 1H), 2.52–2.47 (m, 1H), 2.05 (s, 3H, CH₃), 2.08–2.03 (m, 1H); ¹³C NMR (126 MHz, CDCl₃): δ 202.5, 166.4, 163.3, 143.6, 139.4, 137.6, 136.4, 130.6, 130.5, 129.6 (2C), 128.1 (2C), 127.5, 127.4 (2C), 124.1, 119.3 (2C), 117.1, 59.1, 40.9, 40.6, 32.4, 28.7, 25.9; IR (thin film): 2891, 1675, 1631, 1477, 1356 cm^{−1}; HRMS (TOF MS ES⁺) *m/z*: [M + H]⁺ calcd for C₂₆H₂₄NO₂, 382.1802; found, 382.1807; TLC *R*_f = 0.35 (50% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

4-((3aS*,9bS*)-6-Methyl-3-methylene-2,8-dioxo-9-phenyl-2,3,3a,4,5,7,8,9b-octahydro-1H-azuleno[4,5-b]pyrrol-1-yl)-benzonitrile (50). **50** was prepared according to General Procedure H. Allene-yne **38** (17 mg, 0.045 mmol), rhodium biscarbonyl chloride dimer (2 mg, 0.0045 mmol), and toluene (4.5 mL) provided lactam **50** (13 mg, 72% yield) as a light yellow solid after column chromatography (gradient elution with 15–35% ethyl acetate in hexanes). ¹H NMR (400 MHz, CDCl₃): δ 7.27–7.25 (m, 2H, Ph-H), 7.13–7.03 (m, 5H, Ph-H), 6.61–6.45 (m, 2H, Ph-H), 6.30 (d, *J* = 3.4 Hz, 1H, =CH), 5.55 (d, *J* = 3.4 Hz, 1H, =CH), 5.31 (d, *J* = 8.4 Hz, 1H, CH), 3.35–3.31 (m, 1H, CH), 3.19, 3.10 (AB q, *J* = 21.0, 36.6 Hz, 2H, CH₂), 3.06–3.01 (m, 1H), 2.68–2.59 (m, 1H), 2.55–2.49 (m, 1H), 2.10–2.03 (m, 1H), 2.08 (s, 3H, CH₃); ¹³C NMR (100 MHz, CDCl₃): δ 202.1, 166.6, 162.2, 142.6, 141.4, 139.2, 137.0, 132.1 (2C), 130.3, 130.2, 129.7 (2C), 127.8, 127.5 (2C), 119.6 (2C), 119.0, 118.8, 107.0, 58.9, 40.8, 40.5, 32.2, 28.6, 25.9; IR (thin film): 2925, 2854, 2224, 1704, 1659, 1604, 1509, 1345, 1179 cm^{−1}; HRMS (TOF MS ES⁺) *m/z*: [M + H]⁺ calcd for C₂₇H₂₃N₂O₂, 407.1754; found, 407.1753; TLC *R*_f = 0.27 (50% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

(3aS*,9bS*)-6-Methyl-3-methylene-9-phenyl-1-(4-(trifluoromethyl)phenyl)-1,3a,4,5,7,9b-hexahydro-2H-azuleno[4,5-b]pyrrole-2,8(3H)-dione (51). **51** was prepared according to General Procedure H. Run 1: Allene-yne **39** (17 mg, 0.04 mmol), rhodium biscarbonyl chloride dimer (2 mg, 0.004 mmol), and toluene (4 mL) provided lactam **51** (13 mg, 72% yield) as a white solid after column chromatography (gradient elution with 15–50% ethyl acetate in hexanes). Run 2: Allene-yne **39** (6 mg, 0.01 mmol), rhodium biscarbonyl chloride dimer (0.5 mL of a 1 mg/mL solution in toluene), and toluene (0.9 mL) provided lactam **51** (4 mg, 67% yield) as a white solid after column chromatography. ¹H NMR (500 MHz, CDCl₃): δ 7.20 (d, *J* = 8.5 Hz, 2H, Ph-H), 7.10 (app t, *J* = 7.5 Hz, 1H, Ph-H), 7.00 (d, *J* = 9.0 Hz, 4H, Ph-H), 6.55–6.43 (m, 2H, Ph-H), 6.28 (d, *J* = 3.5 Hz, 1H, =CH), 5.52 (d, *J* = 3.5 Hz, 1H, =CH), 5.33 (d, *J* = 8.5 Hz, 1H, CH), 3.34–3.31 (m, 1H, CH), 3.17, 3.10 (ABq, *J* = 21.0 Hz, 2H, CH₂), 3.09–3.02 (m, 1H), 2.66–2.59 (m, 1H), 2.53–2.49 (m, 1H), 2.10–1.99 (m, 1H), 2.06 (s, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃): δ 202.3, 166.5, 162.5, 143.0, 140.4, 139.3, 136.9, 130.38, 130.36, 129.6 (2C), 127.6 (2C), 127.5 (2C), 125.8, 125.6 (q, *J* = 32.5 Hz, 1C), 125.2 (q, *J* = 3.7 Hz, 1C), 122.8, 119.3, 118.2, 59.1, 40.9, 40.5, 32.3, 28.6, 25.9; IR (thin film): 2930, 1698, 1615, 1325, 1168, 1117 cm^{−1}; HRMS (TOF MS ES⁺) *m/z*: [M + H]⁺ calcd for C₂₇H₂₃NO₂F₃, 450.1675; found, 450.1655; TLC *R*_f = 0.36 (50% EtOAc/hexanes), visualized with UV and vanillin.

(3aS*,9bS*)-1-(4-Methoxyphenyl)-6-methyl-3-methylene-9-phenyl-1,3a,4,5,7,9b-hexahydro-2H-azuleno[4,5-b]pyrrole-2,8(3H)-dione (52). **52** was prepared according to General Procedure I. Run 1: Allene-yne **40** (6 mg, 0.2 mmol), rhodium dicarbonyl chloride dimer (0.5 mL of a solution of Rh(I)-catalyst in toluene, 1 mg Rh/1

mL toluene, 0.001 mmol), and toluene (1.3 mL) provided lactam **52** (5 mg, 76%) as a white solid after column chromatography (gradient elution with 15–35% ethyl acetate in hexanes). Run 2: Allene-yne **40** (25 mg, 0.065 mmol), rhodium biscarbonyl chloride dimer (3 mg, 0.006 mmol), and toluene (6.6 mL) provided lactam **52** (20 mg, 77%) as a white solid after column chromatography. ¹H NMR (500 MHz, CDCl₃): δ 7.12–7.10 (m, 1H, Ph-H), 7.06–7.03 (m, 2H, Ph-H), 6.79–6.77 (m, 2H, Ph-H), 6.58–6.57 (m, 2H, Ph-H), 6.51–6.49 (m, 2H), 6.22 (d, *J* = 3.2 Hz, 1H, =CH), 5.45 (d, *J* = 3.2 Hz, 1H, =CH), 5.31 (d, *J* = 8.5 Hz, 1H, CH), 3.72 (s, 3H, OCH₃), 3.31–3.28 (m, 1H), 3.17, 3.08 (AB q, *J* = 21.0 Hz, 2H, CH₂), 3.09–3.01 (m, 1H), 2.62–2.58 (m, 1H), 2.51–2.46 (m, 1H), 2.10–2.00 (m, 1H), 2.05 (s, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃): δ 202.4, 166.2, 163.4, 156.3, 143.7, 139.3, 136.4, 131.2, 130.7, 129.6 (2C), 127.6, 127.4 (2C), 120.6 (2C), 116.7, 113.5 (2C), 59.2, 55.6, 41.0, 40.6, 32.4, 29.9, 28.7, 25.8; IR (thin film): 2924, 1700, 1511, 1250 cm^{−1}; HRMS (TOF MS ES⁺) *m/z*: [M + H]⁺ calcd for C₂₇H₂₆NO₃, 412.1907; found, 412.1901; TLC *R*_f = 0.23 (50% EtOAc/hexanes) [silica gel, UV, vanillin].

(3aS*,9bS*)-6-Methyl-3-methylene-9-phenyl-1-(thiophen-2-yl)-1,3a,4,5,7,9b-hexahydro-2H-azuleno[4,5-b]pyrrole-2,8(3H)-dione (53). **53** was prepared according to General Procedure I. Run 1: Allene-yne **42** (12 mg, 0.033 mmol), rhodium biscarbonyl chloride dimer (1.3 mL of a solution of Rh(I)-catalyst in toluene, 1 mg Rh/1 mL toluene, 0.003 mmol), and toluene (3.3 mL) provided lactam **53** (7 mg, 54% yield) as a pale yellow solid after column chromatography (gradient elution with 25–50% ethyl acetate in hexanes). Run 2: Allene-yne **42** (18 mg, 0.050 mmol), rhodium biscarbonyl chloride dimer (2 mg, 0.005 mmol), and toluene (5 mL) provided lactam **53** (15 mg, 79%) as a pale yellow solid after column chromatography. ¹H NMR (400 MHz, CDCl₃): δ 7.10–7.04 (m, 3H, Ph-H), 6.70 (dd, *J* = 1.4, 5.4 Hz, 1H, HetAr-H), 6.65–6.59 (m, 3H), 6.25 (d, *J* = 3.6 Hz, 1H, HetAr-H), 6.24 (d, *J* = 3.2 Hz, 1H, =CH), 5.49 (d, *J* = 3.2 Hz, 1H, =CH), 5.16 (d, *J* = 8.0 Hz, 1H, CH), 3.40–3.34 (m, 1H, CH), 3.21, 3.12 (ABq, *J* = 21.0 Hz, 2H, CH₂), 3.08–2.98 (m, 1H), 2.69–2.58 (m, 1H), 2.55–2.48 (m, 1H), 2.06 (s, 3H, CH₃), 2.03–1.99 (m, 1H); ¹³C NMR (100 MHz, CDCl₃): δ 202.4, 164.6, 163.1, 141.8, 139.5, 139.0, 136.2, 130.4, 130.1, 129.3 (2C), 127.8, 127.2 (2C), 123.0, 119.2, 117.8, 111.8, 60.8, 41.9, 40.5, 32.2, 28.5, 25.9; IR (thin film): 3434, 1693, 1575, 1360, 1305 cm^{−1}; HRMS (TOF MS ES⁺) *m/z*: [M + H]⁺ calcd for C₂₄H₂₂NO₂S, 388.1366; found, 388.1357; TLC *R*_f = 0.31 (50% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

(3aS*,9bS*)-6-Methyl-3-methylene-9-phenyl-1-tosyl-1,3a,4,5,7,9b-hexahydro-2H-azuleno[4,5-b]pyrrole-2,8(3H)-dione (54). **54** was prepared according to General Procedure I. Run 1: Allene-yne **43** (4 mg, 0.009 mmol), rhodium biscarbonyl chloride dimer (0.5 mL of a solution of Rh(I)-catalyst in toluene, 1 mg Rh/1 mL toluene, 0.001 mmol), and toluene (1 mL) provided lactam **54** (3 mg, 75% yield) as a white solid after column chromatography (gradient elution with 25–50% ethyl acetate in hexanes). Run 2: Allene-yne **43** (9 mg, 0.02 mmol), rhodium biscarbonyl chloride (1 mg, 0.002), and toluene (2.1 mL) provided lactam **54** (6 mg, 67% yield) as a white solid after column chromatography. ¹H NMR (500 MHz, CDCl₃): δ 7.64 (d, *J* = 8.5 Hz, 2H, Ph-H), 7.40–7.38 (m, 3H, Ph-H), 7.21 (d, *J* = 8.5 Hz, 2H, Ph-H), 7.18–7.14 (m, 2H, Ph-H), 6.16 (d, *J* = 3.5 Hz, 1H, =CH), 5.43 (d, *J* = 3.5 Hz, 1H, =CH), 5.07 (d, *J* = 8.5 Hz, 1H, CH), 3.23–3.22 (m, 1H, CH), 3.23, 3.10 (ABq, *J* = 20.5 Hz, 2H, CH₂), 2.96–2.86 (m, 1H), 2.56–2.47 (m, 1H), 2.43–2.38 (m, 1H), 2.37 (s, 3H, CH₃), 2.01 (s, 3H, CH₃), 1.80–1.74 (m, 1H); ¹³C NMR (125 MHz, CDCl₃): δ 202.6, 166.5, 164.6, 145.3, 140.7, 139.3, 134.8, 134.7, 132.0, 131.4 (2C), 130.9, 130.4, 129.7 (2C), 128.5 (2C), 127.8 (2C), 121.4, 60.2, 41.5, 40.7, 31.9, 29.3, 25.7, 21.8; IR (thin film): 2888, 1714, 1674, 1358, 1160 cm^{−1}; HRMS (TOF MS ES⁺) *m/z*: [M + H]⁺ calcd for C₂₇H₂₆NO₄S, 460.1577; found, 460.1577; TLC *R*_f = 0.23 (50% EtOAc/hexanes), visualized with UV and vanillin.

tert-Butyl (3aS*,9bS*)-6-methyl-3-methylene-2,8-dioxo-9-phenyl-2,3,3a,4,5,7,8,9b-octahydro-1H-azuleno[4,5-b]pyrrole-1-carboxylate (55). **55** was prepared according to General Procedure H.

Run 1: Allene-yne **44** (7 mg, 0.018 mmol), rhodium biscarbonyl chloride dimer (1 mg, 0.002 mmol), and toluene (1.9 mL) provided lactam **55** (3 mg, 40% yield) as a pale yellow solid after column chromatography (gradient elution with 15–50% ethyl acetate in hexanes). Lactam **55** was also prepared according to General Procedure I. Run 2: Allene-yne **44** (16 mg, 0.042 mmol), rhodium biscarbonyl chloride dimer (2 mg, 0.004 mmol), and toluene (4.2 mL) provided lactam **55** (7 mg, 41% yield) as a pale yellow solid after column chromatography. ¹H NMR (500 MHz, CDCl₃): δ 7.35–7.27 (m, 3H, Ph-H), 7.05–7.04 (m, 2H, Ph-H), 6.29 (d, *J* = 3.5 Hz, 1H, =CH), 5.50 (d, *J* = 3.5 Hz, 1H, =CH), 5.12 (d, *J* = 9.0 Hz, 1H, CH), 3.17, 3.06 (AB q, *J* = 21.0 Hz, CH₂), 3.14–3.11 (m, 1H, CH), 3.01–2.92 (m, 1H), 2.56–2.50 (m, 1H), 2.43–2.39 (m, 1H), 2.00 (s, 3H, CH₃), 1.95–1.92 (m, 1H), 1.18 (s, 9H, C(CH₃)₃); ¹³C NMR (125 MHz, CDCl₃): δ 202.7, 165.0, 164.8, 150.4, 142.8, 137.9, 135.7, 131.0 (2C), 130.6, 130.2, 127.9 (2C), 119.9, 83.5, 60.5, 58.5, 40.6, 39.8, 31.9, 28.9, 27.9 (3C), 25.7; IR (thin film): 2969, 2915, 1793, 1757, 1717, 1401, 1302, 1153 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₂₅H₂₈NO₄, 406.2013; found, 406.2013; TLC *R*_f = 0.65 (50% EtOAc/hexanes), visualized with UV and *p*-anisaldehyde.

(3*aS**,9*bS**)-3,6-Dimethyl-1,3*a*,4,5,7,9*b*-hexahydro-2*H*-azuleno[4,5-*b*]pyrrole-2,8(3*H*)-dione (**58**). **58** was prepared according to General Procedure H. Allene-yne **56** (12 mg, 0.059 mmol, 9:1 dr), rhodium biscarbonyl chloride dimer (2 mg, 0.005 mmol), and toluene (5 mL) provided lactam **58** (11 mg, 78% yield, 7.3:1 dr) as a white solid after column chromatography (gradient elution with 25–100% ethyl acetate in hexanes). The diastereomeric ratio was determined based on integration of the resonances corresponding to the lactam methine (4.56, d, CHNH) and lactam methine (**58**, 4.51, d, CHNH). The stereochemistry of the carbon adjacent to the lactam carbonyl was not assigned. ¹H NMR (500 MHz, CDCl₃): δ 7.47 (br s, 1H, NH)*, 7.36 (br s, 1H, NH), 6.10 (s, 1H, CHCO)*, 6.07 (s, 1H, CHCO), 4.56 (d, *J* = 10.5 Hz, 1H, CH)*, 4.51 (d, *J* = 10.0 Hz, 1H, CH), 3.00 (s, 2H, CH₂), 2.76–2.70 (m, 1H), 2.31–2.18 (m, 3H), 1.91 (s, 3H, CH₃), 1.95–1.87 (m, 1H), 1.77–1.70 (m, 1H), 1.20 (d, *J* = 7.0 Hz, 3H, CH₃) *Minor diastereomer where distinguishable; ¹³C NMR (126 MHz, CDCl₃): δ 204.0, 179.5, 173.3, 137.5, 131.2, 126.6, 57.1*, 56.6, 47.3, 45.0, 42.0*, 41.5*, 41.4, 33.3*, 32.7, 28.1, 24.8, 23.7*, 13.8, 12.0*; IR (thin film): 3196, 2893, 1680, 1553, 1438, 1231, 1156 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₁₄H₁₈NO₂, 232.1259; found, 232.1339; TLC *R*_f = 0.11 (100% EtOAc), visualized with UV and *p*-anisaldehyde.

(3*aR**,9*bS**)-3,6-Dimethyl-9-phenyl-1,3*a*,4,5,7,9*b*-hexahydro-2*H*-azuleno[4,5-*b*]pyrrole-2,8(3*H*)-dione (**59**). **59** was prepared according to General Procedure I. Allene-yne **57** (17 mg, 0.061 mmol), rhodium dicarbonyl chloride dimer (3 mg, 0.0061 mmol), and toluene (6 mL) yielded lactam **59** (14 mg, 77% yield) as a white solid after purification by column chromatography (gradient elution with 50–100% ethyl acetate in hexanes). The stereochemistry of the carbon adjacent to the lactam carbonyl was not assigned. ¹H NMR (400 MHz, CDCl₃): δ 7.44–7.38 (m, 3H, Ph-H), 7.17–7.15 (m, 2H, Ph-H), 5.21 (br s, 1H, NH), 4.98 (d, *J* = 7.6 Hz, 1H, CH), 3.19, 3.12 (AB q, *J* = 20.8 Hz, 2H, CH₂), 2.59–2.46 (m, 2H), 2.35–2.24 (m, 2H), 2.09–2.02 (m, 1H), 1.90 (s, 3H, CH₃), 1.86–1.81 (m, 1H), 1.22 (d, *J* = 7.2 Hz, 3H, CH₃); ¹³C NMR (100 MHz, CDCl₃): δ 202.5, 179.8, 163.0, 144.7, 140.9, 131.2, 129.5 (2C), 129.0, 128.8 (2C), 128.7, 55.3, 45.6, 42.4, 42.1, 32.7, 31.9, 24.2, 15.6; IR (thin film): 3374, 2902, 1673, 1431 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₂₀H₂₂NO₂, 308.1645; found, 308.1649; TLC *R*_f = 0.15 (75% EtOAc/hexanes), visualized with UV and vanillin.

General Procedure J. N-Acetyl Lactam Formation from 2° Lactam. (3*aS**,9*bS**)-1-Acetyl-6-methyl-3-methylene-1,3*a*,4,5,7,9*b*-hexahydro-2*H*-azuleno[4,5-*b*]pyrrole-2,8(3*H*)-dione (**60**). A flame-dried test tube equipped with a stir bar, septum, and nitrogen inlet needle was charged with lactam **47** (8 mg, 0.03 mmol) dissolved in DCM (0.4 mL). 4-Dimethylaminopyridine (1 mg, 0.008 mmol) was added, and the solution was cooled to 0 °C. Triethylamine (0.05 mL, 0.3 mmol) was added dropwise via a syringe, followed by dropwise addition of acetic anhydride (0.02 mL, 0.2 mmol) via a syringe. The

reaction was allowed to warm to rt under stirring for 2 h, at which point TLC showed consumption of the starting material. The reaction was diluted with DCM (10 mL) and transferred to a separatory funnel. The organic layer was washed with saturated ammonium chloride (5 mL) and brine (5 mL), dried over magnesium sulfate, gravity filtered, concentrated by rotary evaporation, and purified by silica gel flash column chromatography (gradient elution with 50–75% ethyl acetate in hexanes) to provide N-acetyl lactam **60** (7 mg, 78% yield) as a white, sticky solid. ¹H NMR (400 MHz, CDCl₃): δ 6.27 (d, *J* = 3.6 Hz, 1H, =CH), 5.63 (s, 1H, CHCO), 5.49 (d, *J* = 3.6 Hz, 1H, =CH), 5.03 (d, *J* = 9.2 Hz, 1H, CH), 3.02, 2.94 (AB q, *J* = 20.8 Hz, 2H, CH₂), 2.89–2.75 (m, 2H), 2.65 (s, 3H, CH₃), 2.56–2.41 (m, 1H), 2.37–2.31 (m, 1H), 1.94 (s, 3H, CH₃), 1.87–1.79 (m, 1H); ¹³C NMR (100 MHz, CDCl₃): δ 203.6, 172.4, 171.9, 167.6, 142.4, 136.1, 131.8, 126.9, 120.3, 57.8, 41.0, 39.6, 31.2, 27.7, 25.8, 24.9; IR (thin film): 2890, 1710, 1653, 1555, 1351, 1299, 1145 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₁₆H₁₈NO₃, 272.1281; found, 272.1277; TLC *R*_f = 0.57 (100% EtOAc), visualized with UV and *p*-anisaldehyde.

(3*aS**,9*bS**)-1-Acetyl-6-methyl-3-methylene-9-phenyl-1,3*a*,4,5,7,9*b*-hexahydro-2*H*-azuleno[4,5-*b*]pyrrole-2,8(3*H*)-dione (**59**). **59** was prepared according to General Procedure J. Lactam **45** (9 mg, 0.03 mmol), dimethylaminopyridine (1 mg, 0.008 mmol), triethylamine (0.04 mL, 0.3 mmol), acetic anhydride (0.02 mL, 0.2 mmol), DCM (0.2 mL), and 20 h at rt provided N-acetyl lactam **59** (6 mg, 60% yield) as a pale yellow solid after column chromatography (gradient elution with 15–50% ethyl acetate in hexanes). ¹H NMR (500 MHz, CDCl₃): δ 7.35–7.29 (m, 3H, Ph-H), 7.04–6.49 (m, 2H, Ph-H), 6.29 (d, *J* = 3.2 Hz, 1H, =CH), 5.54 (d, *J* = 3.2 Hz, 1H, =CH), 5.16 (d, *J* = 8.5 Hz, 1H, CH), 3.17, 3.06 (AB q, *J* = 21.0 Hz, 2H, CH₂), 3.12–3.10 (m, 1H, CH), 2.99–2.91 (m, 1H), 2.59–2.51 (m, 1H), 2.44–2.39 (m, 1H), 2.00 (s, 3H, CH₃), 1.91–1.88 (m, 1H), 1.59 (s, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃): δ 202.7, 171.8, 166.4, 164.5, 142.8, 138.3, 135.8, 131.2, 131.0 (2C), 130.6, 127.8 (2C), 120.6, 57.4, 40.5, 39.7, 31.8, 29.7, 28.6, 25.7, 23.9; IR (thin film): 3390, 2892, 1689, 1355, 1273, 1164 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₂₂H₂₂NO₃, 348.1521; found, 348.1518; TLC *R*_f = 0.31 (50% EtOAc/hexanes), visualized with UV and KMnO₄.

(3*aS**,9*bS**)-1-Acetyl-3,6-dimethyl-1,3*a*,4,5,7,9*b*-hexahydro-2*H*-azuleno[4,5-*b*]pyrrole-2,8(3*H*)-dione (**510**). **510** was prepared according to General Procedure J. Lactam **58** (11 mg, 0.05 mmol, 9:1 ratio of diastereomers), dimethylaminopyridine (1 mg, 0.008 mmol), triethylamine (0.07 mL, 0.5 mmol), acetic anhydride (0.02 mL), DCM (0.5 mL), and 6.5 h at rt provided N-acetyl lactam **510** (6 mg, 46% yield) as a white solid after column chromatography (gradient elution with 25–75% ethyl acetate in hexanes). The diastereomeric ratio was determined based on integration of the resonances corresponding to the lactam methine (5.08, d, CHNac) and lactam methine (**510**, 5.02, d, CHNac). ¹H NMR (500 MHz, CDCl₃): δ 5.59 (s, 1H, CHCO), 5.08 (d, *J* = 5.5 Hz, 1H, CH)*, 5.02 (d, *J* = 10.5 Hz, 1H, CH), 3.01, 2.94 (AB q, *J* = 21.0 Hz, 2H, CH₂), 2.87–2.81 (m, 1H), 2.57 (s, 3H, CH₃), 2.41–2.26 (m, 3H), 1.93 (s, 3H, CH₃), 1.87–1.79 (m, 1H), 1.71–1.65 (m, 1H), 1.24 (d, *J* = 7.0 Hz, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃): δ 203.7, 177.1, 172.5, 171.7, 136.5, 131.7, 127.1, 66.0, 58.3, 45.7, 42.3*, 41.0, 31.6, 29.0, 25.4, 24.7, 15.4*, 13.7, * Discernible signals for 1 of 2 diastereomers; IR (thin film): 2893, 1721, 1681, 1661, 1544, 1353, 1254 cm⁻¹; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₁₆H₂₀NO₃, 274.1438; found, 274.1438; TLC *R*_f = 0.46 (75% EtOAc/hexanes), visualized with UV and vanillin.

Synthesis of α-Methylene-γ-lactone **517.** 4-(3-Hydroxybutyl)-3-methylene-5-(phenylethynyl)dihydrofuran-2(3*H*)-one (**513**). A flame-dried, round-bottomed flask equipped with a Teflon-coated stir bar was charged with **512** (1.2 g, 2.9 mmol, E/Z = 2.5:7.5), 3-phenyl-2-propynal (**12**) (757 mg, 5.8 mmol), and toluene (7.8 mL) and cooled to 0 °C. The resulting solution was treated with trifluoromethanesulfonic acid (43.6 mg, 0.29 mmol) and maintained at 0 °C under a nitrogen atmosphere for 12 h. The mixture was then diluted with NH₄Cl (aq):NH₄OH (9:1, v/v, 58 mL) and extracted

with diethyl ether (3 × 60 mL). The combined extracts were washed with brine (2 × 30 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The crude residue was loaded onto a 100 g SNAP column and purified using a Biotage normal phase automated purification system with a gradient of 15–90% diethyl ether/pentane to afford the title compound **S13** (0.375 g, 48%) as a yellow oil. The product was obtained as a mixture of isomers in a *trans/cis* 8.5:2.5 ratio. A pure fraction of the *trans*-lactone was collected for characterization purposes. ¹H NMR (CDCl₃, 500 MHz): δ 7.43–7.42 (m, 2H, Ph-H), 7.32–7.30 (m, 3H, Ph-H), 6.32 (d, *J* = 2.5 Hz, 1H, =CH), 5.68 (d, *J* = 2.5 Hz, 1H, =CH), 4.99–4.98 (m, 1H, CH), 3.87–3.85 (m, 1H), 3.20–3.18 (m, 1H), 2.23 (br s, 1H, OH), 1.83–1.79 (m, 1H), 1.63–1.58 (m, 3H), 1.23–1.21 (m, 3H); ¹³C NMR (CDCl₃, 100 MHz): δ 169.3, 137.4, 131.6 (2C), 129.0, 128.2 (2C), 123.0, 121.3, 87.7, 85.0, 72.3, 67.3, 46.8, 35.3, 29.3, 23.5; IR (thin film): 3465, 2966, 2921, 2855, 2230, 1772, 1670, 1491, 1446, 1405, 1446, 1368, 1269, 1135, 968 cm^{−1}; HRMS (TOF MS ESI+) *m/z*: [M + K]⁺ calcd for C₁₇H₁₈O₃K, 309.0893; found, 309.0901; TLC R_f = 0.3 (80% diethyl ether/pentane), visualized with UV and KMnO₄.

3-Methylene-4-(3-oxobutyl)-5-(phenylethynyl)dihydrofuran-2(3H)-one (S14). A flame-dried, round-bottomed flask equipped with a Teflon-coated stir bar was charged with **S13** (0.11 g, 0.41 mmol, pure *trans*) and CH₂Cl₂ (5.8 mL). The resulting solution was treated with Dess–Martin periodinane (0.21 g, 0.49 mmol). The progress of the reaction was monitored by TLC and upon completion (3 h), and the mixture was concentrated under reduced pressure. The crude residue was loaded onto a 25 g SNAP column and purified using a Biotage normal phase automated purification system with a gradient of 50–100% diethyl ether/pentane to afford the title compound **S14** (96 mg, 88%) as a slightly yellow oil. ¹H NMR (CDCl₃, 500 MHz): δ 7.43–7.31 (m, 5H, Ph-H), 6.32 (d, *J* = 2.5 Hz, 1H, =CH), 5.67 (d, *J* = 2.5 Hz, 1H, =CH), 4.92 (d, *J* = 6 Hz, 1H, CH), 3.22–3.20 (m, 1H), 2.68–2.63 (m, 2H), 2.15 (s, 3H, CH₃), 2.10–2.05 (m, 1H), 1.91–1.84 (m, 1H); ¹³C NMR (CDCl₃, 125 MHz): δ 206.9, 169.8, 137.2, 131.7 (2C), 129.2, 129.4 (2C), 123.0, 121.2, 88.0, 84.8, 72.1, 46.0, 39.4, 30.0, 26.0; IR (thin film): 3056, 2925, 2235, 1773, 1715, 1659, 1487, 1437, 1413, 1368, 1270, 1140, 980 cm^{−1}; HRMS (TOF MS ES+) *m/z*: [M + K]⁺ calcd for C₁₇H₁₆O₃K, 307.0737; found, 307.0745; TLC R_f = 0.7 (80% diethyl ether/pentane), visualized with UV and KMnO₄.

4-Methylene-5-oxo-2-(phenylethynyl)tetrahydrofuran-3-yl)-butan-2-yl Acetate (S15). A flame-dried, 50 mL Schlenk tube equipped with a Teflon-coated stir bar was charged with cerium(III) trichloride (anhydrous beads, 1.13 g, 4.56 mmol) in a nitrogen-filled glovebox. The Schlenk tube was removed from the glovebox, THF (8.7 mL) was added, and the suspension was sonicated for 4 h. The resulting solution was stirred at rt for 12 h, cooled to 0 °C, treated with ethynyl magnesium bromide (0.5 M in THF, 9.62 mL, 4.80 mmol), and stirred at 0 °C for 45 min. The solution was then cooled to −78 °C and cannulated into a round-bottom flask containing a solution of ketone **S14** (0.129 g, 0.48 mmol) in THF (11.7 mL) cooled to −78 °C. After stirring for 30 min at −78 °C, the solution was then diluted with saturated ammonium chloride (5 mL) and diethyl ether (10 mL). After stirring for 1 h at rt, the mixture was filtered through a Celite plug, and the filter cake was rinsed with diethyl ether (60 mL). The organic phase was separated, and the aqueous layer was extracted with diethyl ether (2 × 20 mL). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure to afford **S15** (0.178 g, 0.60 mmol). The propargyl alcohol was not stable to column chromatography, so it was taken on immediately to the next step. Crude **S15** was dissolved in CH₂Cl₂ (2.6 mL), and 4-dimethylaminopyridine (7.42 mg, 0.061 mmol), triethylamine (0.85 mL, 6.07 mmol), and acetic anhydride (0.29 mL, 3.03 mmol) were added. The resulting solution was stirred at rt for 12 h, diluted with ammonium chloride (5 mL), filtered through a Celite plug, and rinsed with CH₂Cl₂. The phases were separated, and the aqueous layer was extracted with CH₂Cl₂ (2 × 20 mL). The combined organic layers were washed with brine (20 mL), dried over Na₂SO₄, filtered,

concentrated under reduced pressure, and purified by flash chromatography (25% diethyl ether/pentane) to afford the title compound **S15** (0.132 g, 82%) as a slightly orange oil. ¹H NMR (CDCl₃, 400 MHz): δ 7.45–7.31 (m, 5H, Ph-H), 6.34 (d, *J* = 2.8 Hz, 1H, =CH), 5.67 (d, *J* = 2.8 Hz, 1H, =CH), 4.98 (d, *J* = 5.2 Hz, 1H, CH), 3.22–3.20 (m, 1H, CH), 2.55 (s, 1H, =CH), 2.07–2.02 (m, 1H), 2.00 (d, *J* = 1.2 Hz, 3H, COCH₃), 1.88–1.70 (m, 3H), 1.68 (d, *J* = 0.8 Hz, 3H, CH₃); ¹³C NMR (CDCl₃, 125 MHz): δ 169.2, 169.0, 137.3, 131.8 (2C), 129.2, 128.4 (2C), 123.3, 121.4, 88.1, 84.9, 83.0, 74.2, 74.1, 72.2, 46.8, 38.1, 27.7, 26.6, 21.9; IR (thin film): 3289, 2929, 2855, 2231, 2116, 1769, 1736, 1667, 1491, 1438, 1368, 1242, 1131, 1103, 1013 cm^{−1}; HRMS (TOF MS ES+) *m/z*: [M + Na]⁺ calcd for C₂₁H₂₀O₄Na, 359.1259; found, 359.1255; TLC R_f = 0.7 (80% diethyl ether/pentane), visualized with silica gel, UV, KMnO₄ stain.

3-Methylene-4-(3-methylpenta-3,4-dienyl)-5-(phenylethynyl)-dihydrofuran-2(3H)-one (S16). A flame-dried, round-bottomed flask equipped with a Teflon-coated stir bar was charged with **S15** (115 mg, 0.33 mmol) and THF (3.3 mL). The resulting solution was treated with dimethylamine (0.45 mL, 0.83 mmol, 2 M in THF), stirred at rt under a nitrogen atmosphere for 11 h and concentrated under reduced pressure to afford the crude Michael adduct (139 mg, 0.37 mmol). After dilution with toluene (5.7 mL, degassed), (triphenylphosphine)copper hydride hexamer (Stryker's reagent) (716 mg, 0.37 mmol) was added, and the resulting solution was stirred at rt under a nitrogen atmosphere for 3 h, quenched with saturated ammonium chloride (5 mL), and stirred opened to air for 12 h. The crude mixture was filtered through a Celite plug and rinsed with diethyl ether. The phases were separated, and the aqueous layer was extracted with diethyl ether (2 × 20 mL). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure to afford crude allene-yne (153 mg, 0.47 mmol). The latter was diluted with CH₂Cl₂ (18 mL), treated with silica gel (4 g, 66 mmol), capped with a glass stopper, and stirred at rt for 11 h. The solution was then filtered, rinsed with CH₂Cl₂, concentrated under reduced pressure, and purified by flash chromatography (10% diethyl ether/pentane) to afford the title compound **S16** (36.1 mg, 39%) as a colorless oil. ¹H NMR (CDCl₃, 500 MHz): δ 7.44–7.32 (m, 5H, Ph-H), 6.35 (d, *J* = 2.5 Hz, 1H, =CH), 5.69 (d, *J* = 2.5 Hz, 1H, =CH), 4.98 (d, *J* = 5.5 Hz, 1H, CH), 4.62–4.60 (m, 2H, =CH₂), 3.20–3.18 (m, 1H, CH), 2.02–2.00 (m, 1H), 1.81–1.76 (m, 2H), 1.69–1.67 (t, *J* = 3.0 Hz, 3H, CH₃), 1.66–1.64 (m, 2H); ¹³C NMR (CDCl₃, 175 MHz): δ 206.0, 169.2, 137.7, 131.7 (2C), 129.0, 128.3 (2C), 122.6, 121.5, 97.6, 87.7, 85.1, 74.6, 72.4, 46.8, 32.8, 32.5, 18.7; IR (thin film): 3289, 2925, 2844, 2231, 1957, 1769, 1491, 1442, 1262, 1131 cm^{−1}; HRMS (TOF MS ES+) *m/z*: [M + H]⁺ calcd for C₁₉H₁₉O₂, 279.1385; found, 279.1375; TLC R_f = 0.27 (20% diethyl ether/pentane), visualized with UV and KMnO₄.

3-Methylene-9-phenyl-3,3a,4,5-tetrahydroazuleno[4,5-b]furan-2,8(7H,9bH)-dione (S17). A flame-dried vial (15 × 45 mm) equipped with a Teflon-coated stir bar and a septum cap was charged with allene-yne **S16** (15 mg, 0.054 mmol, 1 equiv) and toluene (1.6 mL, toluene degassed by bubbling with nitrogen for ~5 min). The tube was placed under vacuum for 3–5 s and refilled with carbon monoxide (3×). To the allene-yne solution was added [Rh(CO)₂Cl]₂ (1 mg, 5.6 × 10^{−3} mmol, 0.10 equiv) in one portion by temporary removal of the septum, and the vial was placed under vacuum and refilled with carbon monoxide (3×). The vial was placed in an oil bath (preheated to 90 °C) and stirred under carbon monoxide. After 25 min, TLC indicated completion, and the mixture was cooled to rt, passed through a short plug of Celite using diethyl ether, and concentrated in vacuo. Purification of the residue by flash chromatography (75% diethyl ether/pentane) afforded the title compound **S17** (8.8 mg, 54%) as a slightly yellow oil. ¹H NMR (CDCl₃, 300 MHz): δ 7.37–7.33 (m, 3H, Ph-H), 7.26–7.23 (m, 2H, Ph-H), 6.22 (d, *J* = 3.3 Hz, 1H, =CH), 5.54 (d, *J* = 3.3 Hz, 1H, =CH), 5.45 (d, *J* = 9.9 Hz, 1H, CH), 3.16–3.13 (m, 2H, COCH₂), 3.13–3.06 (m, 1H, CH), 2.77–2.68 (m, 1H), 2.46–2.35 (m, 2H), 1.99 (s, 3H, CH₃), 1.91–1.85 (m, 1H); ¹³C NMR (175 MHz,

CDCl₃): δ 202.1, 168.1, 162.1, 140.9, 138.9, 135.2, 131.2, 130.2, 129.7 (2C), 127.9, 127.3 (2C), 120.7, 78.3, 43.9, 40.6, 31.3, 25.9, 24.6; IR (thin film): 2929, 2860, 2247, 1777, 1691, 1446, 1311, 1258, 1131, 1037, 1004 cm⁻¹; HRMS (TOF MS ES+) m/z : [M + H]⁺ calcd for C₂₀H₁₉O₃, 307.1334; found, 307.1339; TLC R_f = 0.3 (70% diethyl ether/pentane), visualized with UV and KMnO₄.

General Procedure K: Reaction of α -Methylene- γ -lactams and Lactone **S17 with Cysteamine and Monitoring the Reaction Progress by ¹H NMR.** *Sample Preparation for Pseudo-First-Order Reaction Kinetic Studies.* Pseudo-first-order rate constants were determined using a procedure modified from the original report.⁴² A 1-dram scintillation vial was charged with 1–2 mg of the lactam (1 equiv) and dissolved in CDCl₃ to afford a concentration of 11.5 mM. A second 1-dram scintillation vial was charged with cysteamine (15 equiv) and dissolved in CDCl₃ (The volume of CDCl₃ used for this dilution is equal to that used to dissolve the lactam). These two solutions were transferred *via* a glass pipet to a third 1-dram scintillation vial, making the final lactam concentration 5.7 mM. This vial is capped, mixed for 10 s using a vortex mixer, and transferred to a 5 mm NMR tube.

Monitoring the Reaction Progress by ¹H NMR without an Internal Standard. A ¹H NMR spectrum was obtained immediately after transferring the prepared sample to the NMR tube, and at regular intervals of time thereafter. The solution was maintained at ambient temperature (22 °C) over the course of the reaction. The progress of the reaction was monitored by the disappearance of methylene resonances H_a, H_b, and methine H_c and the simultaneous appearance of H_d of the thio-adduct (see eq S1 for identity of H_a, H_b, H_c, and H_d). For each timepoint, the integration value corresponding to H_a was set to 1.0, and the fraction of remaining lactam was calculated following eq S2 using the integration values of H_a, H_b, H_c, and H_d. The natural log-transformed values of the fraction remaining were plotted against time to afford a linear plot. Using the linear form of the first-order rate equation, the slope of the line-of-best-fit is equal to the pseudo-first-order rate constant ($k_{\text{pseudo1st}}$) (eq S3). The reaction half-life ($t_{1/2}$) is calculated using eq S4.

Monitoring the Reaction Progress by ¹H NMR with an Internal Standard. Two lactams, **49** and **52**, were reanalyzed following the procedure above with an internal standard, hexamethylbenzene (1 equiv). The half-lives for these reactions with and without the internal standard were nearly identical, indicating that monitoring the reactions in the absence of internal standard is reliable (*vide infra*).

Reaction of α -Methylene- γ -lactam **45 with Cysteamine.** This reaction was performed according to General Procedure K. Tricyclic lactam **45** (8 mg, 0.026 mmol), cysteamine (30 mg, 0.39 mmol), and chloroform-*d* (1 mL) were used (see Table S3 and Figures S1–S3).

Reaction of α -Methylene- γ -lactam **49 with Cysteamine.** This reaction was performed according to General Procedure K. Lactam **49** (1.65 mg, 0.0043 mmol), cysteamine (5.0 mg, 0.0649 mmol), and chloroform-*d* (0.76 mL) were used (see Table S4 and Figures S4–S6).

Reaction of α -Methylene- γ -lactam **49 with Cysteamine Using an Internal Standard.** This reaction was performed and monitored using an internal standard according to General Procedure K. Lactam **49** (1.7 mg, 0.0043 mmol), cysteamine (5 mg, 0.065 mmol), hexamethylbenzene (0.7 μ L mmol), and chloroform-*d* (0.76 mL) were used. The fraction remaining was calculated using eq S5 as the initial NMR showed a 2:1 integrative ratio of hexamethylbenzene/lactam (see Table S5 and Figures S7–S9).

Reaction of α -Methylene- γ -lactam **51 with Cysteamine.** This reaction was monitored according to General Procedure K. Lactam **51** (0.98 mg, 0.0022 mmol), cysteamine (2.8 mg, 0.0334 mmol), and chloroform-*d* (0.4 mL) were used (see Table S6 and Figures S10–12).

Reaction of α -Methylene- γ -lactam **52 with Cysteamine.** This reaction was performed according to General Procedure K. Lactam **52** (1.2 mg, 0.0029 mmol), cysteamine (3.4 mg, 0.044 mmol), and chloroform-*d* (0.5 mL) were used (see Table S7 and Figures S13–15).

Reaction of α -Methylene- γ -lactam **52 with Cysteamine Using an Internal Standard.** This reaction was performed and monitored using an internal standard according to General Procedure K. Lactam **52** (1.2 mg, 0.0029 mmol), cysteamine (3.4 mg, 0.044 mmol), hexamethylbenzene (0.4 μ L, 0.0029 mmol), and chloroform-*d* (0.05 mL) were used. The fraction remaining was calculated using eq S6 as the initial NMR showed a 1:1 integrative ratio of hexamethylbenzene/lactam (see Table S8 and Figures S16–18).

Reaction of α -Methylene- γ -lactam **54 with Cysteamine.** Lactam **54** (2 mg, 0.004 mmol) was dissolved in chloroform-*d* (0.7 mL) and transferred to an NMR tube. Cysteamine (5 mg, 0.07 mmol) was dissolved in chloroform-*d* separately and transferred to the tube containing lactam **54** *via* a pipet, and the tube was capped and shaken vigorously by hand for 2 min. The progress of the reaction was monitored ¹H NMR, specifically the disappearance of the methylene resonances (H_a and H_b) that are observed as doublets at δ 6.17 and 5.44. Complete disappearance of these resonances was observed in less than 10 min.

Reaction of α -Methylene- γ -lactam **60 with Cysteamine.** This reaction was monitored according to General Procedure K. Lactam **60** (1.1 mg, 0.0041 mmol), cysteamine (5.0 mg, 0.065 mmol), and chloroform-*d* (0.80 mL) were used (see Table S9 and Figures 19–21).

Reaction of α -Methylene- γ -lactone **S17 with Cysteamine.** This reaction was performed according to General Procedure K. Lactone **S17** (1.0 mg, 0.0033 mmol), cysteamine (3.8 mg, 0.049 mmol), and chloroform-*d* (0.66 mL) were used (see Table S10 and Figures 23–24).

Structural Confirmation of *tert*-Butyl Thiol Adduct. (3aS*,9bS*)-3-((*tert*-Butylthio)methyl)-1,6-dimethyl-9-phenyl-1,3a,4,5,7,9b-hexahydro-2H-azuleno[4,5-*b*]pyrrole-2,8(3H)-dione (**S11**). A 2 mL vial equipped with a magnetic stir bar was charged with tricyclic lactam **48** (10 mg, 0.03 mmol), *tert*-butyl thiol (0.02 mL, 0.2 mmol), acetone (0.5 mL), and phosphate-buffered saline (pH = 8.0, 0.5 mL). The vial was capped with a septum and allowed to stir at rt for 24 h. Only the starting material was detected by TLC analysis, so the pH was increased to 12 (pH paper) by addition of potassium hydroxide (2 mg, 0.04 mmol) and the starting material was consumed within 30 min as observed by TLC. The solution was poured into a separatory funnel containing deionized water (3 mL) and DCM (10 mL). The layers were separated. The aqueous layer was extracted with DCM (2 $\times$ 5 mL). The organic layers were combined, washed with brine (5 mL), dried over magnesium sulfate, filtered, and concentrated by rotary evaporation and purified by flash column chromatography on a silica gel (gradient elution with 25–50% ethyl acetate in hexanes) to provide thiol adduct **S11** (3 mg, 23%, 1.6:1 diastereomeric ratio) as a clear oil. The diastereomeric ratio was determined by integration of resonances corresponding to H_a at δ 4.66 (J = 10.0 Hz) for the minor diastereomer and δ 4.50 (J = 8.5 Hz) for the major diastereomer. ¹H NMR (500 MHz, CDCl₃): δ 7.33–7.27 (m, 3H), 7.23–7.21 (m, 2H), 4.66 (d, J = 10.0 Hz, 1H)*, 4.50 (d, J = 8.5 Hz, 1H), 3.17–3.07 (m, 3H), 2.94–2.79 (m, 2H), 2.75–2.69 (m, 1H), 2.51–2.41 (m, 2H), 2.37–2.30 (m, 2H), 1.98 (s, 3H), 1.90 (s, 3H), 1.37 (s, 9H), *Discernible signals for 1 of 2 diastereomers; ¹³C NMR (100 MHz, CDCl₃): δ 202.92, 202.86*, 175.3, 166.1, 138.5, 137.7, 137.1, 130.8, 130.73*, 130.67 (2C), 128.5, 128.4*, 127.7 (2C), 127.6*, 62.7, 61.7*, 50.3, 44.2, 43.2*, 42.7, 42.6*, 40.6, 34.3, 33.4*, 32.6, 31.1, 31.0*, 30.8, 30.4*, 30.32, 30.30*, 29.4, 25.7, 25.6*, 21.0, *Discernible signals for 1 of 2 diastereomers; IR (thin film): 3407, 2927, 1686, 1445, 1079 cm⁻¹; HRMS TOF MS ES + [M + H]: C₂₅H₃₂NO₂S calcd, 410.2148 Found: 410.2144; TLC R_f = 0.32 (50% EtOAc/hexanes) visualized with UV and vanillin.

Biological Materials and Methods. Preparation and Storage of Compound Stock Solutions. Compound stock solutions were prepared in biological grade DMSO (40–100 mM concentrations) and stored at –20 °C when not in use. Compound purities were assessed prior to conducting biological assays by analytical reverse-phase HPLC. Fresh solutions were prepared as needed.

General Protocol for HPLC Analysis of Synthesized Compounds. DMSO stock solutions of synthesized compounds were dissolved in a

1:1 mixture of methanol and distilled and deionized water (ddH₂O) containing trifluoroacetic acid (TFA, 0.1% v/v) and analyzed on an Agilent 1200 series instrument equipped with a diode array detector and a Zorbax SB-C18 column (4.6 × 150 mm, 5 μm, Agilent Technologies). The analysis method (1 mL/min flow rate) starts with an isocratic eluent system of 1:9 MeCN (containing TFA, 0.1% v/v; solvent A):ddH₂O (containing TFA, 0.1% v/v; solvent B) from 0 to 2 min, followed by a linear gradient of 10:90–85:15 A/B from 2 to 24 min, followed by 85:15–95:5 A/B from 24 to 26 min, and finally an isocratic eluent system of 95:5 A/B from 26 to 30 min. Wavelength monitored = 215 nm (see Table S11 and Figures S25–S32).

Protocol for Mammalian Cell Culture. Cell lines were kept in a humidified 19% O₂, 5% CO₂, and 37 °C environment. A549/NF-κB-luciferase cells were cultured as previously described.⁵³ HEK293/NF-κB-SEAP (Novus Biologicals #NBP2-26260) was cultured in Dulbecco's Modified Eagle's medium (Corning) supplemented with 10% v/v fetal bovine serum (Gibco), 100 IU penicillin (ATCC), 100 μg/mL streptomycin (ATCC), and 0.5 mg/mL G418 (Geneticin).

NF-κB Luciferase Reporter Assay. Performed as previously described by our laboratory,⁵³ with one change: compounds were dosed to cells for 30 min, followed by induction with TNF-α for 8 h.

NF-κB Secreted Placental Alkaline Phosphatase (SEAP) Reporter Assay. HEK293/NF-κB-SEAP cells (Novus Biologicals #NBP2-26260) were seeded in standard 96-well cell culture plates (Costar) at a density of 5000 cells/well (50 μL/well) 24 h before dosing with compounds. Compounds serially diluted in prewarmed media were dosed to cells (50 μL/well, final DMSO concentration: 0.5%), and following a 30 min incubation period, all wells except for the noninduced (N) control were induced with TNF-α (10 μL; final well concentration: 22.5 ng/mL; Invitrogen). Noninduced control wells received 1× PBS (10 μL). After 8 h, 25 μL of media was harvested from each well and plated into a second, 96-well white plate with a clear bottom (Costar). Using the NovaBright Phospha-Light EXP Assay Kit for SEAP Reporter Gene Detection (Invitrogen #N10577) and a BioTek H1 Synergy microplate reader, SEAP was quantified as per the manufacturer's protocol. Each experimental condition had three technical replicates and each experiment was performed in biological triplicate. Activity values were obtained by averaging the mean activity values from each biological replicate. The standard deviation is obtained by propagating the standard deviations of each of the individual biological replicates. Statistical analyses were performed using Microsoft excel and plotted in GraphPad Prism (v. 5.0b).

Vero Cell Viability Assay. Performed as previously described by our laboratory.⁵³

ALARM NMR Assay. Lactone **S17** and lactam **51** were tested by ALARM NMR as previously described with minor modifications.⁵⁸ Briefly, test compounds (400 μM final concentration) were incubated with ¹³C-methyl labeled La antigen (50 μM final concentration) at 37 °C for 90 min. Each compound was tested in the absence and presence of 20 mM DTT. Samples were loaded into Bruker 3 mM SampleJet tubes with 160 μL total sample volumes and stored at ambient temperature while in queue. Data were recorded at 25 °C on a Bruker 700 MHz NMR spectrometer equipped with a cryoprobe (Bruker). Data were collected for each compound every hour for 10 h. Nonreactive compounds were identified by the absence of chemical shifts (¹³C-methyl) independent of the presence of DTT. Reactive compounds were identified by the presence of chemical shifts (¹³C-methyl) and peak attenuations in certain diagnostic peaks in the absence of DTT.

La Antigen Expression. The La antigen protein was expressed and purified similar to a previously reported protocol.⁵⁸ The protocol specifies the expression of ¹³C-labeled protein, but for MS experiments, non-¹³C-labeled (unlabeled) protein was prepared by skipping the secondary culture spin down and resuspension in defined media steps.

La Antigen MS Sample Preparation. The tryptic digestion of purified recombinant protein was adapted from a previously published protocol.⁶⁹ All solutions utilized in this section were purified by centrifugation with a 3 kDa molecular weight cutoff (MWCO) filter

(Amicon, MilliporeSigma) and collection of the flow-through. Recombinant La antigen (2.5 μL; 2 μg/μL in aqueous 25 mM Na₂HPO₄/NaH₂PO₄ buffer, pH 7.0) was added to a solution of the test compound (15.5 μL of 10 mM DMSO stock). The sample was further diluted to 310 μL with 20 mM Tris buffer (pH 8.0) and incubated for 1 h at 37 °C. Next, unlabeled IAD was added to a final concentration of 50 mM, and the solution was incubated in the dark for 1 h at room temperature. The protein was then isolated by spinning down the samples in a 3 kDa MWCO filter, diluting the sample with distilled and deionized H₂O (ddH₂O, 500 μL) and isolating the protein again through the MWCO filter (3× total). The protein was then evaporated to dryness overnight (SpeedVac). The dried protein-compound samples were resuspended in 20 μL of a freshly prepared aqueous denaturing solution (8 M urea, 50 mM ammonium bicarbonate, 50 mM DTT) and incubated at 37 °C for 1 h. A solution of 50 mM d₄-IAD (Cambridge Isotope Laboratories, item #DLM-7249-PK) in ddH₂O was prepared, and 20 μL was added to each sample. The samples were incubated for 1 h in the dark at room temperature. Next, 138 μL of 100 mM ammonium bicarbonate aqueous solution was added to each sample in order to dilute the urea concentration for trypsin digestion. A 1 μg/μL solution of trypsin (Promega, catalog #V5280) in 50 mM acetic acid was diluted to 100 ng/μL with 100 mM aqueous ammonium bicarbonate, and 10 μL of the resulting solution was added to each sample. Acetonitrile was immediately added to each sample to a final concentration of 10% (v/v). Samples were incubated in a rotating (800 rpm) heat block overnight at 37 °C. Glacial acetic acid was added until the pH of each sample was less than 4.0 (~5 μL), and then the samples were desalted using C18 resin pipette tips according to the manufacturer's protocol (Pierce C18 Tips 10 μL bed, ThermoFisher). Desalted samples were evaporated to dryness (SpeedVac) and then reconstituted in 20 μL of 98:2 LC-MS-grade H₂O/MeCN containing 0.1% formic acid (v/v).

La Antigen MS Analysis. Peptides within each sample were separated with a front-end Dionex UltiMate 3000 ultrahigh performance liquid chromatography instrument. Peptides were separated using a home-packed analytical Luna C18 (100 Å pore, 5 μm particles) reverse-phase column (75 μm ID × 200 mm, 10 μm emitter orifice, Phenomenex), where MS-grade water with 0.1% formic acid was eluent A and acetonitrile with 0.1% formic acid was eluent B at room temperature. Initially, an isocratic 2% B elution was run for 5 min (1 μL/min flow rate). Then, the gradient elution (0.3 μL/min flow rate for all gradient steps) began by running from 2% B to 10% B over 5 min, followed by 10% B to 25% B over 40 min, and then 25% B to 40% B over 10 min. The elution gradient further increased from 40% B to 90% B over 1 min, then held at 90% B for 4 min before decreasing from 90% B to 2% B over 0.5 min, and re-equilibrated at 2% B for 4.5 min (1 μL/min flow rate). Eluted peptides were analyzed with a LTQ Orbitrap Velos (ThermoFisher) in the Nth Order Double Play mode. The mass spectrometer utilized an electrospray ionization source with a source voltage of +2.5 kV. MS¹ scan range was *m/z* 220.0–1800.0. For MS², the precursor ion mass of peptides containing the target cysteines in the La antigen were predicted using Skyline software.⁷⁰ The *m/z* of the double and triple-charged peptides were calculated for three peptides: FSGDLDDQTC²⁴⁵R, IGC²³²LLK, and SLEEKIGC²³²LLK. Both IGCLLK and SLEEKIGCLLK were searched as our initial experiments suggested that the SLEEKIGCLLK peptide may be more readily identified with the small carbamidomethylation modifications, but the IGCLLK peptide may be more readily identified with the larger compound modifications. Seven possible cysteine modifications were evaluated: carbamidomethylation (+57.0215 Da), d₂-carbamidomethylation (+59.0340 Da), d₄-carbamidomethylation (+61.0466 Da), 2-chloro-1,4-naphthoquinone (+306.1041 Da), CPM (+402.1580 Da), lactone **S17** (+306.1256 Da), and lactam **51** (+449.1602 Da). In total, 42 precursor ions were selected for MS² analysis (calculated MS¹ precursors are listed in the calculated precursor ions and MS² y- and b-ion list Excel Supporting Information file). Only the most intense ion from the specified parent ion list were selected for MS², repeated for the top 12 peaks. Already examined precursor ions were excluded for 15 s. Ions with an unassigned charge state or a +1 charge

state were rejected. Selected precursor ions (minimum signal = 5000) were analyzed using higher-energy C-trap dissociation (HCD) with a normalized collision energy of 35 V, an isolation width of 2, and the first mass value of m/z 100.00.

Raw data were analyzed manually using Xcalibur Quan Browser software (Thermo Fisher). Raw data files and associated peak list files were uploaded to the MassIVE UCSD public repository and can be downloaded at <ftp://massive.ucsd.edu/MSV000085781/>. MS² spectra of precursor ions were searched for peaks matching the expected y- and b-ion series based on peptide sequences (calculated y- and b-ion masses are identified in the calculated precursor ions and MS² y- and b-ions list Excel file [Supporting Information](#)). Peptides were considered positively identified if all or all but one y-ions and at least two b-ions could be identified in a given spectra. Example spectra are included in SI ([Figures S27–S37](#)). CSV files containing the m/z and intensity of peaks of each positively identified MS² spectrum can be found as [Supporting Information](#) Excel files. Each sample was run in technical triplicate and analyzed independently.

Peptides SLEEKIG²³²LLK and IGC²³²LLK were searched and identified in the raw data, but the results indicate that C232 is less reactive than C245. For example, in the DMSO control samples, C232 peptides are found adducted with both IAD and d₂-IAD, indicating that C232 is inefficiently labeled with IAD in the initial alkylation step with the properly folded La antigen protein. Consequently, C232 thiols are then alkylated with d₄-IAD following protein denaturation. The same pattern of sluggish reactivity of C232 was observed in the fluconazole negative control. These data are also consistent with the published ALARM NMR literature, in which titration experiments indicated that C245 reacts more quickly than C232.⁵⁶ Additional whole-protein MS studies from the same paper indicate both single- and double-adduct species, likely from the partial adduction at one cysteine. Taking these reactivity features into consideration, we elected to only use the FSGDLDDQT²⁴⁵R peptide for our studies. However, the data for C232 peptides are shown in [Table S12](#) for general reference.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.jmedchem.0c01464>.

Structures of Supporting Information compounds; relative NF- κ B activities and relative cellular viabilities for lactams dosed at various concentrations to A549/NF- κ B-luc and HEK293/NF- κ B-SEAP cells; reaction of α -methylene- γ -lactams and lactone S17 with cysteamine: monitoring reaction progress by ¹H NMR; biological materials and methods; peptide mass spectra; HSQC of α -methylene- γ -lactam S2; and ¹H NMR and ¹³C NMR spectra for compounds 9, 11–13, 14a, 14b, 15a, 15b, 18a, 18b, 19a, 19b, 20a, 20b, 21a, 21b, 22a, 22b, 23a, 23b, 24a, 25a, 26, 27a, 28a, 29a, 32a, 30a, 37–40, 42, 43, 44–60, 64, 65, S2–S3, S5, S8–S10, and S13–S17 ([PDF](#))

Molecular formula strings for compounds 9, 11–13, 14a, 14b, 15a, 15b, 18a, 18b, 19a, 19b, 20a, 20b, 21a, 21b, 22a, 22b, 23a, 23b, 24a, 25a, 26, 27a, 28a, 29a, 32a, 31a, 30a, 37–40, 42, 43, 44–60, 64, 65, S2–S3, S5, S8–S11, and S13–S24 ([CSV](#))

Calculated precursor ions and MS² y- and b-ion list ([XLSX](#))

Positively identified MS² spectra ion lists for S17-treated samples ([XLSX](#))

Positively identified MS² spectra ion lists for Fluconazole-treated samples ([XLSX](#))

Positively identified MS² spectra ion lists for DMSO-treated samples ([XLSX](#))

Positively identified MS² spectra ion lists for CPM-treated samples ([XLSX](#))

Positively identified MS² spectra ion lists for Compound S1-treated samples ([XLSX](#))

Accession Codes

MS raw data was deposited at the MassIVE UCSD Public Repository and can be accessed at <ftp://massive.ucsd.edu/MSV000085781>.

■ AUTHOR INFORMATION

Corresponding Authors

Daniel A. Harki – Department of Medicinal Chemistry and Department of Chemistry, University of Minnesota, Minneapolis, Minnesota 55455, United States; [orcid.org/0000-0001-5950-931X](#); Email: daharki@umn.edu

Kay M. Brummond – Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, United States; [orcid.org/0000-0003-3595-6806](#); Email: kbrummon@pitt.edu

Authors

Paul A. Jackson – Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, United States

Henry A. M. Schares – Department of Medicinal Chemistry, University of Minnesota, Minneapolis, Minnesota 55455, United States

Katherine F. M. Jones – Department of Chemistry, University of Minnesota, Minneapolis, Minnesota 55455, United States

John C. Widen – Department of Medicinal Chemistry, University of Minnesota, Minneapolis, Minnesota 55455, United States

Daniel P. Dempe – Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, United States

Francois Grillet – Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, United States

Matthew E. Cuellar – Department of Medicinal Chemistry and Institute for Therapeutics Discovery and Development, University of Minnesota, Minneapolis, Minnesota 55455, United States; [orcid.org/0000-0003-3365-2120](#)

Michael A. Walters – Department of Medicinal Chemistry and Institute for Therapeutics Discovery and Development, University of Minnesota, Minneapolis, Minnesota 55455, United States; [orcid.org/0000-0001-5650-9277](#)

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acs.jmedchem.0c01464>

Funding

This work was supported by the NIH (R01-GM110129 to DAH) and the University of Pittsburgh. Funding for NMR instrumentation used for ALARM NMR assay was provided by the UMN Office of the Vice President for Research, UMN Medical School, UMN College of Biological Science, NIH, NSF and the Minnesota Medical Foundation. Protein MS was performed at the Analytical Biochemistry Core Facility of the UMN Masonic Cancer Center, which is supported by the NIH (P30-CA77598). KFMJ was supported by the National Science Foundation Graduate Research Fellowship Program.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The authors acknowledge Todd Rappe and the Minnesota NMR center for assistance with the ALARM NMR assay. The authors acknowledge Jayme Dahlin for technical assistance and thoughtful discussion. We thank Dr. Steve Geib (Pitt) for X-ray data collection and refinement.

■ ABBREVIATIONS

μ M, micromolar; ALARM MSPS, a La assay to detect reactive molecules by mass spectrometry peptide sequencing; APKR, allenic Pauson–Khand reaction; Bpin, boron pinacolate; Cs_2CO_3 , cesium carbonate; CDCl_3 , deuteriochloroform; CF_3 , trifluoromethyl; CN, cyano; CO, carbon monoxide; CPM, (N-[4-(7-diethylamino-4-methylcoumarin-3-yl)phenyl]-maleimide); CuH, copper hydride; dr, diastereomeric ratio; Et_2O , diethyl ether; EtOAc, ethyl acetate; HCl, hydrochloric acid; HEK, human embryonic kidney; IAD, iodoacetamide; K_2CO_3 , potassium carbonate; k_{inact} , inactivation rate constant; k_{off} , dissociation rate constant; k_{on} , association rate constant; KMnO_4 , potassium permanganate; MeLi, methyl lithium; mg, milligram; Na_2SO_3 , sodium sulfite; NaI, sodium iodide; NEt_3 , trimethylamine; NF- κ B, nuclear factor κ -light-chain-enhancer of activated B cells; ng, nanogram; NH_4Cl , ammonium chloride; NH_4OH , ammonium hydroxide; OMe, methoxy; Ph, phenyl; PTL, parthenolide; PPh_3 , triphenylphosphine; $[\text{Rh}(\text{CO})_2\text{Cl}]_2$, rhodium(bis)carbonylchloride dimer; ScOTf_3 , scandium(III) triflate; SEAP, secreted embryonic alkaline phosphatase; SiMe_3 , trimethylsilane; TIPS, triisopropylsilyl; TNF- α , tumor necrosis factor α

■ REFERENCES

- (1) Kitson, R. R. A.; Millemaggi, A.; Taylor, R. J. K. The Renaissance of α -Methylene- γ -Butyrolactones: New Synthetic Approaches. *Angew. Chem., Int. Ed.* **2009**, *48*, 9426–9451.
- (2) Simonsen, H.; Weitzel, C.; Christensen, S. Guaianolide Sesquiterpenoids: Pharmacology and Biosynthesis. In *Natural Products*; Ramawat, K. G.; Mérillon, J.-M., Eds.; Springer Berlin Heidelberg, 2013; pp 3069–3098.
- (3) Drew, D. P.; Krichau, N.; Reichwald, K.; Simonsen, H. T. Guaianolides in Apiaceae: Perspectives on Pharmacology and Biosynthesis. *Phytochem. Rev.* **2009**, *8*, 581–599.
- (4) Ghantous, A.; Gali-Muhtasib, H.; Vuorela, H.; Saliba, N. A.; Darwiche, N. What Made Sesquiterpene Lactones Reach Cancer Clinical Trials? *Drug Discovery Today* **2010**, *15*, 668–678.
- (5) Kempema, A. M.; Widen, J. C.; Hexum, J. K.; Andrews, T. E.; Wang, D.; Rathe, S. K.; Meece, F. A.; Noble, K. E.; Sachs, Z.; Largaespada, D. A.; Harki, D. A. Synthesis and Antileukemic Activities of C1–C10-Modified Parthenolide Analogues. *Bioorg. Med. Chem.* **2015**, *23*, 4737–4745.
- (6) Abdeldayem, A.; Raouf, Y. S.; Constantinescu, S. N.; Moriggl, R.; Gunning, P. T. Advances in Covalent Kinase Inhibitors. *Chem. Soc. Rev.* **2020**, *49*, 2617–2687.
- (7) Gehringer, M.; Laufer, S. A. Emerging and Re-Emerging Warheads for Targeted Covalent Inhibitors: Applications in Medicinal Chemistry and Chemical Biology. *J. Med. Chem.* **2019**, *62*, 5673–5724.
- (8) Dalton, S. E.; Campos, S. Covalent Small Molecules as Enabling Platforms for Drug Discovery. *ChemBioChem* **2020**, *21*, 1080–1100.
- (9) Lonsdale, R.; Ward, R. A. Structure-Based Design of Targeted Covalent Inhibitors. *Chem. Soc. Rev.* **2018**, *47*, 3816–3830.
- (10) Xu, J.; Chen, A.; Joy, J.; Xavier, V. J.; Ong, E. H. Q.; Hill, J.; Chai, C. L. L. Rational Design of Resorcylic Acid Lactone Analogues as Covalent MNK1/2 Kinase Inhibitors by Tuning the Reactivity of an Enamide Michael Acceptor. *ChemMedChem* **2013**, *8*, 1483–1494.
- (11) Xu, J.; Ong, E. H. Q.; Hill, J.; Chen, A.; Chai, C. L. L. Design, Synthesis and Biological Evaluation of FLT3 Covalent Inhibitors with a Resorcylic Acid Core. *Bioorg. Med. Chem.* **2014**, *22*, 6625–6637.
- (12) Cee, V. J.; Volak, L. P.; Chen, Y.; Bartberger, M. D.; Tegley, C.; Arvedson, T.; McCarter, J.; Tasker, A. S.; Fotsch, C. Systematic Study of the Glutathione (GSH) Reactivity of N-Arylacrylamides: 1. Effects of Aryl Substitution. *J. Med. Chem.* **2015**, *58*, 9171–9178.
- (13) Flanagan, M. E.; Abramite, J. A.; Anderson, D. P.; Aulabaugh, A.; Dahal, U. P.; Gilbert, A. M.; Li, C.; Montgomery, J.; Oppenheimer, S. R.; Ryder, T.; Schuff, B. P.; Uccello, D. P.; Walker, G. S.; Wu, Y.; Brown, M. F.; Chen, J. M.; Hayward, M. M.; Noe, M. C.; Obach, R. S.; Philippe, L.; Shanmugasundaram, V.; Shapiro, M. J.; Starr, J.; Stroth, J.; Che, Y. Chemical and Computational Methods for the Characterization of Covalent Reactive Groups for the Prospective Design of Irreversible Inhibitors. *J. Med. Chem.* **2014**, *57*, 10072–10079.
- (14) Jackson, P. A.; Widen, J. C.; Harki, D. A.; Brummond, K. M. Covalent Modifiers: A Chemical Perspective on the Reactivity of α,β -Unsaturated Carbonyls with Thiols via Hetero-Michael Addition Reactions. *J. Med. Chem.* **2017**, *60*, 839–885.
- (15) Martin, J. S.; MacKenzie, C. J.; Fletcher, D.; Gilbert, I. H. Characterising Covalent Warhead Reactivity. *Bioorg. Med. Chem.* **2019**, *27*, 2066–2074.
- (16) Konaklieva, M.; Plotkin, B. Lactones: Generic Inhibitors of Enzymes? *Mini-Rev. Med. Chem.* **2005**, *5*, 73–95.
- (17) Merfort, I. Perspectives on Sesquiterpene Lactones in Inflammation and Cancer. *Curr. Drug Targets* **2011**, *12*, 1560–1573.
- (18) Ma, G.-H.; Chen, K.-X.; Zhang, L.-Q.; Li, Y.-M. Advance in Biological Activities of Natural Guaiane-Type Sesquiterpenes. *Med. Chem. Res.* **2019**, *28*, 1339–1358.
- (19) Siedle, B.; García-Piñeres, A. J.; Murillo, R.; Schulte-Mönting, J.; Castro, V.; Rüngeler, P.; Klaas, C. A.; Da Costa, F. B.; Kisiel, W.; Merfort, I. Quantitative Structure–Activity Relationship of Sesquiterpene Lactones as Inhibitors of the Transcription Factor NF- κ B. *J. Med. Chem.* **2004**, *47*, 6042–6054.
- (20) Büchele, B.; Zugmaier, W.; Lunov, O.; Syrovets, T.; Merfort, I.; Simmet, T. Surface Plasmon Resonance Analysis of Nuclear Factor- κ B Protein Interactions with the Sesquiterpene Lactone Helenalin. *Anal. Biochem.* **2010**, *401*, 30–37.
- (21) Hayden, M. S.; Ghosh, S. Shared Principles in NF- κ B signaling. *Cell* **2008**, *132*, 344–362.
- (22) Chaturvedi, M. M.; Sung, B.; Yadav, V. R.; Kannappan, R.; Aggarwal, B. B. NF- κ B Activation and its Role in Cancer: ‘One Size Does Not Fit All’. *Oncogene* **2011**, *30*, 1615–1630.
- (23) Wen, B.; Hexum, J. K.; Widen, J. C.; Harki, D. A.; Brummond, K. M. A Redox Economical Synthesis of Bioactive 6,12-Guaianolides. *Org. Lett.* **2013**, *15*, 2644–2647.
- (24) Wells, S. M. The Synthesis of 6,12-Guaianolide Analogs and Related Chemical Tools for Probing their Mechanism of NF- κ B Inhibition. Ph.D. Thesis, University of Pittsburgh, 2016.
- (25) Grillet, F.; Huang, C.; Brummond, K. M. An Allenic Pauson–Khand Approach to 6,12-Guaianolides. *Org. Lett.* **2011**, *13*, 6304–6307.
- (26) Brummond, K. M.; Lu, J. A Short Synthesis of the Potent Antitumor Agent ($\pm$)-Hydroxymethylacetylfulvene Using an Allenic Pauson–Khand Type Cycloaddition. *J. Am. Chem. Soc.* **1999**, *121*, 5087–5088.
- (27) Tsuji, J.; Sugiura, T.; Minami, I. Preparation of 1,2-Dienes by the Palladium-Catalyzed Hydrogenolysis of 3-Methoxycarbonyloxy-1-alkynes with Ammonium Formate. *Synthesis* **1987**, 603–606.
- (28) Elford, T. G.; Hall, D. G. Imine Allylation Using 2-Alkoxy carbonyl Allylboronates as an Expedient Three-Component Reaction to Polysubstituted α -exo-Methylene- γ -Lactams. *Tetrahedron Lett.* **2008**, *49*, 6995–6998.
- (29) To obtain high yields consistently for this reaction, HMPA was distilled from CaH_2 , then stored over 4 Å molecular sieves, and chloromethyl pinacolboronate was freshly distilled.
- (30) Chataigner, I.; Zammattio, F.; Lebreton, J.; Villiéras, J. Enantioselective Addition of β -Functionalized Allylboronates to

Aldehydes and Aldimines. Stereocontrolled Synthesis of α -Methylene- γ -lactones and Lactams. *Tetrahedron* **2008**, *64*, 2441–2455.

(31) Elford, T. G.; Ulaczyk-Lesanko, A.; De Pascale, G.; Wright, G. D.; Hall, D. G. Diversity-Oriented Synthesis and Preliminary Biological Screening of Highly Substituted Five-Membered Lactones and Lactams Originating From an Allylboration of Aldehydes and Imines. *J. Comb. Chem.* **2009**, *11*, 155–168.

(32) Chataigner, I.; Zammattio, F.; Lebreton, J.; Villieras, J. Enantioselective Synthesis of α -Methylene- γ -Lactams. Nucleophilic Addition of a Chirally Modified β -Functionalized Allylboronate Reagent to Imines. *Synlett* **1998**, 275–276.

(33) Daeuble, J. F.; McGettigan, C.; Stryker, J. M. Selective Reduction of Alkynes to Cis-Alkenes by Hydrometallation Using $[(\text{Ph}_3\text{P})\text{CuH}]_6$. *Tetrahedron Lett.* **1990**, *31*, 2397–2400.

(34) Petasis, N. A.; Hu, Y.-H. Allenation of Carbonyl Compounds with Alkenyltitanocene Derivatives. *J. Org. Chem.* **1997**, *62*, 782–783.

(35) Tsuji, J.; Sugiura, T.; Yuhara, M.; Minami, I. Palladium-Catalysed Preparation of 1,2-Dienes by Selective Hydrogenolysis of Alk-2-ynyl Carbonates with Ammonium Formate. *J. Chem. Soc., Chem. Commun.* **1986**, 922–924.

(36) Liu, Z.; Liao, P.; Bi, X. Lewis and Bronsted Acid Cocatalyzed Reductive Deoxygenation of Propargylic Alcohols with 2-Nitrobenzenesulfonylhydrazide. *Chem.—Eur. J.* **2014**, *20*, 17277–17281.

(37) Yin, J.; Buchwald, S. L. Pd-Catalyzed Intermolecular Amidation of Aryl Halides: The Discovery that Xantphos can be Trans-Chelating in a Palladium Complex. *J. Am. Chem. Soc.* **2002**, *124*, 6043–6048.

(38) Klapars, A.; Huang, X.; Buchwald, S. L. A General and Efficient Copper Catalyst for the Amidation of Aryl Halides. *J. Am. Chem. Soc.* **2002**, *124*, 7421–7428.

(39) Wells, S. M.; Brummond, K. M. Conditions for a Rh(I)-catalyzed [2+2+1] Cycloaddition Reaction with Methyl Substituted Allenes and Alkynes. *Tetrahedron Lett.* **2015**, *56*, 3546–3549.

(40) Kallepalli, V. A.; Shi, F.; Paul, S.; Onyeozili, E. N.; Maleczka, R. E.; Smith, M. R. Boc Groups as Protectors and Directors for Ir-Catalyzed C–H Borylation of Heterocycles. *J. Org. Chem.* **2009**, *74*, 9199–9201.

(41) Petri, L.; Ábrányi-Balogh, P.; Varga, P. R.; Imre, T.; Keserű, G. M. Comparative Reactivity Analysis of Small-Molecule Thiol Surrogates. *Bioorg. Med. Chem.* **2020**, *28*, 115357.

(42) Avonto, C.; Tagliatalata-Scafati, O.; Pollastro, F.; Minassi, A.; Di Marzo, V.; De Petrocellis, L.; Appendino, G. An NMR Spectroscopic Method to Identify and Classify Thiol-Trapping Agents: Revival of Michael Acceptors for Drug Discovery. *Angew. Chem., Int. Ed.* **2011**, *50*, 467–471.

(43) Spartan'16 MMFF was used to determine the lowest energy conformer and equilibrium geometries were established using DFT calculations (B3LYP) and 6-31G* basis set. All calculations were performed in the gas phase.

(44) Castañeda-Acosta, J.; Fischer, N. H.; Vargas, D. Biomimetic Transformations of Parthenolide. *J. Nat. Prod.* **1993**, *56*, 90–98.

(45) Widen, J. C.; Kempema, A. M.; Villalta, P. W.; Harki, D. A. Targeting NF- κ B p65 with a Helenalin Inspired Bis-electrophile. *ACS Chem. Biol.* **2017**, *12*, 102–113.

(46) Rana, S.; Blowers, E. C.; Tebbe, C.; Contreras, J. I.; Radhakrishnan, P.; Kizhake, S.; Zhou, T.; Rajule, R. N.; Arnst, J. L.; Munkarah, A. R.; Rattan, R.; Natarajan, A. Isatin Derived Spirocyclic Analogues with α -Methylene- γ -Butyrolactone as Anticancer Agents: A Structure–Activity Relationship Study. *J. Med. Chem.* **2016**, *59*, S121–S127.

(47) García-Piñeres, A. J.; Lindenmeyer, M. T.; Merfort, I. Role of Cysteine Residues of p65/NF- κ B on the Inhibition by the Sesquiterpene Lactone Parthenolide and N-Ethyl Maleimide, and on its Transactivating Potential. *Life Sci.* **2004**, *75*, 841–856.

(48) Pande, V.; Sousa, S.; Ramos, M. Direct Covalent Modification as a Strategy to Inhibit Nuclear Factor- κ B. *Curr. Med. Chem.* **2009**, *16*, 4261–4273.

(49) Widen, J. C.; Kempema, A. M.; Baur, J. W.; Skopec, H. M.; Edwards, J. T.; Brown, T. J.; Brown, D. A.; Meece, F. A.; Harki, D. A. Helenalin Analogues Targeting NF- κ B p65: Thiol Reactivity and

Cellular Potency Studies of Varied Electrophiles. *ChemMedChem* **2018**, *13*, 303–311.

(50) Kwok, B. H. B.; Koh, B.; Ndubuisi, M. I.; Elofsson, M.; Crews, C. M. The Anti-Inflammatory Natural Product Parthenolide from the Medicinal Herb Feverfew Directly Binds to and Inhibits I κ B Kinase. *Chem. Biol.* **2001**, *8*, 759–766.

(51) Freund, R. R. A.; Gobrecht, P.; Fischer, D.; Arndt, H.-D. Advances in Chemistry and Bioactivity of Parthenolide. *Nat. Prod. Rep.* **2020**, *37*, 541–565.

(52) Lindenmeyer, M. T.; García-Piñeres, A. J.; Castro, V.; Merfort, I. Sesquiterpene Lactones Inhibit Luciferase but not β -Galactosidase Activity in vitro and ex vivo. *Anal. Biochem.* **2004**, *328*, 147–154.

(53) Hexum, J. K.; Tello-Aburto, R.; Struntz, N. B.; Harned, A. M.; Harki, D. A. Bicyclic Cyclohexenones as Inhibitors of NF- κ B Signaling. *ACS Med. Chem. Lett.* **2012**, *3*, 459–464.

(54) Moraski, G. C.; Markley, L. D.; Hipskind, P. A.; Boshoff, H.; Cho, S.; Franzblau, S. G.; Miller, M. J. Advent of Imidazo[1,2-a]pyridine-3-carboxamides with Potent Multi- and Extended Drug Resistant Antituberculosis Activity. *ACS Med. Chem. Lett.* **2011**, *2*, 466–470.

(55) Li, X.-C.; Babu, K. S.; Jacob, M. R.; Khan, S. I.; Agarwal, A. K.; Clark, A. M. Natural Product-Based 6-Hydroxy-2,3,4,6-tetrahydropyrrolo[1,2-a]pyrimidinium Scaffold as a New Antifungal Template. *ACS Med. Chem. Lett.* **2011**, *2*, 391–395.

(56) Huth, J. R.; Mendoza, R.; Olejniczak, E. T.; Johnson, R. W.; Cothron, D. A.; Liu, Y.; Lerner, C. G.; Chen, J.; Hajduk, P. J. ALARM NMR: A Rapid and Robust Experimental Method to Detect Reactive False Positives in Biochemical Screens. *J. Am. Chem. Soc.* **2005**, *127*, 217–224.

(57) Huth, J. R.; Song, D.; Mendoza, R. R.; Black-Schaefer, C. L.; Mack, J. C.; Dorwin, S. A.; Lador, U. S.; Severin, J. M.; Walter, K. A.; Bartley, D. M.; Hajduk, P. J. Toxicological Evaluation of Thiol-Reactive Compounds Identified Using A La Assay to Detect Reactive Molecules by Nuclear Magnetic Resonance. *Chem. Res. Toxicol.* **2007**, *20*, 1752–1759.

(58) Dahlin, J. L.; Cuellar, M.; Singh, G.; Nelson, K. M.; Strasser, J. M.; Rappe, T.; Xia, Y.; Veglia, G.; Walters, M. A. ALARM NMR for HTS Triage and Chemical Probe Validation. *Curr. Protoc. Chem. Biol.* **2018**, *10*, 91–117.

(59) Potter, Z. E.; Lau, H.-T.; Chakraborty, S.; Fang, L.; Guttman, M.; Ong, S.-E.; Fowler, D. M.; Maly, D. J. Parallel Chemoselective Profiling for Mapping Protein Structure. *Cell Chem. Biol.* **2020**, *27*, 1084–1096.

(60) Görl, C.; Alt, H. G. The Combination of Mononuclear Metallocene and Phenoxyimine Complexes to Give Trinuclear Catalysts for the Polymerization of Ethylene. *J. Organomet. Chem.* **2007**, *692*, S727–S753.

(61) Abidi, S. L. Direct Conversion of Terpenylalkanolamines to Ethylidene N-Nitroso Compounds. *J. Org. Chem.* **1986**, *51*, 2687–2694.

(62) Balog, A.; Curran, D. P. Ring-Enlarging Annulations. A One-Step Conversion of Cyclic Silyl Acyloins and ω -Alkynyl Acetals to Polycyclic Enediones. *J. Org. Chem.* **1995**, *60*, 337–344.

(63) Whiting, A. A Convenient Preparation of β -boronate Carbonyl Derivatives. Evidence For the Intervention of Boronate “ate”-Complexes in Enolate Alkylations. *Tetrahedron Lett.* **1991**, *32*, 1503–1506.

(64) Grillet, F.; Huang, C.; Brummond, K. M. An Allenic Pauson-Khand Approach to 6,12-Guaianolides. *Org. Lett.* **2011**, *13*, 6304–6307.

(65) Serra, S.; Fuganti, C. Benzannulation of Substituted 3-Alkoxy-carbonylhex-3-en-5-ynoic Acids: A New Route to 4-Substituted 3,5-Dihydroxybenzoic Acids Derivatives. *Synlett* **2002**, 1661–1664.

(66) Fatiadi, A. J. Active Manganese Dioxide Oxidation in Organic Chemistry - Part I. *Synthesis* **1976**, 65–104.

(67) Kern, N.; Hoffmann, M.; Blanc, A.; Weibel, J.-M.; Pale, P. Gold(I)-Catalyzed Rearrangement of N-Aryl 2-Alkynylazetidines to Pyrrolo[1,2-a]indoles. *Org. Lett.* **2013**, *15*, 836–839.

- (68) Lee, D.-w.; Yun, J. Direct Synthesis of Stryker's Reagent from a Cu(II) Salt. *Tetrahedron Lett.* **2005**, *46*, 2037–2039.
- (69) Carlson, T. L.; Moini, M.; Eckenrode, B. A.; Allred, B. M.; Donfack, J. Protein Extraction from Human Anagen Head Hairs 1-Millimeter or Less in Total Length. *BioTechniques* **2018**, *64*, 170–176.
- (70) MacLean, B.; Tomazela, D. M.; Shulman, N.; Chambers, M.; Finney, G. L.; Frewen, B.; Kern, R.; Tabb, D. L.; Liebler, D. C.; MacCoss, M. J. Skyline: An Open Source Document Editor for Creating and Analyzing Targeted Proteomics Experiments. *J. Bioinf.* **2010**, *26*, 966–968.