

Photoreaction of 3-Azido-5,7-di-*t*-butyl-2,1-benzisothiazole.
Spectroscopic Detection of a Thionitroso Intermediate

Renji OKAZAKI,^{*} Masaki TAKAHASHI, Naoki INAMOTO,
Tadashi SUGAWARA,^{†,††} and Hiizu IWAMURA^{†,†††}


Department of Chemistry, Faculty of Science,
The University of Tokyo, Hongo, Tokyo 113

[†] Division of Applied Molecular Science,
Institute of Molecular Science, Okazaki 444

Photoreaction of the title compound at cryogenic temperatures produced a highly reactive 2,4-di-*t*-butyl-6-cyanothionitrosobenzene which showed a visible absorption around 470 nm.

Although thiocarbonyl compounds have extensively been studied in recent years, our knowledge on thionitroso compounds, the corresponding nitrogen analog, is relatively scarce.¹⁾ In connection with our continuing work on nitrogen-sulfur compounds,²⁾ we report here on the spectroscopic detection (UV-VIS and IR) of a thionitroso compound using the photoreaction of azidoisothiazole **1**. Although Joucla and Rees reported the generation of a thionitroso intermediate in the photoreaction of azidoisothiazole derivatives, its intermediacy was confirmed only by chemical trapping.^{1a)}

Compound **1** was synthesized from **2**^{2b)} by amination,³⁾ diazotization and subsequent reaction with sodium azide (20% from **2**).⁴⁾




(a) *n*-BuLi. (b) $\text{N}_3\text{CH}_2\text{SiMe}_3$. (c) H_3O^+ . (d) $\text{NaNO}_2\text{-HCl}$. (e) NaN_3 .

^{††} Present address: Department of Pure and Applied Sciences, Faculty of Arts and Sciences, The University of Tokyo, Komaba, Tokyo 153.


^{†††} Present address: Department of Chemistry, Faculty of Science, The University of Tokyo, Hongo, Tokyo 113.

Irradiation of a hexane solution of **1** with a high pressure Hg lamp at 0 °C gave **3** (78%) and **4** (16%).^{4,5)}



When the photoreaction (≥ 320 nm) of **1** in an argon matrix at 12 K was followed by electronic spectroscopy, a new band was observed around 470 nm (456 and 483 nm) at the expense of a strong 353 nm absorption of **1** (Fig. 1a). Monitoring by IR spectroscopy of the reaction under similar conditions revealed appearance of a band due to a cyano group (2225 cm^{-1}) along with concomitant decrease of the azide absorption (2130 cm^{-1}) (Fig. 1b). This strongly suggests that the initial intermediate, which has an absorption around 470 nm (ϵ 4700), can be assigned to thionitroso compound **5**, excluding the possibility of nitrene **6**. The band around 470 nm was also observed in the photoreactions in organic glass (70–80 K) such as EPA (ether:pentane:ethanol 5:5:2), 2-methyltetrahydrofuran, 3-methylpentane, and isopentane, although it disappeared above 90 K, suggesting the high reactivity of **5**.

These observations lead us to the conclusion that the photoreaction of **1** most likely proceeds via the direct formation of thionitrosobenzene **5** from **1** without the intermediacy of nitrene **6** because of the unique structure of **1** with a probable great resonance contribution of **1'**.






Fig. 1. Photolysis of 1 in argon matrix (12 K) monitored by electronic (a) and IR (b) spectroscopies.

Previously, Pedersen et al. reported that the photolysis of benzo[c]-1,2,5-thiadiazole 2-oxide resulted in the reversible formation of 2-thionitroso-nitrosobenzene to which they assigned a visible absorption at 485 nm (ϵ 8000-9000) (in an argon matrix at 20 K).⁶⁾ This interpretation was later questioned, however, by Mayer et al. who carried out extensive calculations on the electronic absorption of thionitroso compounds as well as related sulfur containing species.⁷⁾

We believe that close similarity in the position and coefficient of the absorptions between Pedersen's compound and ours strongly suggests that thionitrosobenzene derivatives have a rather strong absorption band around 470-490 nm in contrast to Mayer's calculations.

References

- 1) a) M. F. Joucla and C. W. Rees, *J. Chem. Soc., Chem. Commun.*, **1984**, 374; b) M. R. Bryce and P. C. Taylor, *Tetrahedron Lett.*, **30**, 3835 (1989) and references cited in these papers.
- 2) a) Y. Inagaki, R. Okazaki, and N. Inamoto, *Bull. Chem. Soc. Jpn.*, **52**, 1998 (1979); b) Y. Inagaki, R. Okazaki, and N. Inamoto, *ibid.*, **52**, 2002 (1979); c) S. Nakamura, M. Takahashi, R. Okazaki, and K. Morokuma, *J. Am. Chem. Soc.*, **109**, 4142 (1987); d) R. Okazaki, M. Unno, and N. Inamoto, *Chem. Lett.*, **1987**, 2293.
- 3) K. Nishiyama and N. Tanaka, *J. Chem. Soc., Chem. Commun.*, **1983**, 1322.
- 4) **1**: mp 77-79 °C(dec); ¹H NMR(CDCl₃) δ 1.21(s, 9H), 1.44(s, 9H), 7.17(ABq, 2H, $\Delta\delta=0.08$, J=1.9 Hz); UV-VIS(hexane) $\lambda_{\max}$ 228 nm (ϵ 18400), 255(sh, 6890), 297(4280), 312(4260), 356(7720); high MS 288.1403(calcd for C₁₅H₂₀N₄S 288.1408). **3**: mp 175.0-175.5 °C; ¹H NMR(CDCl₃) δ 1.17(s, 36H), 7.38(ABq, $\Delta\delta=0.15$, J=2.4 Hz); UV-VIS(pentane) $\lambda_{\max}$ 284 nm (ϵ 10300), 416(5400); high MS 488.2954(calcd for C₃₀H₄₀N₄S 488.2972). **4**: mp 80.0-80.5 °C; ¹H NMR(CDCl₃) δ 1.13(s, 9H), 1.28(s, 9H), 4.33(bs, 2H), 7.15(ABq, 2H, $\Delta\delta=0.19$, J=2.0 Hz); UV-VIS(pentane) $\lambda_{\max}$ 246 nm (ϵ 6300), 319(4000); high MS 230.1773(calcd for C₁₅H₂₂N₂ 230.1781).
- 5) Products **3** and **4** were isolated by silica gel chromatography (hexane-ether 5:1) and recrystallization.
- 6) C. L. Pedersen, C. Lohse, and M. Poliakoff, *Acta Chem. Scand., Ser. B*, **32**, 625 (1978).
- 7) A. Mehlhorn, J. Sauer, J. Fabian, and R. Mayer, *Phosphorus and Sulfur*, **11**, 325 (1981).

(Received September 12, 1989)