

Diels–Alder Intramolecular Cycloaddition of Vinylallenyl Cycloalken-3-yl Ethers

J. P. Dulcère, V. Agati and R. Faure

Faculté des Sciences et Techniques de St Jérôme, Laboratoire de Synthèse Organique URA CNRS 1411, av. Esc. Normandie-Niemen, D 12, F-13397-Marseille Cedex 20, France

Cohalogenation of cycloalkenes **1** by *N*-bromosuccinimide (NBS) in 4-methylpent-4-en-2-yn-1-ol affords β -bromopent-4-en-2-ynyl ethers **2** which undergo facile base-promoted consecutive dehydrohalogenation–isomerization–intramolecular Diels–Alder cycloaddition leading to tricyclic conjugated enol ethers **6**.

The ability of vinylallenes to participate as dienes in Diels–Alder type cycloadditions¹ constitutes a convenient route for the construction of complex ring systems.

Although intramolecular cycloadditions of allenyl ether dienophiles have been well studied,² reactions involving vinyl allenyl ether dienes have received little attention.³ Our interest in the reactivity of allyl allenyl ethers⁴ has prompted us to prepare and study the intramolecular cycloaddition of a series of allyl vinylallenyl ethers **5**.


It has been shown that allyl-4-methylpent-4-en-2-ynyl ethers do not undergo base-catalysed cycloadditions;³ we report here the synthesis and the extra reactivity of the cyclic analogues **3**, which, under the same base-catalysed reaction conditions, lead to tricyclic conjugated enol ethers **6** by consecutive isomerization [4 + 2] Diels–Alder reaction.

Halogenation of cycloalkenes **1** by *N*-bromosuccinimide (NBS) in 4-methylpent-4-en-2-yn-1-ol afforded the β -bromopent-4-en-2-ynyl ethers **2** in 70–75% yield, the reaction being

regio- and stereo-selective.[†] Treating **2** with 1.3 equiv. of Bu^tOK in refluxing benzene for 2 h resulted in the formation of **6**‡ in 80–85% yield; dehydrobromination of **2** in diazabicycloundecene (DBU, 3 equiv. 60 °C, 3 h) gave rise to allyl pent-4-en-2-ynyl ethers **3** (72–97%), which undergo catalytic isomerization (Bu^tOK, 0.3 equiv., in refluxing pentane) into enol ethers **6** (90–95%). On the other hand, treatment of **2a, b** under the same catalytic conditions resulted in the isomeriza-

[†] Satisfactory analytical and spectral data were obtained for all compounds.

‡ **6b**: ¹H NMR (CDCl₃) (200 MHz) δ 6.16 (s, br, 1H); 5.90 (s, br, 1H); 4.46 (q, *J* 7.0 Hz, 1H); 2.96 (t, br, *J* 7.0 Hz, 1H); 2.15 (m, 1H); 1.8–1.7 (m, 2H); 1.65 (s, 3H); 1.62 (m, 1H); 1.45 (m, 1H); 1.33 (m, 1H); 1.2–1.1 (m, 2H); 0.85 (m, 1H). ¹³C NMR (50.32 MHz) 137.72, 130.65, 114.84, 114.78, 80.53, 42.42, 37.13, 31.49, 28.41, 27.92, 24.05, 21.64. IR ν /cm^{–1} 3100, 2995, 1650, 1610, 1070.



Scheme 1 Reagents and conditions: i, $H_2C=C(Me)C\equiv CCH_2OH$, NBS ($-15^\circ C$, 3 h); ii, Bu^tOK (1.3 equiv.), benzene, reflux, 2 h; iii, DBU (3 equiv., $110^\circ C$, 3 h); iv, Bu^tOK (0.3 equiv., pentane, reflux, 0.5–2 h); v, MeOH, camphorsulfonic acid (CSA) 3%, 2 h; vi, Jones' reagent

tion of the acetylenic unit,⁵ leading to β -bromovinyl allenyl ethers **4a, b** (65–67%) and then to **6a, b** (78–82%), upon treatment with DBU. This base-catalysed isomerization into a vinylallenyl moiety accounts for the clean intramolecular cycloaddition of **3**. The structures of $[4 + 2]$ cycloaddition adducts **6** were confirmed by selective conversion into acetals **7** and the corresponding polycyclic fused lactones **8**, the structures of which constitute the framework of a number of natural terpenes.⁶

In conclusion, the base-promoted consecutive dehydrobromination–isomerization–intramolecular cycloaddition of readily available derivatives **2** provides a selective means for the functionalization of cycloalkenes **1** into tricyclic structures **6** with high synthetic potential.

Received, 9th November 1992; Com. 2/05958D

References

- 1 M. Bertrand, J. Grimaldi and B. Waegell, *Bull. Soc. Chim. Fr.*, 1971, 962; E. A. Deutsch and B. B. Snider, *J. Org. Chem.*, 1982, **47**, 2682; K. Hayakawa, M. Yodo, S. Ohsuki and K. Kanematsu, *J. Am. Chem. Soc.*, 1984, **106**, 6735.
- 2 K. Hayakawa, K. Aso, M. Shiro and K. Kanematsu, *J. Am. Chem. Soc.*, 1989, **111**, 5312 and references cited therein.
- 3 A. J. Bartlett, T. Laird and W. D. Ollis, *J. Chem. Soc., Perkin Trans. 1*, 1975, 1315; *J. Chem. Soc., Chem. Commun.*, 1974, 496.
- 4 J. P. Dulcère, M. N. Mihoubi and J. Rodriguez, *J. Chem. Soc., Chem. Commun.*, 1988, 237; J. P. Dulcère and J. K. Crandall, *J. Chem. Soc., Chem. Commun.*, 1990, 561.
- 5 J. P. C. M. van Dongen, A. J. De Jong, W. A. Selling, P. P. Montijn, J. H. van Boom and L. Brandsma, *Recl. Trav. Chim. Pays-Bas*, 1967, **86**, 1077; H. A. Selling, J. A. Rompes, P. P. Montijn, S. Hoff, J. H. van Boom, L. Brandsma and J. F. Arens, *Recl. Trav. Chim. Pays-Bas*, 1969, **88**, 119; L. Brandsma and H. D. Verkrujine, in *Studies in Organic Chemistry 8, Synthesis of Acetylenes, Allenes and Cumulenes*, Elsevier, 1981, p. 82.
- 6 K. Hayakawa, S. Ohsuki and K. Kanematsu, *Tetrahedron Lett.*, 1986, **27**, 947; S. Hashimoto, S. Sakata, M. Sonogawa and S. Ikegami, *J. Am. Chem. Soc.*, 1988, **110**, 3670; E. J. Corey, P. D. S. Jardine and J. C. Rohloff, *J. Am. Chem. Soc.*, 1988, **110**, 3672; K. Kanematsu and S. Nagashima, *J. Chem. Soc., Chem. Commun.*, 1989, 1028; M. J. Begley, D. R. Cheshire, T. Harrison, J. H. Hutchinson, P. L. Myers and G. Pattenden, *Tetrahedron*, 1989, **45**, 5215; M. E. Jung and C. N. Zimmerman, *J. Am. Chem. Soc.*, 1991, **113**, 7813; E. Kitazawa, A. Sato, S. Takahashi, H. Kuwano and A. Ogiso, *Chem. Pharm. Bull.*, 1980, **28**, 227; J. R. Hanson, in *Terpenoids and Steroids*, RSC, 1983, vol. 12.