

*Acta Cryst.* (1994). **C50**, 1812–1814

## 5-Phenyl-3-oxa-4-azatricyclo[5.2.1.0<sup>2,6</sup>]-dec-4-ene

A. THIRUVALLUVAR AND V. PARTHASARATHI\*

*Department of Physics, Bharathidasan University,  
Tiruchirapalli 620 024, India*

A. NAGARAJAN AND M. KRISHNA PILLAY

*Department of Chemistry, Bharathidasan University,  
Tiruchirapalli 620 024, India*


(Received 16 August 1993; accepted 28 January 1994)

### Abstract

A molecule of the title compound, C<sub>14</sub>H<sub>15</sub>NO, consists of an isoxazole ring fused to norbornane. The conformation of the five-membered isoxazole ring is nearly planar and its fusion onto the norbornane moiety is *exo*.

### Comment

Isoxazolines have become versatile intermediates in the synthesis of a variety of complex biomolecules as a result of their relatively facile ring cleavage to afford  $\beta$ -hydroxy ketones and  $\gamma$ -amino alcohols (Curran, 1988; Torssell, 1988; Caramella & Grunanger, 1984; Lang & Lin, 1984). Nitrile oxide cycloaddition with alkenes is a simple method of obtaining isoxazolines (Grundmann & Grunanger, 1971). A view of the title compound, (I), is given in Fig. 1.



5-Phenyl-3-oxa-4-azatricyclo[5.2.1.0<sup>2,6</sup>]-dec-4-ene forms a special type of compound since the bicyclic moiety (norbornane unit) can be fused to the isoxazole ring in two different ways, either in an *exo* or an *endo* fashion. In order to know which type of fusion has taken place, proton NMR spectra for the title compound were obtained.

The title compound was found to possess an *exo*-fused isoxazoline ring from analysis of the coupling




Fig. 1. Structure of the molecule showing the atomic numbering scheme.

constants of the H2 and H6 protons (Fig. 1); the signals were found to be doublets with  $J_{2,6} = 8.2$  Hz. The value for *exo-exo* vicinal coupling in the norbornane moiety is 12 Hz, while the value for *endo-endo* vicinal coupling is 9 Hz. This serves to indicate that the fusion of the isoxazoline ring onto norbornane is *exo*, leading to an *endo* orientation of the H2 and H6 protons. According to the Karplus relationship (Kemp, 1986), the dihedral angle corresponding to 8.2 Hz is near 0°. From the X-ray structure analysis of the title compound, the dihedral angle for H2—C2—C6—H6 is found to be 5.6 (4)°, thereby the *exo* orientation of the isoxazole to norbornane is confirmed.

From the <sup>13</sup>C NMR data, it was found that the C=N values were shifted upfield (~4 p.p.m.) when the substituent attached to the C=N double bond was changed from methyl to phenyl (Nagarajan & Krishna Pillay, 1993a). This means that when an aryl group is present at the oximino linkage, there is an effective conjugation. This would require the phenyl ring to be coplanar with the isoxazole ring. The phenyl ring makes an angle of 4.6 (1)° with the isoxazole ring and this enables both groups to establish conjugation. It was found that a weak C—H...N intramolecular interaction exists between the proton on C52 of the phenyl and atom N4 [H...N = 2.504 (3) Å], which contributes to the stabilization of the planar conformation. Dapporto, Paoli, Brandi, De Sarlo, Goti & Guarna (1992) have also reported the presence of conjugation in a similar situation.

Although the isoxazoline ring has been shown to possess a puckered conformation analogous to pyra-

zoline and 2,3-dihydrofuran on the basis of NMR spectral characteristics (Nagarajan & Krishna Pillay, 1993*b*), it adopts a nearly planar conformation in the title compound [puckering parameters:  $q_2 = 0.044(3) \text{ \AA}$ ,  $\varphi_2 = 145(4)^\circ$ ; C2 deviates  $0.072(3) \text{ \AA}$  from the least-squares plane formed by O3, N4, C5, C6]. The fusion of the isoxazole to norbornane prevents it from achieving a puckered conformation.

The structure of the norbornane moiety may be viewed as two interlocked five-membered rings. C10 deviates  $-0.851(3)$  and  $-0.853(3) \text{ \AA}$  from the least-squares planes of C7, C8, C9 and C1 and C1, C2, C6 and C7, respectively, thus constituting the common flap of the two envelopes. Similar values of this deviation [ $0.86(1)$  and  $0.84(1) \text{ \AA}$ ] have been reported for the norbornane moiety in 6,6-dimethyl-1-phenyl-7-thiatricyclo[3.2.1.1<sup>3,8</sup>]nonane 7,7-dioxide (Estienne, Pepe, Siri, Baldy & Samat, 1992). In the title compound, the norbornane moiety is found to be rigid, as expected.

## Experimental

The title compound was obtained, employing Torssell's one-pot synthesis (Larsen & Torssell, 1984), by the cycloaddition of norbornene with benzonitrile oxide. *N*-Chlorosuccinimide (2.66 g, 0.02 mol) and dichloromethane (75 ml) were added to a round-bottomed flask, fitted with a reflux condenser. To the suspension, pyridine (1 ml) and benzaldoxime (2.42 g, 0.02 mol) were added. The mixture was heated under reflux for 1 h and allowed to cool. Norbornene (1.88 g, 0.02 mol), in 10 ml of  $\text{CH}_2\text{Cl}_2$ , and triethylamine (2 ml in 5 ml of  $\text{CH}_2\text{Cl}_2$ ) were added and heated under reflux for 1 h. The mixture, after being allowed to cool, was washed with water (to remove succinimide, pyridine.HCl) and dried over anhydrous  $\text{Na}_2\text{SO}_4$ . Concentration *in vacuo* yielded the title compound (70%). Recrystallization from ethanol afforded colourless crystals, m.p. 382 K. Density  $D_m$  was measured by flotation in aqueous KBr.

## Crystal data

$\text{C}_{14}\text{H}_{15}\text{NO}$   
 $M_r = 213.3$   
 Monoclinic  
 $P2_1/n$   
 $a = 12.456(8) \text{ \AA}$   
 $b = 6.279(2) \text{ \AA}$   
 $c = 14.409(3) \text{ \AA}$   
 $\beta = 92.19(3)^\circ$   
 $V = 1126.1(8) \text{ \AA}^3$   
 $Z = 4$   
 $D_x = 1.26 \text{ Mg m}^{-3}$   
 $D_m = 1.27 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation  
 $\lambda = 0.71069 \text{ \AA}$   
 Cell parameters from 22 reflections  
 $\theta = 10.1\text{--}18.6^\circ$   
 $\mu = 0.07 \text{ mm}^{-1}$   
 $T = 298 \text{ K}$   
 Plate  
 $0.3 \times 0.25 \times 0.1 \text{ mm}$   
 Colourless

## Data collection

Enraf–Nonius CAD-4 diffractometer  
 $\omega$ – $2\theta$  scans  
 Absorption correction: none

$R_{\text{int}} = 0.04$   
 $\theta_{\text{max}} = 25^\circ$   
 $h = 0 \rightarrow 14$   
 $k = 0 \rightarrow 7$   
 $l = -17 \rightarrow 17$

2328 measured reflections  
 1488 independent reflections  
 1368 observed reflections  
 $[F > 6\sigma(F)]$

2 standard reflections monitored every 100 reflections  
 intensity variation: none

## Refinement

Refinement on  $F$   
 $R = 0.076$   
 $wR = 0.073$   
 $S = 2.49$   
 1368 reflections  
 145 parameters  
 H-atom parameters not refined

$w = 1/[\sigma^2(F) + 0.001F^2]$   
 $(\Delta/\sigma)_{\text{max}} = 0.001$   
 $\Delta\rho_{\text{max}} = 0.43 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.29 \text{ e \AA}^{-3}$   
 Extinction correction: none  
 Atomic scattering factors from SHELX76 (Sheldrick, 1976)

Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

$$U_{\text{eq}} = (1/3)\sum_i \sum_j U_{ij} a_i^* a_j^* a_i \cdot a_j.$$

|     | <i>x</i>    | <i>y</i>    | <i>z</i>   | $U_{\text{eq}}$ |
|-----|-------------|-------------|------------|-----------------|
| C1  | 0.3484 (2)  | −0.0519 (5) | 1.0109 (2) | 0.060 (1)       |
| C2  | 0.2337 (2)  | −0.0282 (5) | 0.9764 (2) | 0.050 (1)       |
| O3  | 0.2068 (2)  | 0.1926 (3)  | 0.9547 (1) | 0.064 (1)       |
| N4  | 0.1262 (2)  | 0.2636 (4)  | 1.0129 (2) | 0.055 (1)       |
| C5  | 0.1043 (2)  | 0.1157 (4)  | 1.0706 (2) | 0.042 (1)       |
| C6  | 0.1676 (2)  | −0.0831 (4) | 1.0605 (2) | 0.041 (1)       |
| C7  | 0.2564 (2)  | −0.1239 (4) | 1.1369 (2) | 0.051 (1)       |
| C8  | 0.3073 (3)  | −0.3364 (5) | 1.1153 (2) | 0.065 (1)       |
| C9  | 0.3683 (3)  | −0.2896 (6) | 1.0269 (3) | 0.072 (1)       |
| C10 | 0.3429 (2)  | 0.0317 (5)  | 1.1100 (2) | 0.061 (1)       |
| C51 | 0.0224 (2)  | 0.1526 (4)  | 1.1402 (2) | 0.044 (1)       |
| C52 | −0.0263 (3) | 0.3501 (5)  | 1.1494 (2) | 0.062 (1)       |
| C53 | −0.1024 (3) | 0.3822 (6)  | 1.2142 (3) | 0.077 (1)       |
| C54 | −0.1305 (3) | 0.2181 (8)  | 1.2717 (3) | 0.082 (1)       |
| C55 | −0.0853 (3) | 0.0217 (7)  | 1.2633 (2) | 0.072 (1)       |
| C56 | −0.0076 (2) | −0.0128 (5) | 1.1976 (2) | 0.055 (1)       |

Table 2. Selected geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

|              |            |               |           |
|--------------|------------|---------------|-----------|
| C1—C2        | 1.502 (4)  | C7—C8         | 1.515 (4) |
| C1—C9        | 1.529 (5)  | C7—C10        | 1.516 (4) |
| C1—C10       | 1.525 (4)  | C8—C9         | 1.536 (5) |
| C2—O3        | 1.457 (4)  | C51—C52       | 1.389 (4) |
| C2—C6        | 1.530 (4)  | C51—C56       | 1.388 (4) |
| O3—N4        | 1.405 (4)  | C52—C53       | 1.371 (5) |
| N4—C5        | 1.283 (4)  | C53—C54       | 1.376 (6) |
| C5—C6        | 1.487 (4)  | C54—C55       | 1.363 (7) |
| C5—C51       | 1.476 (4)  | C55—C56       | 1.396 (4) |
| C6—C7        | 1.552 (4)  |               |           |
| C9—C1—C10    | 102.0 (3)  | C6—C7—C10     | 101.9 (2) |
| C2—C1—C10    | 101.5 (2)  | C6—C7—C8      | 107.0 (2) |
| C2—C1—C9     | 107.1 (3)  | C8—C7—C10     | 102.0 (2) |
| C1—C2—C6     | 104.4 (2)  | C7—C8—C9      | 103.1 (3) |
| C1—C2—O3     | 111.9 (2)  | C1—C9—C8      | 103.3 (3) |
| O3—C2—C6     | 104.9 (2)  | C1—C10—C7     | 94.3 (2)  |
| C2—O3—N4     | 109.7 (2)  | C5—C51—C56    | 119.8 (3) |
| O3—N4—C5     | 109.4 (2)  | C5—C51—C52    | 121.4 (3) |
| N4—C5—C51    | 119.8 (2)  | C52—C51—C56   | 118.9 (3) |
| N4—C5—C6     | 114.7 (3)  | C51—C52—C53   | 120.8 (3) |
| C6—C5—C51    | 125.5 (2)  | C52—C53—C54   | 119.9 (4) |
| C2—C6—C5     | 101.1 (2)  | C53—C54—C55   | 120.6 (4) |
| C5—C6—C7     | 115.8 (2)  | C54—C55—C56   | 120.0 (4) |
| C2—C6—C7     | 102.0 (2)  | C51—C56—C55   | 119.8 (3) |
| C9—C1—C10—C7 | −54.2 (3)  | C51—C5—C6—C2  | 177.9 (3) |
| C2—C1—C10—C7 | 56.2 (3)   | C6—C5—C51—C52 | 174.5 (3) |
| C10—C1—C9—C8 | 32.9 (13)  | C6—C5—C51—C56 | −5.9 (4)  |
| C2—C1—C9—C8  | −73.2 (3)  | C51—C5—C6—C7  | −72.8 (4) |
| C9—C1—C2—C6  | 69.6 (3)   | C5—C6—C7—C8   | 177.9 (2) |
| C9—C1—C2—O3  | −177.4 (3) | C2—C6—C7—C8   | −73.4 (3) |
| C10—C1—C2—C6 | −36.9 (3)  | C5—C6—C7—C10  | −75.4 (3) |
| C10—C1—C2—O3 | 76.0 (3)   | C2—C6—C7—C10  | 33.3 (3)  |

|               |            |                 |            |
|---------------|------------|-----------------|------------|
| C1—C2—C6—C7   | 2.3 (3)    | C6—C7—C10—C1    | −54.5 (2)  |
| C1—C2—C6—C5   | 122.0 (2)  | C8—C7—C10—C1    | 56.0 (3)   |
| C1—C2—O3—N4   | −117.3 (3) | C6—C7—C8—C9     | 69.6 (3)   |
| O3—C2—C6—C7   | −115.5 (2) | C10—C7—C8—C9    | −37.0 (3)  |
| O3—C2—C6—C5   | 4.2 (3)    | C7—C8—C9—C1     | 2.2 (3)    |
| C6—C2—O3—N4   | −4.6 (3)   | C5—C51—C56—C55  | −179.8 (3) |
| C2—O3—N4—C5   | 3.1 (3)    | C5—C51—C52—C53  | 179.9 (3)  |
| O3—N4—C5—C6   | −0.1 (3)   | C52—C51—C56—C55 | −0.2 (5)   |
| O3—N4—C5—C51  | 179.3 (2)  | C56—C51—C52—C53 | 0.3 (5)    |
| N4—C5—C6—C2   | −2.7 (3)   | C51—C52—C53—C54 | 0.7 (6)    |
| N4—C5—C51—C52 | −4.8 (4)   | C52—C53—C54—C55 | −1.7 (6)   |
| N4—C5—C51—C56 | 174.8 (3)  | C53—C54—C55—C56 | 1.8 (6)    |
| N4—C5—C6—C7   | 106.6 (3)  | C54—C55—C56—C51 | −0.8 (5)   |

Structure solution was by direct methods using *DIRDIF* (Beurskens *et al.*, 1990). Non-H atoms were located with *DIRDIF* using the *ORIENT* option with the norbornane rigid group, taken from *DIRDIF ORBASE*, as an input model. Refinement was performed by full-matrix least squares using *SHELX76* (Sheldrick, 1976). All the non-H atoms were refined with anisotropic displacement parameters. H atoms were located from a difference Fourier map and fixed at a distance of 0.95 Å from their parent atom. H atoms were included in the structure-factor calculations and given displacement parameters equal to 1.1U<sub>eq</sub> of their parent atom, but their parameters were not refined. *PARST* (Nardelli, 1983) was used for molecular geometry calculations and molecular graphics were prepared using *PLUTO* (Motherwell & Clegg, 1978).

AT thanks the University Grants Commission, India, for the award of a teaching fellowship. AN wishes to thank UGC for the award of a senior research fellowship.

---

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates, angles involving H atoms and least-squares-planes data have been deposited with the IUCr (Reference: AL1074). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

---

## References

- Beurskens, P. T., Bosman, W. P., Doesburg, H. M., Gould, R. O., van den Hark, Th. E. M., Prick, P. A. J., Noordik, J. H., Beurskens, G., Parthasarathi, V., Bruins Slot, H. J., Haltiwanger, R. C., Strumpel, M. K., Smits, J. M. M., Garcia-Granda, S., Smykalla, C., Behm, H. J. J., Schäfer, G. & Admiraal, G. (1990). *The DIRDIF Program System*. Crystallography Laboratory, Univ. of Nijmegen, The Netherlands.
- Caramella, P. & Grunanger, P. (1984). In *1,3-Dipolar Cycloaddition Chemistry*, Vol. I, edited by A. Padwa. New York: Wiley.
- Curran, D. P. (1988). In *Advances in Cycloaddition*. New York: JAI.
- Dapporto, P., Paoli, P., Brandi, A., De Sarlo, F., Goti, A. & Guarna, A. (1992). *Acta Cryst.* **B48**, 234–238.
- Estienne, J., Pepe, G., Siri, D., Baldy, A. & Samat, A. (1992). *Acta Cryst.* **C48**, 352–354.
- Grundmann, C. & Grunanger, P. (1971). In *The Nitrile Oxides, Versatile Tools of Theoretical and Preparative Chemistry*. Berlin: Springer-Verlag.
- Kemp, W. (1986). In *NMR in Chemistry*. New York: Macmillan.
- Lang, S. A. & Lin, Y. (1984). In *Comprehensive Heterocyclic Chemistry*, Vol. 6, edited by A. Katritzky & C. W. Rees. London: Pergamon.
- Larsen, K. E. & Torrsell, K. B. G. (1984). *Tetrahedron*, **40**, 2985.

- Motherwell, W. D. S. & Clegg, W. (1978). *PLUTO. Program for Plotting Molecular and Crystal Structures*. Univ. of Cambridge, England.
- Nagarajan, A. & Krishna Pillay, M. (1993a). *Indian J. Chem.* **32B**, 471–474.
- Nagarajan, A. & Krishna Pillay, M. (1993b). *Indian J. Chem.* **32B**, 938–941.
- Nardelli, M. (1983). *Comput. Chem.* **7**, 95–98.
- Sheldrick, G. M. (1976). *SHELX76. Program for Crystal Structure Determination*. Univ. of Cambridge, England.
- Torrsell, K. B. G. (1988). In *Nitrile Oxides, Nitrones and Nitronates in Organic Synthesis*. New York: Verlag Chemie.

*Acta Cryst.* (1994). **C50**, 1814–1818

## Influence of Crystal Environment on Molecular Conformation: *p*-Bromo-*N*-(*p*-dimethylaminobenzylidene)aniline

MUSTAFA T. AHMET, JACK SILVER AND  
ANDREW HOULTON

*Department of Chemistry and Biological Chemistry,  
University of Essex, Colchester CO4 3SQ, England*

(Received 29 September 1993; accepted 25 March 1994)

## Abstract

*p*-Bromo-*N*-(*p*-dimethylaminobenzylidene)aniline, C<sub>15</sub>H<sub>15</sub>BrN<sub>2</sub>, (1), comprises two independent molecules with markedly different conformations. Molecule *A* is essentially planar along its entire length (including the dimethylamino group) whereas molecule *B* deviates considerably from planarity with an interplanar angle of 145.8 (3)° between the two phenyl rings and the NMe<sub>2</sub> group twisted by 10 (2)° out of the plane of the phenyl ring to which it is bound. While polymorphism and crystal packing effects are well documented for benzylidene derivatives, this is an interesting example with two quite distinct conformations in the same crystal phase. The differences are accounted for in terms of crystal-packing effects. The structure highlights the ability of particular molecular arrangements to stabilize a less favourable molecular conformation.

## Comment

Recently, we embarked on a programme to develop molecular materials which exhibit non-linear optical properties (Houlton, Jassim, Roberts, Silver, McArdle, Cunningham & Higgins, 1992; Houlton, Miller, Silver, Jassim, Ahmet, Axon, Bloor & Cross, 1993). Two classes of compound have been considered: organometallic and organic derivatives.