This article was downloaded by: [Statsbiblioteket Tidsskriftafdeling] On: 25 April 2014, At: 02:23 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lncn20

2′-0-

Hydroxyalkoxymethylribonucleosides and their Incorporation into Oligoribonucleotides

Sergey N. Mikhailov^a, Georgii V. Bobkov^a, Kyrill V. Brilliantov^a, Jef Rozenski^b, Arthur Van Aerschot^b, Piet Herdewijn^b, Michael H. Fisher^c & Rudolph L. Juliano^c

^a Engelhardt Institute of Molecular Biology, Russian Academy of Sciences , Moscow, Russia

^b Laboratory of Medicinal Chemistry, Rega Institute for Medical Research, Katholieke Universiteit Leuven, Leuven, Belgium

^c Department of Pharmacology, School of Medicine, University of North Carolina, Chapel Hill, North Carolina, USA Published online: 05 Dec 2007.

To cite this article: Sergey N. Mikhailov, Georgii V. Bobkov, Kyrill V. Brilliantov, Jef Rozenski, Arthur Van Aerschot, Piet Herdewijn, Michael H. Fisher & Rudolph L. Juliano (2007) 2'-O-Hydroxyalkoxymethylribonucleosides and their Incorporation into Oligoribonucleotides, Nucleosides, Nucleotides and Nucleic Acids, 26:10-12, 1509-1512, DOI: <u>10.1080/15257770701543007</u>

To link to this article: http://dx.doi.org/10.1080/15257770701543007

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



2'-O-HYDROXYALKOXYMETHYLRIBONUCLEOSIDES AND THEIR INCORPORATION INTO OLIGORIBONUCLEOTIDES

Sergey N. Mikhailov, Georgii V. Bobkov, and Kyrill V. Brilliantov *Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Moscow, Russia*

Jef Rozenski, Arthur Van Aerschot, and Piet Herdewijn *Laboratory of Medicinal Chemistry, Rega Institute for Medical Research, Katholieke Universiteit Leuven, Leuven, Belgium*

Michael H. Fisher and Rudolph L. Juliano Department of Pharmacology, School of Medicine, University of North Carolina, Chapel Hill, North Carolina, USA

 \Box A simple and efficient method for the preparation of pyrimidine 2'-Ohydroxyethoxymethylribonucleosides and 2'-O-hydroxypropoxymethylribonucleosides has been developed. These modified nucleosides were incorporated into oligoribonucleotides, which were shown to form stable RNA/RNA duplexes. The effect of 2'-O-modification in the antisense and sense strands of small interference RNA was evaluated in multi-drug resistant NIH 3T3 cells.

Keywords Oligoribonucleotides; ONs; RNA/RNA duplexes

Modified oligonucleotides (ONs) are widely used in molecular biology and medicinal chemistry. Antisense agents are valuable tools to inhibit the expression of a target gene in a sequence specific manner, and may be used for functional genomics, target validation and therapeutic purposes.^[1] Research in this field increased in impact by the discovery of RNA interference.^[2,3]

Numerous attempts have been made to improve the properties of natural ONs. It is believed that the 2'-carbohydrate modifications are the most universal and promising.^[4–6] A typical strategy for the preparation of such ONs is the synthesis of modified nucleoside followed by conversion to the corresponding phosphoramidite suitable for automated ON

Financial support of the Russian Foundation for Basic Research, Programme "Molecular and Cellar Biology," INTAS and KUL Research Council is acknowledged.

Address correspondence to Piet Herdewijn, Laboratory of Medicinal Chemistry, Rega Institute for Medical Research, Katholieke Universiteit Leuven, Minderbroedersstraat 10, Leuven B 3000, Belgium. E-mail: piet herdewijn@rega.kuleuven.ac.be



synthesis. Most of these modifications were achieved via alkylation reactions of partially protected ribonucleosides. The heterocyclic bases should be protected in order to avoid their alkylation. For each nucleoside the specific blocking groups are used,^[7] which in most cases are not compatible with standard automated ON synthesis. Very recently for the preparation of 2'-O-modified nucleosides we have developed another type of chemistry based on O-glycosylation reactions, which have several important advantages, such as increased yields and simplification.^[8,9] A simple and effective method for the preparation of 2'-O-(β -D-ribofuranosyl)nucleosides starting from readily available 3',5'-O-blocked N-acylribonucleosides and 1-O-acetyl-2,3,5-tri-O-benzoyl- β -D-ribofuranose preactivated with tin tetrachloride in 1,2-dichloroethane at 0°C, has been recently developed.^[10-12]

RESULTS AND DISCUSSION

Here we present our recent results on the preparation of 2'-Ohydroxyalkoxymethylribonucleosides and their incorporation into oligoribonucleotides.^[13] This scheme is nearly the same as for the preparation of disaccharide nucleosides and their incorporation into ONs.^[14] The starting compounds 1(n = 1, 2)^[15] were easily prepared from 1,3-dioxolane and 1,3-dioxalane, respectively, and were condensed with 3',5'-O-blocked ribonucleosides **2** (B = Ura, Cyt^{Bz}) in the presence of tin tetrachloride in 1,2-dichloroethane at -12° C for 20 minutes (Figure 1). The yields (85–87%) of products **3** were even higher than in the case of preparation of above mentioned disaccharide nucleosides.^[10] The silyl group was cleaved to yield partially protected **4** and subsequent deblocking with ammonia in methanol gave free 2'-O-substituted nucleosides **5** in high overall yield. Nucleosides **4** were converted using standard procedures to the corresponding monomethoxytrityl derivatives **6** and their phosphoramidites **7**. With these phosphoramidites, several oligoribonucleotides with one or more modifications were assembled. It was shown that modified ONs form stable RNA/RNA duplexes, with about 0.5°C destabilization per modification.

In addition several siRNAs comprising one or more of the new modifications have been prepared and are currently under evaluation. The effect of 2'-O-modification in the antisense and sense strands of small interfering RNA targeting the MDR1 gene was evaluated in multi-drug resistant NIH 3T3 cells using previously described techniques.^[16] Standard siRNA or siRNAs with 2'-O-hydroxyalkoxymethylribonucleosides in the sense strand effectively inhibited P-glycoprotein expression, whereas modifications in the antisense strand were not tolerated.

CONCLUSIONS

The general method for the preparation of $2'-O-\beta$ -D-ribofuranosylnucleosides was found to be applicable for the synthesis of pyrimidine 2'-O-hydroxyalkoxymethylribonucleosides. The 2'-O-substitutent was found to be stable during oligonucleotide synthesis. Additional work on the preparation of other 2'-O-functionalized nucleosides and oligonucleotides bearing amino groups is in progress and will be published shortly.

REFERENCES

- Kurrech, J. Antisence technologies. Improvements through novel chemical modifications. *Eur. J. Biochem.* 2003, 270, 1628–1644.
- Sioud, M.; Iversen, P.O. Ribozymes, DNAzymes and small interfering RNAs as therapeutics. *Curr. Drug Targets* 2005, 6, 647–653.
- Coppelli, F.M.; Grandis, J.R. Oligonucleotides as anticancer agents: From the benchside to the clinical and beyond. *Curr. Pharm. Des.* 2005, 11, 2825–2840.
- Manoharan, M. 2'-Carbohadrate modification in antisense oligonucleotide therapy: Importance of conformation, configuration and conjugation. *Biochim. Biophys. Acta* 1999, 1489, 117–130.
- Freier, S.M.; Altmann, K.-H. The ups and downs of nucleic acid duplex stability: Structure-stability studies on chemically-modified DNA:RNA duplexes. *Nucleic Acids Res.* 1997, 25, 4429–4443.
- Jin, S.; Miduturu, C.V.; McKinney, D.C.; Silverman S.K. Synthesis of amine- and thiole-modified nucleoside phosphoramidites for site-specific introduction of biophysical probes into RNA. *J. Org. Chem.* 2005, 70, 4284–4299.

- Zatsepin, T.S.; Romanova, E.A.; Oretskaya, T.S. Synthesis of 2'-O-alkylnucleosides. *Russ. Chem. Rev.* 2002, 71, 513–534.
- 8. Efimtseva, E.V.; Mikhailov, S.N. Disaccharide nucleosides. Russ. Chem. Rev. 2004, 73, 401-414.
- Nauwelaerts, K.; Efimtseva, E.V.; Mikhailov, S.N.; Herdewijn, P. Disaccharide nucleosides, an important group of natural compounds. In *Frontiers in Nucleosides and Nucleic Acids*, Schinazi, R.F. Liotta, D.C. eds. IHL Press, Tucker, GA, pp. 187–220 2004.
- Mikhailov, S.N.; Efimtseva, E.V.; Gurskaya, G.V.; Zavodnik, V.E.; De Bruyn, A.; Janssen, G.; Rozenski, J.; Herdewijn, P. An efficient synthesis and physico-chemical properties of 2'-O-β-D-ribofuranosylnucleosides, minor tRNA components. *J. Carbohydr. Chem.* **1997**, 16, 75–92.
- Mikhailov, S.N.; Rodionov, A.A.; Efimtseva, E.V.; Ermolinsky, B.S.; Fomitcheva, M.V.; Padyukova N.Sh.; Rothenbacher, K.; Lescrinier, E.; Herdewijn, P. Formation of Trisaccharide Nucleoside during disaccharide nucleoside synthesis. *Eur. J. Org. Chem.* **1998**, 2193–2199.
- Mikhailov, S.N.; Efimtseva E.V. Disaccharide nucleosides and oligonucleotides on their basis. *Collect. Czech. Chem. Commun. Symp. Ser.* 2002, 5, 181–194.
- Bobkov, G.V.; Brilliantov, K.V.; Mikhailov, S.N.; Rozenski, J.; Van Aerschot, A.; Herdewijn, P. Synthesis of oligoribonucleotides containing pyrimidine 2'-O-[(hydroxyalkoxy)methyl] ribonucleosides. *Collection Czech. Chem. Comm.* 2006, 71, 804–819.
- Efimtseva, E.V.; Bobkov, G.V.; Mikhailov, S.N.; Van Aerschot, A.; Schepers, G.; Busson, R.; Rozenski, J.; Herdewijn, P. Oligonucleotides containing disaccharide nucleosides. *Helv. Chim. Acta* 2001, 84, 2387–2397.
- 15. Senkus, M. Reaction of some cyclic acetals with acetic anhydride. J. Am. Chem. Soc. 1946, 68, 734-736.
- Xu, D.; Kang, H.; Fisher, M.; Juliano, R.L. strategies for the inhibition of MDR1 gene expression. *Mol. Pharmacol.* 2004, 66, 268–275.