Nucleosides. XLV. $1-\alpha-L-aldo$ -Pentofuranosylpyrimidines¹

NAOTAKA YAMAOKA, BRIAN A. OTTER, AND JACK J. FOX

Division of Biological Chemistry, Sloan-Kettering Institute for Cancer Research, Sloan-Kettering Division of Cornell University Medical College, New York 21, New York

Received July 6, 1967

Methods are described for the synthesis of the 1- α -L-arabinofuranosyl derivatives of thymine, cytosine, and uracil. The uracil and thymine derivatives were converted *via* anhydronucleoside intermediates to nucleosides of the α -L-xylo and α -L-lyxo configurations. Two syntheses of 1- α -L-xylofuranosylcytosine (the C-4' epimer of the biologically active 1- β -D-arabinofuranosylcytosine) from 2,2'-anhydro-1-(3',5'-di-O-benzoyl- α -L-xylofuranosyl)uracil are described. Preliminary screening studies of these L nucleosides showed no significant activity against L1210 mouse leukemia or Burkitt's tumor cells in tissue culture. The cytosine nucleosides are not de-aminated by human liver or mouse kidney homogenates, nor do they inhibit the deamination of 1- β -D-arabino-furanosylcytosine in these systems.

The sugar moieties of all nucleosides hitherto found in nature (*i.e.*, those derived from the nucleic acids or from nucleoside antibiotics) are of the D configuration.^{2,3} This report deals with the synthesis of pyrimidine nucleoside analogs which differ from certain biologically important nucleosides *only* in the configuration of C-4'. Such C-4' epimers are the hitherto unknown 1- α -L-aldo-pentofuranosylpyrimidines. Of special interest are the synthesis and biological study of the C-4' epimer of the chemotherapeutically active 1- β -D-arabinofuranosylcytosine⁴ and of the tRNA component, 1- β -D-ribofuranosylthymine.⁵

Direct access to α -L-thymine nucleosides (without the production of anomeric mixtures) was readily achieved by utilization of the mercuri procedure⁶ (see Scheme I). Condensation of an anomeric mixture of tri-O-benzoyl-L-arabinofuranosyl bromides⁷ (1) with dithyminyl mercury⁶ in refluxing toluene afforded 50-70% yield of 1-(tri-O-benzoyl- α -L-arabinofuranosyl)thymine (2) which was debenzoylated with NaOH in 50% ethanol to $1-\alpha$ -L-arabinofuranosylthymine (3a). The L-ara-thymine nucleoside (3a) showed, as expected, ultraviolet absorption properties generally similar to those for 1-glycosylthymines⁸ and, when treated with metaperiodate, **3a** consumed 1 mole/mole, slowly (40 hr), in accord with a trans-glycol system. The fact that the optical rotation of **3a** was of equal magnitude to but of opposite sign from the known enantiomorphic nucleoside, 1-α-D-arabinofuranosylthymine,⁹ establishes the α -L configuration for **3a** and for all the nucleosides subsequently derived from it $(3a \rightarrow 9a, 10a)$. Mesylation of **3a** with methanesulfonyl chloride in pyridine afforded $1-(tri-O-mesyl-\alpha-L-arabinofuranosyl)$ thymine (4a). Large-scale preparation of 4a may be achieved from 1 in $\sim 35\%$ over-all yield without isolation of 2 and 3a.

- (4) S. S. Cohen, *ibid.*, **5**, 2 (1966).
- (4) B. B. Conen, *ibid.*, **9**, 2 (1966).
 (5) P. R. Srinivasan and E. Borek, *ibid.*, **5**, 157 (1966).
- (6) J. J. Fox, N. Yung, J. Davoll, and G. B. Brown, J. Am. Chem. Soc., 78, 2117 (1956).
- (7) Tri-O-benzoyl- α,β -L-arabinofuranosyl bromide was prepared according to the method described for the enantiomorph; see H. G. Fletcher, Jr., Methods Carbohydrate Chem., **2**, 228 (1963); R. K. Ness and H. G. Fletcher, Jr., J. Am. Chem. Soc., **80**, 2007 (1958).

(8) J. J. Fox and D. Shugar, Biochim. Biophys. Acta, 9, 369 (1952).

(9) T. Nishimura and B. Shimizu, Chem. Pharm. Bull. (Tokyo), 7, 803 (1965).

For the conversion of **4a** to the α -L-xylo and α -L-lyxo isomers (**9a** and **10a**), approaches previously employed in this laboratory^{10,11} were adapted. Treatment of **4a** with 1 equiv of NaOH in 50% ethanol yielded the 2,2'-anhydro derivative **5a** which, after treatment with sodium benzoate in DMF, gave the 2,2'-anhydro 5'-benzoate (**6a**, X = O). Reaction of **6a** with sodium benzoate in DMF and benzoic acid gave, as major products, **7a** and **8a**.

A unified pathway for the conversion of anhydro nucleosides of type $\mathbf{6}$ ((X = O) to nucleosides of type $\mathbf{7}$ and $\mathbf{8}$ has been discussed.¹¹ Formation of anhydro nucleoside 7 probably proceeds via benzoate attack at C-2' of the protonated nucleoside 6, followed by the intramolecular displacement of the 3'-mesyloxy function with formation of a 2',3'-orthoester ion. Attack by the 2-carbonyl on this orthoester would produce 7a. Attack of benzoate ion on 7a produced the α -L-lyxo nucleoside 8a. Alkali treatment of 7a and 8a in dilute alcohol gave the unblocked crystalline nucleosides 9a and 10a, respectively. Nucleosides 9a and 10a differed in melting point and electrophoretic mobility from 3a but showed ultraviolet absorption properties expected for 1-substituted glycosylthymines.8 When 9a was treated with metaperiodate, 1 equiv of oxidant was consumed. This consumption was very slow (170 hr) indicative of a *trans*-glycol system. These data es-tablish the L-xylo configuration of **9a**. Nucleoside **10a** showed a rapid uptake of metaperiodate (within 3 min), similar to that reported for $1-\alpha$ -D-ribofuranosylthymine,⁶ consonant with a *cis*-glycol system. Nucleoside 10a exhibited the same melting point as the enantiomorphic nucleoside $1-\alpha$ -D-lyxofuranosylthymine⁹ and gave identical nmr data. These data establish 10a as 1- α -L-lyxofuranosylthymine. It should be noted that the L-lyxo (10a) and L-xylo (9a) nucleosides are the C-4' epimers of $1-\beta$ -D-ribofuranosylthymine⁶ and of "spongothymidine" (1- β -D-arabinofuranosylthymine).¹²

Easy access to the uracil nucleoside (3b) was accomplished by the deamination of 11 which was prepared by the procedure of Yamaoka, *et al.*^{13,14} Condensa-

⁽¹⁾ This investigation was supported in part by funds from the National Cancer Institute, National Institutes of Health, U. S. Public Health Service (Grant No. CA 08748).

⁽²⁾ J. J. Fox and I. Wempen, Advan. Carbohydrate Chem., 14, 283 (1959).
(3) J. J. Fox, K. A. Watanabe, and Λ. Bloch, Progr. Nucleic Acid Res. Mol. Biol., 5, 252 (1966).

⁽¹⁰⁾ J. F. Codington, R. Fecher, and J. J. Fox, J. Am. Chem. Soc., 82, 2794 (1960).

⁽¹¹⁾ J. J. Fox and N. C. Miller, J. Org. Chem., 28, 936 (1963).

⁽¹²⁾ J. J. Fox, N. Yung, and A. Bendich, J. Am. Chem. Soc., 79, 2775 (1957).

⁽¹³⁾ N. Yamaoka, K. Aso, and K. Matsuda, J. Org. Chem., 30, 149 (1965).
(14) The direct synthesis of uracil nucleosides by this procedure is currently under investigation in this laboratory.



tion of N⁴-benzoylcytosine¹⁵ with tri-O-benzoyl- α,β -Larabinofuranosyl bromide (1) in nitromethane containing mercuric cyanide afforded a high yield (80-90%) of the tetrabenzoylated nucleoside (11). Treatment of 11 with alkali gave $1-\alpha$ -L-arabinofuranosylcytosine (12) in quantitative yield. Conventional deamination of 12 with nitrous acid did yield 3; however, the method was cumbersome and the yields were low. Selective N⁴-benzoylation¹⁶ of 12 gave 13 in good yield which was smoothly converted to **3b** by refluxing with hot aqueous acetic acid.¹⁷ It was found more convenient to prepare **3b** by treatment of **11** with hot aqueous acetic acid and then with 1 N NaOH. The arabino nucleoside (**3b**) formed an amorphous tri-O-mesylate (4b) which was converted with alkali to the crystalline anhydro-L-ribo nucleoside (5b). Reaction of 5b with sodium benzoate in DMF at 100° gave 6b (X = O). Treatment of 6b (X = O) with

sodium benzoate in DMF with benzoic acid (as described above for the thymine nucleoside series) yielded a mixture of **7b** and **8b**, the latter of which was converted to $1-\alpha-L-lyxofuranosyluracil (10b)$ by alkali. Nucleoside **10b** showed physical properties expected for the enantiomorph of the known $1-\alpha-D-lyxofuranosyluracil.⁹$

Nucleoside 7b was converted to the crystalline 1- α -L-xylofuranosylcytosine (16) in 27% over-all yield in four steps (via 9b, 14, and 15) without isolation of the intermediates. When treated with metaperiodate, nucleoside 16 consumed 1 equiv slowly (as did 12) indicative of a *trans*-glycol system. Compound 16 differed from 12 in melting point, optical rotation, and paper-electrophoretic migration. All of the data permit the assignment of only the L-xylo configuration to 16.

Earlier studies¹⁸ had shown that certain anhydro nucleosides could be thiated to their 4-thiono derivatives, and when **6b** (X = O) was thiated with P₂S₅ in pyridine under mild anhydrous conditions, **6b** (X = S) was obtained. Similarly, thiation of the di-O-benzoate (**7b**) gave a 64% yield of **17**. Treatment

(18) N. Yung and J. J. Fox, J. Org. Chem., 27, 1447 (1962).

⁽¹⁵⁾ D. M. Brown, A. R. Todd, and S. Varadarajan, J. Chem. Soc., 2384 (1956).

⁽¹⁶⁾ K. A. Watanabe and J. J. Fox, Angew Chem. Intern. Ed. Engl., 5, 579 (1966); B. A. Otter and J. J. Fox in "Synthetic Procedures in Nucleic Acid Chemistry," Vol. 1, W. W. Zorbach and R. S. Tipson, Ed., Interscience Publishers, Inc., New York, N. Y., in press.

⁽¹⁷⁾ Brown, et al., 15 had demonstrated that N⁴-benzoylcytosine is converted to uracil and cytosine (ratio 33:1) by this procedure.

of 17 with dilute alkali afforded 18 which was converted to 1- α -L-lyxofuranosylcytosine (16) in $\sim 45\%$ over-all yield from 7b. Nucleosides 10b and 16 are the C-4' epimers of uridine and $1-\beta$ -D-arabinofuranosylcytosine, respectively.

Screening Studies.^{19,20}—In preliminary studies the 1- α -L derivatives of arabinosyl- and xylosylcytosine (12 and 16) were inactive against L1210 mouse leukemia at doses of 120 mg/kg given daily intraperitoneally, starting 24 hr after inoculation of leukemia and continued for 10 consecutive days. None of the $1-\alpha-L$ nucleosides (3a, 9a, 10a, 12, and 16) were inhibitory to Burkitt's tumor cells in culture in doses as high as 30 mg/ml. The cytosine L-nucleosides 12 and 16 (unlike $1-\beta$ -D-arabinofuranosylcytosine) were not deaminated to their uracil analogs by human liver or mouse kidney homogenates, nor did they inhibit the deamination of $1-\beta$ -D-arabinofuranosylcytosine in these systems even with a substrate to inhibitor ratio of 10:1. These data attest to the importance of the configuration at C-4' in the biological activity of 1- β -D-arabinofuranosylcvtosine.

Experimental Section²¹

1-(Tri-O-benzoyl- α -L-arabinofuranosyl)thymine (2).—Powdered dithyminylmercury⁶ (2.15 g, 4.76 mmoles) was added to 150 ml of dry toluene. The vigorously stirred suspension was dried by azeotropic distillation of about 50 ml of solvent. A solution of tri-O-benzoyl- α,β -L-arabinofuranosyl bromide⁷ (5.0 g, 9.52 mmoles) in 30 ml of dry toluene was added and a further 30 ml of solvent removed by distillation. Refluxing was continued for 1 hr during which time most of the solid material dissolved. The warm solution was filtered from 0.5 g of insoluble material, and the filtrate was concentrated in vacuo to ca. 50 ml. The solution was treated with 500 ml of petroleum ether (bp $30-60^{\circ}$) and the resulting precipitate was collected and dissolved in CHCl₃. After filtration, the CHCl₃ solution was washed with freshly prepared 30% aqueous KI then H₂O and dried (Mg-SO₄). Removal of solvent in vacuo left a yellow, viscous syrup which contained 2 and several minor components. A solution of the syrup in C_6H_6 -Et₂O (2:1) was placed on a column of silica gel (2.5 \times 30 cm). The column was eluted with C₆H₆-Et₂O (2:1) until the minor components had been removed, then with EtOAc. The syrup, obtained by concentration of the EtOAc solution, was chromatographically pure but resisted all attempts at crystallization. When stored *in vacuo*, it changed to a colorless, amorphous solid (3.9 g, 72%): $\lambda_{\text{max}}^{\text{dil EtoH}}$ 235, 268 mµ; λ_{min} 210, 257 mµ.

Anal. Calcd for C₃₁H₂₆N₂O₉: C, 65.26; H, 4.95; N, 4.91. Found: C, 65.29; H, 4.75; N, 4.99.

1- α -L-Arabinofuranosylthymine (3a).—Amorphous 2 (1.0 g 1.75 mmoles) was dissolved in 15 ml of warm, 50% aqueous EtOH containing 6.0 ml of 1 N NaOH. The solution was refluxed for 30 min, cooled, and neutralized with HOAc. The concentrated solution (ca. 7 ml) was treated with Dowex 50 (H⁺) (ca. 1 g of moist resin). The precipitated benzoic acid was removed along with the resin, and the resin treatment was repeated. The last traces of benzoic acid were removed by extraction of the filtrate with CHCl₃. Concentration of the aqueous phase afforded 423 mg (90%) of an amorphous solid. The product, in 90% aqueous EtOH, crystallized after several months at 4°. Even with seed crystals, cystallization of 3a (from several solvent systems) was difficult. Crystalline material melted indistinctly at 115-125°

57

(eff) and showed $[\alpha]^{25}D - 37^{\circ}$ (c 0.7, H₂O). The enantiomorph, solid with $[\alpha]_{D} + 40^{\circ}$ (H₂O), $\lambda_{max}^{H_2O}$ 267 m μ , λ_{min} 235 m μ . Anal. Calcd for C₁₀H₁₄N₂O₆: C, 46.51; H, 5.47; N, 10.85.

Found: C, 45.90; H, 5.50; N, 10.45.

1-(Tri-O-mesyl- α -L-arabinofuranosyl)thymine (4a). Method A.-MeSO₂Cl (0.06 ml, 0.77 mmole) was added to a solution of 56.0 mg (0.22 mmole) of amorphous 3a in 2 ml of dry pyridine. After 24 hr at room temperature, the dark solution was poured into 50 ml of ice-H₂O. Crystalline material slowly formed, and after ca. 30 min, the solid was collected and washed with H_2O . The solid was dissolved in hot 50% aqueous EtOH and the solution was treated with charcoal. The product (50 mg, 49%) crystallized from the cold filtrate as fine needles. Analytically pure material, mp 125-126°, was obtained after a further re-crystallization; $\lambda_{\max}^{dil E:OH}$ 266 m μ , λ_{\min} 233 m μ .

Anal. Calcd for $C_{13}H_{20}N_2O_{12}S_3$: C, 31.70; H, 4.09; N, 5.69; S, 19.53. Found: C, 31.59; H, 4.29; N, 5.69; S, 19.29.

Method B.—Crude, syrupy 2 (obtained as above from 5.60 g of dithyminylmercury and 13.0 g of 1) was dissolved in 150 ml of 50% aqueous EtOH containing 60 ml of 1 N NaOH; this was refluxed for 30 min and processed as described above for the preparation of 3a. The resulting syrupy residue was dissolved in EtOH and dried by azeotropic distillation of the solvent. The syrup was dissolved in pyridine (100 ml) and the distillation was repeated. The dry syrup was dissolved in 100 ml of dry pyridine and the solution was cooled to 5° . MeSO₂Cl (8.0 ml) was added dropwise, and the solution was kept at room temperature for 2 hr. The dark reaction mixture was poured into 700 ml of cold H_2O . After 2 hr at 4°, the crystalline material was colletted and recrystallized from 50% aqueous EtOH (char-The yield was 4.0 g (33%) of material (mp 123-125°) coal). suitable for transformation to 5a.

2,2'-Anhydro-1-(3',5'-di-O-mesyl- α -L-ribofuranosyl)thymine (5a).—A hot solution of 1.0 g (2.03 mmoles) of 4a in 100 ml of 50% aqueous EtOH was cooled quickly to avoid crystallization. When the temperature reached 40° , 2.08 ml of NaOH (1.0 N) was added slowly with stirring. Within 5 min of the addition, crystalline material formed. The mixture was kept at 5° for 1 hr then filtered. Recrystallization of the product (700 mg, 87%) from dilute EtOH afforded colorless needles: mp 246-247°; $\lambda_{\text{max}}^{\text{dil EtOH}}$ 227, 248 m μ ; λ_{min} 214, 238 m μ .

Anal. Caled for $C_{12}H_{16}N_2O_9S_2$: C, 36.38; H, 4.07; N, 7.07; S, 16.18. Found: C, 36.46; H, 3.96; N, 7.01; S, 15.90.

2,2'-Anhydro-1-(5'-O-benzoyl-3'-O-mesyl-α-L-ribofuranosyl)thymine (6a, X = O).—Compound 5a (2.0 g, 5.04 mmoles) was added to a hot solution of 805 mg (5.6 mmoles) of sodium benzoate in 200 ml of dry DMF. A clear solution was obtained after ca. 30 min of heating on a steam bath. After 4 hr of heating, the solution was concentrated in vacuo (oil pump, bath 55°) to ca. 50 ml. The solution was poured into 400 ml of cold H_2O and, after 1 hr of cooling, the solid was collected and washed with H₂O. Recrystallization from EtOAc afforded 1.6 g (75%) of pure material: mp 234-235°; $\lambda_{max}^{dil EtOH}$ 232, shoulder 250-270 m μ ; λ_{min} 212 mu.

Anal. Calcd for $C_{18}H_{18}N_2O_8S$: C, 51.18; H, 4.29; N, 6.63; S, 7.58. Found: C, 51.09; H, 4.33; N, 6.47; S, 7.70.

2,2'-Anhydro-1- $(3',5'-di-O-benzoyl-\alpha-L-xylofuranosyl)$ thymine (7a) and Tri-O-benzoyl- α -L-lyxofuranosylthymine (8a).—A solution of 2.0 g (4.73 mmoles) of **6a** (X = O) in 400 ml of dry DMF containing 2.72 g (18.82 mmoles) of sodium benzoate and 580 mg (4.73 mmoles) of benzoic acid was heated at ca. 150° for 4.5 hr. The dark reaction mixture was cooled and concentrated to dryness *in vacuo* (oil pump) at 50°. The solid residue was partitioned between 100 ml of H₂O and 100 ml of CHCl₃, and the organic layer was washed with H₂O and dried (MgSO₄). Removal of the solvent afforded a thin syrup from which the last traces of DMF were removed by codistillation with xylene. The residue was dissolved in 70 ml of hot EtOH and the solution cooled slowly to room temperature. Crystalline 7a (1 g) was collected and washed with cold EtOH. Recrystallization from EtOH afforded 920 mg (43%) of pure material: mp 251-253°; $\lambda_{max}^{dil EtOH}$ 233, shoulder 250–270 m μ ; λ_{\min} 212 m μ .

Anal. Caled for C24H20N2O7: C, 64.27; H, 4.50; N, 6.25. Found: C, 63.87; H, 4.41; N, 5.90.

⁽¹⁹⁾ The authors are indebted to Drs. J. H. Burchenal, M. R. Dollinger and R. W. Mackey of this institute for these preliminary data

⁽²⁰⁾ See M. R. Dollinger, J. H. Burchenal, W. Kreis, and J. J. Fox. Biochem. Pharmacol., 16, 689 (1967), and references therein for a description of the experimental systems used in these screening studies.

⁽²¹⁾ Microanalyses were performed by the Spang Microanalytical Laboratories, Ann Arbor, Mich., and by Galbraith Laboratories, Inc., Knoxville, Tenn. Melting points, determined on a Thomas-Hoover capillary melting point apparatus, are corrected.

The mother liquors and washings obtained from the preparation of 7a were concentrated to dryness, and the residue was dissolved in the minimum of EtOAc-MeOH (20:1). The solution was placed on a column of silica gel $(2.5 \times 30 \text{ cm})$ and the column was eluted with the same solvent. Concentration of the 200-

350-ml portion of the eluate afforded 160 mg (7.5%) of pure 7a mp and mmp 251-253°. Concentration of the 65-140-ml portion and recrystallization of the residue from EtOH afforded 270 mg $(10^{e_{\ell}}_{e_{\ell}})$ of **8a**: mp 183–184°; $\lambda_{\max}^{dil EOH}$ 228, 265 m μ ; λ_{\min} 215, 255 mµ.

Anal. Caled for $C_{31}H_{26}N_2O_9$: C, 65.26; H, 4.59; N, 4.91. Found: C, 65.35; H, 4.65; N, 4.69.

Treatment of 400 mg (0.9 mmole) of 7a with sodium benzoate (520 mg, 3.6 mmoles) and benzoic acid (100 mg, 0.9 mmole) in 40 ml of DMF at 150° for 24 hr and work-up of the solution as described above afforded a syrup which contained starting material (7a) and 8a. Fractional crystallization of the syrup from EtOH afforded 120 mg of 7a. Column chromatography on silica gel, as described above, afforded 150 mg of 8a, mp 183-184°, and a further 30 mg of starting material. The yield of 8a was 45^{ee}_{ee} (based on the amount of used starting material).

1- α -L-Xylofuranosylthymine (9a).—Compound 7a (600 mg, 1.34 mmoles) was dissolved in 20 ml of hot 50% aqueous EtOH (300 ml) and 4.1 ml of 1 N NaOH was added to the clear solution. The solution was refluxed for 30 min and processed as described for the preparation of $1-\alpha$ -L-arabinofuranosylthymine (3a). The syrup residue was dissolved in ca. 3 ml of EtOH and the solution was refrigerated. The crystalline product (242 mg, 70%)in two crops) had mp 192–194°. A single recrystallization afforded material with mp 194–195°, $\lambda_{\max}^{\text{H}_{2O}}$ 267 m μ , λ_{\min} 236.

Anal. Caled for C₁₀H₁₄N₂O₆: C, 46.51; H, 5.47; N, 10.85. Found: C, 46.28; H, 5.60; N, 10.87.

 $1-\alpha-L-Lyxofuranosylthymine$ (10a).—Compound 8a (600 mg, 1.05 mmoles) was dissolved in 20 ml of warm, 50% aqueous EtOH containing 3.5 ml of 1 N NaOH. The solution was refluxed for 30 min and processed as described in the preparation of 3a. The residue obtained was recrystallized twice from 5 ml of hot EtOH. The yield of pure material, mp 200-202°, was 200 mg (82%). The melting point and uv and nmr spectral data found for 10a were in close agreement with values reported⁹ for the enantiomorphic $1-\alpha$ -p-lyxofuranosylthymine. Furthermore, the ir spectra of these compounds were identical.

Anal. Caled for C₁₀H₁₄N₂O₆: C, 46.51; H, 5.47; N, 10.85. Found: C, 46.32; H, 5.58; N, 10.79.

 $1 \text{-} (Tri \text{-} \textit{O}\text{-} benzoyl \text{-} \alpha \text{-} \textbf{L}\text{-} arabino fur anosyl) \text{-} \mathcal{N}^4 \text{-} benzoyl cytosine$ (11) -- A mixture of 6.5 g of N4-benzoylcytosine15 and 4 g of Hg(CN)₂ was added to 3 l. of MeNO₂, and the suspension was dried azeotropically by distillation of 150 ml of solvent. Tri-Obenzoyl- α,β -L-arabinofuranosyl bromide (16 g) was added to the hot, stirred suspension, and the reaction mixture was refluxed. During the reflux period, 100 ml of solvent was distilled from the reaction mixture. After 2 hr of refluxing, another charge of $Hg(CN)_2$ (4 g) was added, and the mixture was dried azeotropically by distillation of some of the solvent. An additional 16 g of halogenose was added and refluxing was continued for a total of 4 hr. The clear reaction mixture was concentrated to dryness, and the residue was dissolved in CHCl₃. The CHCl₃ solution was washed with 30% freshly prepared aqueous KI and then with H_2O . The organic layer was dried (Na₂SO₄), filtered, and concentrated in vacuo to a syrup. In order to obtain seed crystals, a small amount of this syrup was dissolved in C_6H_6 -Et₂O (1:1). The solution was absorbed on a silica gel column and washed with C_6H_6 -Et₂O (1:1). The product was eluted from the column with EtOAc-MeOH(20;1) and the eluate evaporated to dryness. The syrup crystallized from hot EtOH. The syrup obtained by concentration of the CHCl₃ solution was dissolved in hot EtOH, and the solution was seeded with crystals of 11. After slow cooling to $\sim 40-50^{\circ}$, fine crystals of 11 were obtained (90% based on N^4 -benzoyleytosine, mp 197-198°). Recrystallization from EtOH gave pure product: mp 198–199°; $\lambda_{\rm max}^{\rm dif\,ErOH}$ 232, 260, 302 mµ; $\lambda_{\rm min}$ 249, 290 mµ.

Anal. Caled for $C_{37}H_{29}N_3O_9$: C, 67.37; H, 4.43; N, 6.37. Found: C, 67.43; H, 4.57; N, 6.40.

1- α -L-Arabinofuranosylcytosine (12).— Compound 11 (6.6 g) was added to a solution of 200 ml of EtOH and 100 ml of 1 N NaOH and the stirred mixture was heated for 1 hr at 75°. The solution was cooled to room temperature and added to a column of Dowex 50 (H⁺) (100 g wet weight). The column was washed with 50% aqueous EtOH (500 ml), then with H₂O (400 ml) to remove benzoic acid. The unblocked nucleoside 12 was eluted from the column with 2.6 N NH₄OH (400 ml) and the eluate was concentrated to dryness in vacuo. The residue was dissolved in 50 ml of hot EtOH and cooled overnight in the refrigerator. The crystalline material (2.0 g plus an additional 0.4 g obtained from the mother liquor) was recrystallized from hot EtOH.

The product, which was dried over P_2O_5 in vacuo at 80°, melted at 115-116° and showed $[\alpha]^{25}\mathfrak{v} = +2$ (c/3, H₂O), $\lambda_{\max}^{\mathrm{B}_{2O}} = 271 \text{ m}\mu$. $\lambda_{\min} 249 \ \mathrm{m}\mu$.

Anal. Caled for C₉H₄₈N₃O₅: C, 44.44; H, 5.38; N, 17.28. Found: C, 44.42; H, 5.47; N, 17.30.

1- α -L-Arabinofuranosyl- N^4 -benzoylcytosine (13). A solution of 1.5 g of 12 in 150 ml of MeOH was treated with 1.5 g of benzoic anhydride and refluxed for 5 hr. Additional charges of 1.5 g of anhydride were added at 1-hr intervals during the reflux period according to the procedure described.¹⁶ The hot reaction mixture was filtered from some insoluble material and the filtrate was concentrated in vacuo to a small volume which crystallized after treatment with Et₂O; yield 1.5 g (70%), mp 219-220°. Pure treatment with 154_{2} O; yield 1.5 g +70° /), hip 219-220°. Fure material, mp 228-229°, was obtained after recrystallization from hot MeOH; $\lambda_{\text{max}}^{\text{Ho}2}$ 258, 303 mµ; λ_{min} 230, 290 mµ. Anal. Calcd for C₁₆H₁₇N₃O₆; C, 55.32; H, 4.93; N, 42.09.

Found: C, 55.27: H, 4.90; N, 12.14.

1- α -L-Arabinofuranosvluracil (3b). Method A from 13. Compound 13 (1 g) was refluxed for 3 hr in 100 ml of 85% aqueous AcOH. The reaction mixture was evaporated to dryness in racuo at $\sim 40^\circ$, and the residue was dissolved in H₂O and added to a column containing Dowex 50 (H^+) . The column was eluted with H₂O until the eluate was free of uv-absorbing material. The eluate was concentrated to dryness in vacuo. The residue was dissolved in H₂O and washed with CHCl₃. The aqueous layer was evaporated to dryness in vacuo. The glassy residue was chromatographically pure (silica gel the in n-BuOH-H₂O, 86:14) but resisted all attempts at crystallization. The product, not analyzed, was converted directly to its tri-O-mesylate (4b).

Method B from 11. The fully benzoylated nucleoside (11, 5 g) was refluxed for 3 hr in 85% aqueous AcOH and the reaction mixture evaporated to dryness in vacuo at 40°. The residue was added to a solution of 50 ml of EtOH and 25 ml of 1 N NaOH and the mixture was heated for 1 hr at 70°. The reaction mixture was worked up as in method A above. A chromatographically pure glass, 1.7 g (89%), was obtained: $\lambda_{\max}^{\text{HgO}}$ 262 m μ , λ_{\min} 230 m μ . This product **3b** was used directly for the synthesis of **5b**.

2,2'-Anhydro-1-(3,5-di-O-mesyl- α -L-ribofuranosyl)uracil (5b). The glassy residue **3b** was azeotropically dried by addition of toluene and distillation of solvent in vacuo. The residue (from 37 g of blocked nucleoside 11) was then dissolved in 250 ml of dry pyridine, and the solution was cooled in an ice bath to $0-5^{\circ}$. The solution was treated dropwise with 33 g (twofold excess) of MeSO₂Cl, and the reaction mixture was allowed to remain overnight at room temperature, concentrated to a small volume, and poured into stirred ice-H₂O. The brown precipitate which formed was collected and washed several times with cold H₂O. The product (crude 4b) was dried (P_2O_3) to yield 20 g of a dark residue. A portion of crude residue (5.4 g) was dissolved in hot anhydrous MeOH (700 ml) and the mixture was decanted from a small amount of a black syrup. The cooled decantate was treated dropwise at room temperature with almost 1 equiv of 1 N NaOH. A fine crystalline product (3.7 g) was filtered after 1 hr and an additional 0.4 g was obtained from the mother liquor (total yield $94\frac{c_c}{c}$). After recrystallization from hot $80\frac{c_c}{c}$ aqueous EtOH, pure material was obtained with mp 250-251°; $\lambda_{\text{max}}^{\text{diff}}$ 225, 247 mµ; λ_{min} 234 mµ.

Anal. Caled for CuH14N2O5S2: C, 34.55; H, 3.69; N, 7.33; S. 16.77. Found: C, 34.70; H, 3.82; N, 7.26; S, 16.66.

 $2,2'-Anhydro-1-(3-O-mesyl-5-O-benzoyl-\alpha-L-ribofuranosyl)$ uracil (6b, X = O). Treatment of 2,2'-anhydro(3,5-di-O-mesyl- α -L-ribofuranosyl)uracil (5b) (3.0 g, 7.8 mmoles) with sodium benzoate (1.3 g, 9 mmoles) in dry DMF, as described above for the preparation of **6a**, afforded **6b** (94%, from hot EtOH), mp 233-234°, $\lambda_{\text{max}}^{\text{EOH}}$ 232 m μ , λ_{min} 208 m μ .

 $\begin{array}{c} \text{mp } 255 \ \text{Ext} + m_{\text{max}} \ 252 \ \text{mp}, \\ \text{Maal} \ \text{Calcd for } C_{17} \text{H}_{16} \text{N}_2 \text{Os}^2 \text{S}; \ \text{C}, \ 49.99; \ \text{H}, \ 3.95; \ \text{N}, \ 6.86; \\ \text{S}, \ 7.85; \ \text{Found}; \ \text{C}, \ 50.07; \ \text{H}, \ 4.05; \ \text{N}, \ 6.81; \ \text{S}, \ 7.93. \end{array}$

2.2'-Anhydro-1-(3-()-mesyl-5-()-benzoyl-a-L-ribofuranosyl)-4thiouracil (6b, X = S).--P₂S₆ (2.2 g, 0.01 mole) was added to 35 ml of dry pyridine, and the mixture was refluxed until solution occurred. Compound **6b** (X = O) (1.4 g, 3.7 mmoles) was added, and the solution was refluxed for 0.5 hr. The reaction mixture was concentrated in racno to a thin syrup which was then poured into stirred ice- H_2O . The brown solid material was filtered and washed well with cold H₂O. Recrystallization from hot 50 $_{\ell}^{c}$ aqueous EtOH gave 1.0 g of **6b** (X = S) (69 $_{\ell}^{c}$). A further recrystallization from the same solvent (including treatment with charcoal) gave fine yellow crystals: mp 200-201°: $\chi_{max}^{dif\,EtOH}$ 232, 328 m μ ; λ_{max} 212, 258 m μ .

Anal. Calcd for $C_{17}H_{16}N_2O_7S_2$: C, 48.10; H, 3.80; N, 6.60; S, 15.11. Found: C, 48.18; H, 3.97; N, 6.62; S, 15.09.

2,2'-Anhydro-1-(3,5-di-O-benzoyl- α -L-xylofuranosyl)uracil (7b) and Tri-O-benzoyl- α -L-lyxofuranosyluracil (8b).—A solution of 2.0 g (4.9 mmoles) of 6b (X = O) in 400 ml of dry DMF containing 3.0 g (20.8 mmoles) of sodium benzoate and 600 mg (4.9 mmoles) of benzoic acid was heated at 150° for 4.5 hr. The dark reaction mixture was processed as described previously for the preparation of 7a and 8a. Fractional crystallization of the residue from hot EtOH afforded 0.8 g (37%) of 7b which, after a further recrystallization, had mp 131-132°, $\lambda_{max}^{dil EtOH}$ 231 m μ , λ_{min} 212 m μ .

Anal. Calcd for $C_{23}H_{18}N_2O_7$: C, 63.59; H, 4.17; N, 6.44. Found: C, 63.63; H, 4.02; N, 6.39.

Compound **8b** was isolated from the mother liquors as described above in the preparation of **8a**. Fractions of the column effluent (EtOAc-MeOH, 15:1) containing mostly **8b** were combined and concentrated to dryness *in vacuo*. Recrystallization of the residue from EtOH-Et₂O afforded pure **8b**: mp 146-148°; $\lambda_{max}^{dit EtOH}$ 230, 256 m μ ; λ_{min} 211, 252 m μ .

Anal. Calcd for $C_{30}H_{24}N_2O_9$: C, 64.74; H, 4.34; N, 5.03. Found: C, 64.77; H, 4.46; N, 4.96.

1- α -L-Lyxofuranosyluracil (10b).—Compound 8b (0.1 g) was suspended in a solution of EtOH (10 ml) and 1 N NaOH (5.4 ml), and the mixture was warmed at 70° for 1 hr. The clear solution was treated with Dowex 50 (H⁺) and then filtered. The filtrate was evaporated to dryness *in vacuo*, and the residue was dissolved in a small amount of H₂O, then washed with CHCl₃ (three 20-ml portions). The H₂O layer was evaporated to dryness *in vacuo*, and the syrup residue was crystallized from EtOH. The yield of pure product, mp 202-203°, was 43 mg. For the enantiomorphic 1- α -D-lyxofuranosylthymine, Nishimura⁹ reported mp 203-204.5°. The uv and nmr spectral data were in close agreement with the values reported for the enantiomorph and the infrared spectra were identical.

Anal. Calcd for $C_9H_{12}N_2O_6$: C, 44.26; H, 4.95; N, 11.47. Found: C, 44.27; H, 4.89; N, 11.31. 2,2'-Anhydro-1-(3,5-di-O-benzoyl- α -L-xylofuranosyl)-4-thio-

2,2'-Anhydro-1-(3,5-di-O-benzoyl- α -L-xylofuranosyl)-4-thiouracil (17).—Compound 7b (1.5 g, 3.3 mmoles) was added to a refluxing solution of P₂S₅ (2.2 g, 10 mmoles) in 35 ml of pyridine. After 30 min of refluxing, the dark brown solution was evaporated to dryness *in vacuo* to a dark syrup which crystallized when triturated in cold H₂O. Recrystallization of the crude product (1 g, 64%) from 50% aqueous EtOH afforded yellow needles of 17: mp 236-237°; $\lambda_{max}^{di \ EtOH}$ 231, 283, 329 m μ , shoulder 275 m μ ; λ_{min} 258, 287 m μ .

Anal. Calcd for $C_{23}H_{18}N_2O_6S$: C, 61.32; H, 4.03; N, 6.22; S, 7.12. Found: C, 61.48; H, 4.05; N, 6.10; S, 7.15.

1- α -L-Xylofuranosylcytosine (16). Method A.—To a suspension of 800 mg (1.77 mmoles) of 17 in 100 ml of 50% aqueous EtOH was added 1 N NaOH (10 ml). The mixture was warmed

at 70° for 5 min and then allowed to cool. After 1 hr, the clear yellow solution was treated with Dowex 50 (H^+) (13 g of moist resin). The resin was removed by filtration and the vellow filtrate was concentrated to dryness. A solution of the residue in H_2O was extracted with CHCl₃ (three 50-ml portions). Concentration of the aqueous layer afforded a yellow amorphous solid (18) $(\lambda_{\max}^{\text{difEtoH}} 332 \text{ m}\mu, \lambda_{\min} 250 \text{ m}\mu)$ which was used directly for the synthesis of 17. A sealed tube containing the above product (18) in 40 ml of ethanolic NH₃ was heated at 100° for 17 hr. The redbrown reaction mixture was evaporated to dryness in vacuo at 40°, and the residue was dissolved in 100 ml of H_2O . This solution was added to a column of Dowex 50 (H+) (10 g of moist resin), and the column was eluted with 400 ml of H_2O . Compound 16 was eluted from the column by 400 ml of 2 N NH4OH. The effluent was treated with charcoal and evaporated in vacuo. Crystallization of the product occurred in the concentrated solution. The solid (310 mg) was collected and recrystallized from dilute EtOH. The over-all yield from 17 was 68-70%. Pure **16** had mp 267–268° dec and $[\alpha]^{25}D + 118° (c \ 0.1, H_2O); \lambda_n^{\text{H}}$ 273, shoulder 225 mµ; λ_{\min} 246 mµ. The spectra at various pH values were closely similar to those previously reported for $1-\beta$ p-ribofuranosylcytosine.8

Anal. Caled for C_8H_1_3N_3O_5: C, 44.43; H, 5.38; N, 17.27. Found: C, 44.56; H, 5.33; N, 17.32.

Method B.-Treatment of a solution of 2.0 g (7b) in 50% aqueous EtOH with NaOH and isolation of the product as described previously for the preparation of 9a afforded 1.2 g of syrup which contained 9b. The syrup was dissolved in 20 ml of pyridine and dried by azeotropic distillation of the solvent. The syrupy residue was dissolved in 10 ml of Ac₂O containing 1 ml of pyridine, and the solution was heated at 100° for 1 hr. Removal of the solvents in vacuo left a syrup containing the triacetate 14. The syrup in 100 ml of pyridine was treated with 4.2 g of P_2S_5 and 0.09 ml of H_2O . The stirred mixture was refluxed for 6 hr and concentrated in vacuo to dryness. The residue was triturated with 50 ml of cold H₂O and the resulting brown solid was removed by filtration. The solid was dissolved in 300 ml of CHCl₃ and washed with H₂O. Concentration of the organic layer afforded semisolid material which contained thione 15. Crude 15 in 80 ml of ethanolic NH₃ was heated in a sealed tube at 100° for 20 hr, and the brown reaction mixture was processed as described in method A above. The yield of pure 16, mp and mmp 267-268°, was 300 mg (over-all yield from 7b, 27%). The product had uv absorption characteristics identical with a sample prepared according to method A above.

Acknowledgment.—The authors express their appreciation to Dr. T. Nishimura for samples of $1-\alpha$ -D-lyxofuranosylthymine and $1-\alpha$ -D-lyxofuranosyluracil used in this study.