

solution and extracted with CHCl_3 . The extract was washed with H_2O , dried on K_2CO_3 and distilled *in vacuo* to give the crude compound (XI), whose recrystallization from EtOH gave 100 mg. of XI as colorless plates, m.p. 122~124°. Beilstein test of this compound was negative. IR cm^{-1} (KBr): ν_{NH} 3440, 3340; $\nu_{\text{C=N}}$ 1622; δ_{NH} 1642. NMR (τ) (in CDCl_3): 7.20 (3H, singlet, $\text{C}_1\text{-Me}$); 5.50~6.00 (2H, broad, NH_2); 3.47 (1H, singlet, $\text{C}_4\text{-H}$). Anal. Calcd. for $\text{C}_{10}\text{H}_{10}\text{N}_2$: C, 75.92; H, 6.38; N, 17.71. Found: C, 75.72; H, 6.42; N, 17.96. Recrystallization of the picrate from EtOH gave yellow needles, m.p. 232~233° (decomp.). Anal. Calcd. for $\text{C}_{10}\text{H}_{10}\text{N}_2 \cdot \text{C}_6\text{H}_3\text{O}_7\text{N}_3$ (XI): C, 49.62; H, 3.38; N, 18.08. Found: C, 49.75; H, 3.64; N, 17.65.

b) Ammonolysis of 1.0 g. of Xb under the same conditions as the method a) gave 120 mg. of XI. Furthermore, ammonolysis of VI under the same conditions as the method a) also afforded 0.4 g. (25%) of XI. Both specimens were identical with the sample obtained by the procedure a) by mixed melting point test and infrared spectrum.

N-[3-(1-methyloquinolyl)]acetamide (XIa)—A mixture of 100 mg. of XI with an excess of Ac_2O was heated on a water-bath for 1 hr. After the excess of Ac_2O was removed by distillation *in vacuo*, the residue was basified with 10% aq. NaOH solution and extracted with CHCl_3 . The extract was washed with H_2O , dried on Na_2SO_4 and distilled. Recrystallization of the resultant residue from *iso*-PrOH gave 70 mg. of XIa as colorless plates, m.p. 202~203°. Anal. Calcd. for $\text{C}_{12}\text{H}_{12}\text{ON}_2$ (XIa): C, 71.84; H, 6.04; N, 13.99. Found: C, 72.02; H, 6.24; N, 13.35. IR cm^{-1} (KBr): ν_{NH} 3220; $\nu_{\text{C=O}}$ 1658; $\nu_{\text{C=N}}$ 1625 (shoulder).

We thank President A. Yanagisawa, Mr. O. Takagi, and Grelan Pharmaceutical Co., Ltd., for their grateful assistance. We also thank Mrs. F. Seto, Miss N. Nanjo, and Miss R. Kobayashi for microanalyses.

[Chem. Pharm. Bull.
15(5) 707~710 (1967)]

UDC 547.964.07

Kazumi Ogata and Satoru Ishii: Syntheses of Ophthalmic Acid and its Analogues.*¹

(Research Laboratory, Senju Pharmaceutical Co., Ltd.*²)

(Received March 18, 1966)

In 1956, Waley^{1~3)} isolated an acidic tripeptide, γ -L-glutamyl-L- α -amino-*n*-butyryl-glycine, named ophthalmic acid, and an analogous tripeptide, γ -L-glutamyl-L-alanyl-glycine, named norophthalmic acid from calf lens. Afterward, he synthesized⁴⁾ ophthalmic acid by the reaction of N-benzoyloxycarbonyl-L- γ -glutamyl azide with L- α -amino-*n*-butyryl-glycine. Enzymatic synthesis of ophthalmic acid was reported by the same authors.⁵⁾

At present, several syntheses of ophthalmic, norophthalmic acids and their analogue are reported. Kermack, *et al.*⁶⁾ synthesized DL-norophthalmic acid (γ -DL-glutamyl-DL-alanylglycine) from phthalylglutamic anhydride and alanylglycine. He obtained the optically active acid by the treatment of glutathione with Raney Nickel in poor yield. By using α -*tert* butyl N-benzoyloxycarbonylglutamate and (α -amino-*n*-butyryl)glycine *tert* butyl ester, Taschner, *et al.*⁷⁾ synthesized ophthalmic acid. Shchukina, *et al.*⁸⁾

*¹ Presented at the 21th Meeting of the Pharmaceutical Society of Japan, at Tokushima, October, 1965 and the 4th Peptide Symposium, at Institute for Protein Research, University of Osaka, December, 1965.

*² Iwazono-cho 416, Ashiya, Hyogo Pref. (緒方一美, 石井 覚).

1) S. G. Waley: Biochem. J., **64**, 715 (1956).

2) *Idem*: *Ibid.*, **67**, 172 (1957).

3) *Idem*: Proc. Chem. Soc., **1957**, 25.

4) S. G. Waley: Biochem. J., **68**, 189 (1958).

5) E. E. Cliffe, S. G. Waley: *Ibid.*, **79**, 669 (1961).

6) W. O. Kermack, N. A. Matheson: *Ibid.*, **65**, 45 (1957).


7) E. Taschner, T. Sokolowska, J. F. Biernat: Collection Czch. Chem. Commun., **27**, 2237 (1962).

8) L. A. Shchukina, A. L. Jouse: "Peptides," Ed. by G. T. Young, p. 189 (1963), Pergamon Press, London.

prepared the same acid with *p*-nitrobenzyloxycarbonylglutamate and α -amino-*n*-butyrylglycine *p*-nitrobenzyl ester. Losse, *et al.*⁹⁾ prepared γ -L-glutamyl-DL- α -amino-*n*-butyrylglycine and γ -L-glutamyl-L-alanylglycine in an active ester method.

The concentration of ophthalmic acid was reported as approximately 0.2 mg. per gram of the lens. This is about one-tenth that of glutathione content in the lens. It seems interest that the two tripeptides, ophthalmic and norophthalmic acid, structurally related to glutathione were found, by the present time, only a few animal tissues, though glutathione distributes in almost tissues of animals and plants. Their physiological functions are still unknown.

In order to explore their biological activities, we are in need of these compounds. Here we wish to record an alternate syntheses of ophthalmic acid and norophthalmic acid. In addition, the threonine analogue of ophthalmic acid was also prepared. Synthetic schema of γ -L-glutamyl-L- α -amino-*n*-butyrylglycine, γ -L-glutamyl-L-alanylglycine and γ -L-glutamyl-L-threonylglycine are illustrated in Chart 1. The synthetic procedures employed are: (1) coupling of α -benzyl N-benzyloxycarbonyl-L-glutamate¹⁰⁾ with dipeptides in a mixed anhydride method, (2) followed by catalytic hydrogenation over a Pd-C catalyst to obtain the γ -L-glutamyl-L-peptides.



Cbz = Benzyloxycarbonyl, Bz = Benzyl, DCC = Dicyclohexylcarbodiimide.

Chart 1.

N-Benzyloxycarbonyl-L- α -amino-*n*-butyric acid (Ia)⁴⁾ and N-benzyloxycarbonyl-L-alanine (Ib)^{11,12)} were reacted with glycine benzyl ester (II) by dicyclohexylcarbodiimide

- 9) G. Losse, H. Jeschkeit, R. Höhn : Ann., **676**, 222 (1964).
- 10) E. Klieger, H. Gibian : *Ibid.*, **655**, 195 (1962).
- 11) M. Bergmann, L. Zervas : Ber., **65**, 1192 (1932).
- 12) M. Hunt, du Vigneaud : J. Biol. Chem., **124**, 699 (1938).

to give N-benzyloxycarbonyl-L- α -amino-n-butyrylglycine benzyl ester (IIIa) and N-benzyloxycarbonyl-L-alanylglycine benzyl ester (IIIb)¹⁴ respectively. These compounds were hydrogenated catalytically over a Pd-C catalyst, to give the corresponding dipeptides (IVa)⁴ and (IVb)¹⁴⁻²⁰ respectively. Then, these dipeptides were reacted with α -benzyl N-benzyloxycarbonyl-L-glutamate (V)¹⁰ by the mixed anhydride method to give α -benzyloxycarbonyl-L-glutamyl-L-peptide (VIa) and (VIb) respectively. Finally, catalytic hydrogenation of (VIa) and (VIb) gave γ -L-glutamyl-L- α -amino-n-butyrylglycine (ophthalmic acid) (VIIa) and γ -L-glutamyl-L-alanylglycine (norophthalmic acid) (VIIb) in nearly quantitative yield. These synthetic compounds, (VIIa) and (VIIb), exhibited a single spot on paper chromatography. The acid hydrolysates gave respectively the constituent amino acids in the ratios predicted by theory. All physical constants were quite identical with those reported in the literatures.^{4,6,9} We have also prepared the threonine analogue of ophthalmic acid in essentially the same method stated above starting with N-benzyloxycarbonyl-L-threonine (Ic).¹³ Examination of the product by paper chromatography and the result of amino acid analysis seems to verify the conclusion that the synthetic compound is homogeneous. Biological examination of those compounds is under progress, these results will be reported in future.

Experimental*3

N-Benzyloxycarbonyl-L- α -amino-n-butyrylglycine Benzyl Ester (IIIa)—N-Benzyloxycarbonyl-L- α -amino-n-butyric acid (8.8 g.), glycine benzyl ester-toluene-*p*-sulfonate (10.9 g.) and dry triethylamine (3.3 g.) were dissolved in anhydrous tetrahydrofuran (70 ml.). To this ice-cooled solution, dicyclohexylcarbodiimide (6.7 g.) was added. The mixture was stirred for 1 hr. and left overnight. Dicyclohexylurea formed during the reaction was filtered off and the filtrate was evaporated *in vacuo*. The resulting oil was dissolved in AcOEt, which was washed successively with 2% HCl, 4% NaHCO₃ and H₂O, and dried over anhydrous Na₂SO₄. The solvent was removed by evaporation. The resulting solid was washed with light petroleum, collected by filtration and recrystallized from a mixture of MeOH and H₂O; yield 9.6 g. (77%). m.p. 115~116°, $[\alpha]_D^{25}$ -26.5° (c=4.0, MeOH). *Anal.* Calcd. for C₂₁H₂₄O₅N₂: C, 65.6; H, 6.3; N, 7.3. Found: C, 65.6; H, 6.4; N, 7.2.

N-Benzyloxycarbonyl-L-alanylglycine Benzyl Ester (IIIb)—This compound was prepared from N-benzyloxycarbonyl-L-alanine (11.2 g.), glycine benzyl ester-toluene-*p*-sulfonate (16.9 g.), triethylamine (5.1 g.) and dicyclohexylcarbodiimide (10.3 g.) in essentially the same manner as described above in the preparation of IIIa; yield 15.0 g. (81%), m.p. 111~112°, [lit.⁹] m.p. 111°, $[\alpha]_D^{25}$ -23.4° (c=4.0, MeOH). *Anal.* Calcd. for C₂₀H₂₂O₅N₂: C, 64.9; H, 6.0; N, 7.6. Found: C, 65.0; H, 6.2; N, 7.5.

N-Benzyloxycarbonyl-L-threonylglycine Benzyl Ester (IIIc)—This compound was prepared from N-benzyloxycarbonyl-L-threonine (8.0 g.), glycine benzyl ester-toluene-*p*-sulfonate (10.7 g.), triethylamine (3.2 g.) and dicyclohexylcarbodiimide (6.5 g.) in essentially the same manner as described above in the preparation of IIIa; yield 10.0 g. (79%), m.p. 102~104°, $[\alpha]_D^{25}$ -13.1° (c=4.9, MeOH). *Anal.* Calcd. for C₂₁H₂₄O₅N₂: C, 63.0; H, 6.0; N, 7.0. Found: C, 62.7; H, 6.2; N, 7.0.

L- α -Amino-n-butyrylglycine (IVa)—N-Benzyloxycarbonyl-L- α -amino-n-butyrylglycine benzyl ester (6.7 g.) was hydrogenated over a Pd-C catalyst in a mixture of 10% *tert*-BuOH (150 ml.) and AcOH (2 ml.) until evolution of CO₂ ceased. The catalyst was removed by filtration and the filtrate evaporated *in vacuo*. The residual oil was crystallized by addition of MeOH. The product was collected and recrystallized from aqueous MeOH; yield 2.3 g. (83%), m.p. 225~227° (decomp.), $[\alpha]_D^{25}$ +82.5° (c=5.0, H₂O), [lit.⁴] m.p. 225° (decomp.), $[\alpha]_D^{18}$ +72°, Rf 0.78 (phenol-0.2% NH₄OH). *Anal.* Calcd. for C₈H₁₂O₃N₂: C, 45.0; H, 7.6; N, 17.5. Found: C, 45.3; H, 7.7; N, 17.3.

*3 All melting points are uncorrected.

13) R. B. Merrifield: *Ibid.*, **232**, 43 (1958).

14) B. F. Erlanger, E. Brand: *J. Am. Chem. Soc.*, **73**, 3508 (1951).

15) E. Fischer: *Ber.*, **41**, 850 (1908); *Ibid.*, **38**, 2914 (1905).

16) J. L. Bailey: *Nature*, **164**, 889 (1949); *J. Chem. Soc.*, **1950**, 3461.

17) M. Bergmann, L. Zervas, J. S. Fruton, F. Schneider, H. Schleich: *J. Biol. Chem.*, **109**, 325 (1935).

18) L. Zervas, D. M. Theodoropoulos: *J. Am. Chem. Soc.*, **78**, 1359 (1956).

19) W. Grassmann, E. Wunsch: *Ber.*, **91**, 449 (1958).

20) E. Brand, B. F. Erlanger: *J. Am. Chem. Soc.*, **72**, 3314 (1950).

L-Alanylglycine (IVb)—This compound was prepared from N-benzyloxycarbonyl-L-alanylglycine benzyl ester (13.4 g.) by hydrogenation as described above in the preparation of IVa; yield 4.9 g. (93%), m.p. 232~234°(decomp.), $[\alpha]_D^{25} + 49.5^\circ$ (c=2.0, H₂O), [lit. ¹⁴] $[\alpha]_D^{25} + 50.6^\circ$, lit. ¹⁸) $[\alpha]_D^{25} + 51.3^\circ$, lit. ¹⁶) $[\alpha]_D^{25} + 49.1^\circ$, lit. ¹⁵) $[\alpha]_D^{25} + 50.2^\circ$, lit. ¹⁹) $[\alpha]_D^{17} + 51.44^\circ$, Rf 0.70 (phenol-0.2% NH₄OH). *Anal.* Calcd. for C₅H₁₀O₃N₂: C, 41.1; H, 6.9; N, 19.2. Found: C, 40.9; H, 7.1; N, 19.2.

L-Threonylglycine (IVc)—This compound was prepared from N-benzyloxycarbonyl-L-threonylglycine benzyl ester (7.0 g.) by hydrogenation as described above in the preparation of IVa; yield 2.5 g. (81%), m.p. 244~246°(decomp.), $[\alpha]_D^{25} + 53.7^\circ$ (c=3.7, H₂O), Rf 0.63 (phenol-0.2% NH₄OH). *Anal.* Calcd. for C₆H₁₂O₄N₂: C, 40.9; H, 6.9; N, 15.9. Found: C, 41.0; H, 7.2; N, 15.8.

α -Benzyl N-Benzyloxycarbonyl-L-glutamate (V)—This compound was prepared according to the method of Klieger, *et al.*¹⁰); m.p. 98~99°, $[\alpha]_D^{25} - 23.8^\circ$ (c=2.0, MeOH), [lit. ¹⁰] m.p. 97~98°, $[\alpha]_D^{25} - 23.8^\circ$. *Anal.* Calcd. for C₂₀H₂₁O₆N: C, 64.7; H, 5.7; N, 3.8. Found: C, 64.8; H, 5.9; N, 4.0.

α -Benzyl N-Benzyloxycarbonyl-L-glutamyl-L- α -amino-n-butyrylglycine (VIa)—A mixed anhydride was prepared in an ice-bath from α -benzyl N-benzyloxycarbonyl-L-glutamate (4.6 g.) in anhydrous tetrahydrofuran (30 ml.) with triethylamine (1.3 g.) and ethyl chloroformate (1.4 g.). This solution was added slowly with stirring to a chilled solution of L- α -amino-n-butyrylglycine (2.0 g.) and triethylamine (1.3 g.) in H₂O (10 ml.). The mixture was stirred in an ice-bath for 1 hr. and at room temperature for 2 hr. The solvent was evaporated *in vacuo*. The residue was acidified with 1N HCl and the resulting precipitate was collected by filtration, washed successively with H₂O and ether, and recrystallized from MeOH; yield 4.1 g. (64%), m.p. 178~179°, $[\alpha]_D^{25} - 25.0^\circ$ (c=3.3, AcOH). *Anal.* Calcd. for C₂₆H₃₁O₈N₃: C, 60.8; H, 6.1; N, 8.2. Found: C, 61.0; H, 6.4; N, 8.2.

α -Benzyl N-Benzyloxycarbonyl-L-glutamyl-L-alanylglycine (VIb)—This compound was prepared from α -benzyl N-benzyloxycarbonyl-L-glutamate (3.7 g.) and L-alanylglycine (1.5 g.) by the mixed anhydride method as described above in the preparation of VIa; yield 3.2 g. (64%), m.p. 191~192°, $[\alpha]_D^{25} - 26.9^\circ$ (c=2.0, AcOH). *Anal.* Calcd. for C₂₅H₂₉O₈N₃: C, 60.1; H, 5.9; N, 8.4. Found: C, 60.3; H, 6.0; N, 8.4.

α -Benzyl N-Benzyloxycarbonyl-L-glutamyl-L-threonylglycine (VIc)—This compound was prepared from α -benzyl N-benzyloxycarbonyl-L-glutamate (3.7 g.) and L-threonylglycine (1.8 g.) by the mixed anhydride method as described above in the preparation of VIa. It was recrystallized from EtOH; yield 3.3 g. (62%), m.p. 148~149°, $[\alpha]_D^{25} - 16.0^\circ$ (c=2.0, AcOH). *Anal.* Calcd. for C₂₆H₃₁O₈N₃: C, 59.0; H, 5.9; N, 7.9. Found: C, 58.8; H, 6.0; N, 8.0.

γ -L-Glutamyl-L- α -amino-n-butyrylglycine (ophthalmic acid) (VIIa)— α -Benzyl N-benzyloxycarbonyl-L-glutamyl-L- α -amino-n-butyrylglycine (3.0 g.) was hydrogenated over a Pd-C catalyst in a mixture of 10% (v/v) *tert*-BuOH (100 ml.) and AcOH (2 ml.) until the evolution of CO₂ ceased. The catalyst was removed by filtration and the filtrate was evaporated *in vacuo*. The residual oil precipitated by addition of Me₂CO was collected and recrystallized from H₂O by addition of Me₂CO; yield 1.2 g. (68%), m.p. 177~179°(decomp.), $[\alpha]_D^{25} - 28.4^\circ$ (c=2.1, H₂O), [lit. ⁴] $[\alpha]_D^{25} - 29^\circ$, Rf 0.59 (phenol-0.2% NH₄OH), single ninhydrin positive spot. Amino acid ratios in an acid hydrolysate Glu_{1.00} α -amino-n-but_{1.01}Gly_{0.98} (average recovery 92%). *Anal.* Calcd. for C₁₁H₁₉O₆N₃·H₂O: C, 43.0; H, 6.9; N, 13.7. Found: C, 43.2; H, 7.3; N, 13.8.

γ -L-Glutamyl-L-alanylglycine (norophthalmic acid) (VIIb)—This compound was prepared from α -benzyl N-benzyloxycarbonyl-L-glutamyl-L-alanylglycine (5.0 g.) by hydrogenation as described above in the preparation of VIIa; yield 2.0 g. (71%), m.p. 197~199°(decomp.), $[\alpha]_D^{25} - 29.8^\circ$ (c=2.4, H₂O), Rf 0.51 (phenol-0.2% NH₄OH), single ninhydrin positive spot. Amino acid ratios in an acid hydrolysate Glu_{1.00}Ala_{1.08}Gly_{0.93} (average recovery 93%). *Anal.* Calcd. for C₁₀H₁₇O₆N₃· $\frac{1}{2}$ H₂O: C, 42.2; H, 6.4; N, 14.8. Found: C, 41.8; H, 6.6; N, 14.8.

γ -L-Glutamyl-L-threonylglycine (VIIc)—This compound was prepared from α -benzyl N-benzyloxycarbonyl-L-glutamyl-L-threonylglycine (2.5 g.) by hydrogenation as described above in the preparation of VIIa; yield 1.0 g. (68%), $[\alpha]_D^{25} - 16.5^\circ$ (c=2.9, H₂O), Rf 0.38 (phenol-0.2% NH₄OH), single ninhydrin positive spot. Amino acid ratios in an acid hydrolysate Glu_{1.02}Thr_{1.00}Gly_{0.98} (average recovery 80%). *Anal.* Calcd. for C₁₁H₁₉O₇N₃· $\frac{1}{2}$ H₂O: C, 42.0; H, 6.4; N, 13.4. Found: C, 42.2; H, 6.3; N, 13.3.

The authors are deeply indebted to associate professor H. Yajima of the University of Kyoto for his courteous and helpful guidance and encouragement throughout the experiment, and also indebted to the members of the Elemental Analytical Center of Faculty of Pharmaceutical Sciences, University of Kyoto and University of Osaka for elemental analyses.