

Antidiabetic Ellagitannins from Pomegranate Flowers: Inhibition of α -Glucosidase and Lipogenic Gene Expression


Tao Yuan,[†] Yuanqing Ding,[‡] Chunpeng Wan,[†] Liya Li,[†] Jialin Xu,[†] Ke Liu,[§]
Angela Slitt,[†] Daneel Ferreira,^{*,‡} Ikhlas A. Khan,^{*,‡} and Navindra P. Seeram^{*,†}

Department of Biomedical and Pharmaceutical Sciences, College of Pharmacy,
University of Rhode Island, Kingston, Rhode Island 02881, United States,
National Center for Natural Products Research and Department of Pharmacognosy,
School of Pharmacy, University of Mississippi, University, Mississippi 38677,
United States, and Department of Medicinal Chemistry, University of Kansas,
Lawrence, Kansas 66045, United States

dferreir@olemiss.edu; ikhan@olemiss.edu; nseeram@uri.edu

Received September 15, 2012

ABSTRACT



Two new ellagitannins containing a rare 3-oxo-1,3,3a,8b-tetrahydrofuro[3,4-*b*]benzofuran moiety, namely punicatannins A (1) and B (2), were isolated from pomegranate (*Punica granatum*) flowers. Their structures with absolute configuration were determined by detailed analysis of spectroscopic data, electronic circular dichroism (ECD) calculation, and chemical hydrolysis. A plausible biogenetic route involving a key enzymatic 1,4-Michael addition is proposed. Punicatannin A showed potent inhibition of α -glucosidase and lipogenic gene expression.

The prevalence of diabetes is increasing rapidly worldwide. The International Diabetes Federation predicts that over 435 million people worldwide will be struggling with this disease by 2030 if no effective prevention and control programs are implemented. This global pandemic is driven by type 2 diabetes which contributes to the vast majority of all cases of diabetes.¹ While several synthetic drugs including sulfonylureas, biguanides, thiazolidinediones, and α -glucosidase inhibitors are available to treat type 2 diabetes, many have limited efficacy with significant associated side

effects. This has led to the search for more effective compounds with less side effects, particularly natural products and dietary agents, for the management of type 2 diabetes.

Punica granatum L. (Punicaceae), commonly known as pomegranate, is commercially cultivated for its edible fruit throughout the drier parts of Southeast Asia, the Mediterranean region, and the United States. The flowers of this plant were used as a remedy for diabetes in the Unani and Ayurvedic medicinal systems.² Moreover, a pomegranate flower extract was reported to markedly lower blood

[†] University of Rhode Island.

[‡] University of Mississippi.

[§] University of Kansas.

(1) Zimmet, P.; Alberti, K. G.; Shaw, J. *Nature* **2001**, 414, 782–787.

(2) Huang, T. H. W.; Peng, G.; Kota, B. P.; Li, G. Q.; Yamahara, J.; Roufogalis, B. D.; Li, Y. *Toxicol. Appl. Pharmacol.* **2005**, 207, 160–169.

(3) Li, Y.; Wen, S.; Kota, B. P.; Peng, G.; Li, G. Q.; Yamahara, J.; Roufogalis, B. D. *J. Ethnopharmacol.* **2005**, 99, 239–244.

glucose levels in the Zucker diabetic fatty rat model.³ However, there is limited knowledge on pomegranate flower constituents⁴ and the bioactive antidiabetic compounds present therein remain unknown.

In continuation of our characterization of new antidiabetic compounds from natural sources,⁵ this study sought to identify the bioactive compounds present in pomegranate flowers. Preliminary screening of a pomegranate flower extract showed promising α -glucosidase inhibitory activity ($IC_{50} = 3.2 \mu\text{g/mL}$). Subsequent bioassay-guided isolation afforded two unprecedented ellagitannins, named punica-tannins A (**1**) and B (**2**), both containing a rare 3-oxo-1,3,3a,8b-tetrahydrofuro[3,4-*b*]benzofuran moiety attached to a (¹C₄)-glucopyranose core, along with a biosynthetically related compound, isocorilagin (**3**).⁶ Herein we present the details of the isolation and structure elucidation of **1** and **2**, the evaluation of their antidiabetic activities, and a plausible biosynthetic pathway for these two compounds.




Figure 1. ¹H–¹H COSY (bold line) and key HMBC correlations (H→C) of **1**.

Compound **1** was isolated as a tan amorphous powder. The standard color reaction with ferric chloride (dark blue) and sodium nitrite–acetic acid (reddish brown) suggested that compound **1** was most likely an ellagitannin. Its molecular formula was established as C₄₃H₃₄O₂₈ with an index of hydrogen deficiency of 27 based on the HR-ESI-MS exhibiting an [M–H][–] ion at *m/z* 997.1151 (calcd for C₄₃H₃₃O₂₈, 997.1158). The ¹H and ¹³C NMR spectra [Supporting Information (SI) Table S2] of **1** showed signals due to a highly acylated hexopyranosyl moiety along with aromatic and carboxylic carbon resonances, strongly indicative of compound **1** being a hydrolyzable tannin. The presence of galloyl and hexahydroxydiphenoyl (HHDP) groups was readily determined by the ¹H NMR signals at $\delta_{\text{H}} = 7.08$ (2H, s), $\delta_{\text{H}} = 6.72$ (s), 6.83 (s) and characteristic ¹³C NMR signals in the aromatic region. The

The remaining part of the molecule was established by analysis of 1D and 2D NMR data. Apart from the above functionalities, carbon resonances for six aromatic carbons (δ_{C} 112.7, 118.6, 119.1, 136.4, 147.8, 148.6), four ester carbonyls (δ_{C} 165.4, 169.9, 171.1, 175.0), two methines (δ_{C} 51.5, 80.9), an oxygenated quaternary (δ_{C} 89.6), one methylene (δ_{C} 39.3), and two *O*-methyls (δ_{C} 52.8, 53.0) were observed in the ¹³C NMR spectrum. The chemical shifts of the aromatic carbon signals implied the presence of a pyrogallol ring. The HSQC spectrum permitted assignment of the protons to their bonding carbons, and the ¹H–¹H COSY and HMBC spectra (Figure 1) were applied to construct the structure of the remaining part of the molecule. The ¹H–¹H COSY correlation between H-8b'' and H-1'' revealed the linkage of C-8b'' and C-1''. In the HMBC spectrum, the mutual ¹H–¹³C correlation between an aromatic singlet ($\delta_{\text{H}} = 7.12$, H-7'') and the C-12'' ester carbonyl carbon (δ_{C} 165.4), C-5'', and C-8a'', indicated that the C-12'' ester carbonyl was linked at C-8'' of a pyrogallol-type ring. The HMBC correlations from H-8b'' to C-4a'', C-8'', and C-8a'', suggested that the CH-8b'' methine group was attached at C-8a'' of the pyrogallol-type ring. The HMBC correlations from H₂-10'' to C-3'' (δ_{C} 175.0), C-3a'' (δ_{C} 89.6), and C-8b'' (δ_{C} 51.5) indicated the contiguous linkage of C-3'', C-3a'', and C-8b'' and also

(4) (a) Xie, Y.; Morikawa, T.; Ninomiya, K.; Imura, K.; Muraoka, O.; Yuan, D.; Yoshikawa, M. *Chem. Pharm. Bull.* **2008**, *56*, 1628–1631. (b) Wang, R.; Wang, W.; Wang, L.; Liu, R.; Ding, Y.; Du, L. *Fitoterapia* **2006**, *77*, 534–537.

(5) (a) Yuan, T.; Wan, C.; Liu, K.; Seeram, N. P. *Tetrahedron* **2012**, *68*, 959–964. (b) Wan, C.; Yuan, T.; Li, L.; Kandhi, V.; Cech, N. B.; Xie, M.; Seeram, N. P. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 597–600.

(6) Liu, X.; Cui, C.; Zhao, M.; Wang, J.; Luo, W.; Yang, B.; Jiang, Y. *Food Chem.* **2008**, *109*, 909–915.

(7) Immel, S.; Khanbabaee, K. *Tetrahedron: Asymmetry* **2000**, *11*, 2495–2507.

allowed the attachment of CH₂-10'' to C-3a''. A methoxycarbonyl group was attached to both C-10'' and C-1'' based on the HMBC correlations from H₂-10'' and the *O*-methyl proton signals ($\delta_{\text{H}} = 3.79$) to C-11'' and from H-8b'', H-1'', and the *O*-methyl proton signals ($\delta_{\text{H}} = 3.31$) to C-9''. The linkage of C-1'' and C-3'' via oxygen was determined by the HMBC correlation between H-1'' and C-3''. The index of hydrogen deficiency of 26 in compound **1** was accounted for by the isocorilagin moiety (17), four carbonyl groups (4), a pyrogallol ring (4), and a furan ring (1). The remaining index of hydrogen deficiency required the presence of an additional ring. Analysis of the ¹³C NMR data suggested the presence of an ether bridge between oxygenated quaternary carbon C-3a'' and aromatic carbon C-4a'' to furnish a ring. Thus, the molecular structure of the remaining part was determined as the rare 5,6-dihydroxy-3a-(2-methoxy-2-oxoethyl)-1-(methoxycarbonyl)-3-oxo-1,3,3a,8b-tetrahydrofuro[3,4-*b*]benzofuran-8-carboxylic acid moiety.⁸ Finally, the HMBC correlation between H-4 and C-12'' facilitated the placement of this structural unit at C-4 of the glucopyranosyl core via an ester bridge.


The relative configuration of the 5,6-dihydroxy-3a-(2-methoxy-2-oxoethyl)-1-(methoxycarbonyl)-3-oxo-1,3,3a,8b-tetrahydrofuro[3,4-*b*]benzofuran-8-carboxylic acid moiety was determined by the ROESY spectrum and the H-1''–H-8b'' coupling constant (9.6 Hz) reminiscent of the *cis*-configuration of these protons.⁹ The ROESY correlation between H-8b'' and H-10'' indicated that H-8b'' and the C-3a'' 2-methoxy-2-oxoethyl group were cofacial.

A combination of electronic circular dichroism (ECD) analyses, chemical hydrolysis, and calculated ECD spectra were used to elucidate the absolute configuration of compound **1**. First, the HHDP moiety was assigned an *R*-configuration by the negative Cotton effect at 234 nm. Notably, a Cotton effect at 235 nm is a diagnostic criterion in determining the absolute configuration of HHDP units in ellagitannins, which is unaffected by the presence or absence of galloyl groups and the conformation of the glucose core.¹⁰ Acid hydrolysis of **1** afforded D-glucose, which was identified by direct comparison with an authentic sample (SI S1). However, the challenging issue to determine the absolute configuration of the acyl group linked at C-4 in compound **1** still remained. Hydrolysis of **1** in hot water yields **1a**, which was confirmed by the ¹H NMR and mass spectra (SI S20 and S21). As the absolute configuration of **1a** could not be resolved directly by the analysis of its ECD curve, comparison between experimental and calculated ECD spectra using the time-dependent DFT method was successfully applied to resolve this

issue.¹¹ The mandatory assigned (1''*R*, 3a''*R*, 8b''*R*)-configuration was selected for the conformational search of **1a** by using the OPLS2005 force field in the Schrodinger 9.2 MacroModel software package. After the detailed calculated ECD analysis (SI S4.1), the (1''*R*, 3a''*R*, 8b''*R*) absolute configuration of partial structure **1a** was unequivocally confirmed.

Compound **2** was isolated from the same fraction as **1** and had the same molecular formula (C₄₃H₃₄O₂₈) and similar UV, IR, ¹H, and ¹³C NMR spectra as compound **1**. A combined analysis of ¹H, ¹³C NMR (SI Table S2), ¹H–¹H COSY, HSQC, and HMBC data of **2** indicated that it possessed the same molecular structure as **1**. The glucopyranosyl core also had the same conformation and D-configuration. The experimental ECD spectrum exhibited a negative Cotton effect at 233 nm indicative of an HHDP moiety with an (*R*)-configured biphenyl bond. The anomeric galloyl group was assigned an α -equatorial orientation based on the ca. 1.0 Hz coupling constant of the anomeric proton. The 5,6-dihydroxy-3a-(2-methoxy-2-oxoethyl)-1-(methoxycarbonyl)-3-oxo-1,3,3a,8b-tetrahydrofuro[3,4-*b*]benzofuran-8-carboxylic acid moiety also showed the same relative configuration as **1**. Thus, **1** and **2** only differ as far as the absolute configuration of this rare structural moiety is concerned. Compound **2** is, thus, a (C-1'', C-3a'', C-8b'') quasi-enantiomer of compound **1** and accordingly named punicatannin B.

Scheme 1. Postulated Biosynthesis of the 4-Acyl Group of Punicatannins A (**1**) and B (**2**)



A possible biosynthetic pathway to the 5,6-dihydroxy-3a-(2-methoxy-2-oxoethyl)-1-(methoxycarbonyl)-3-oxo-1,3,3a,8b-tetrahydrofuro[3,4-*b*]benzofuran-8-carboxylic acid moiety of compounds **1** and **2** is postulated in Scheme 1. Its biosynthetic precursor is proposed to be an HHDP group, similar to that in, e.g., compound **1**, which is converted into **A** by oxidation and keto–enol tautomerism. Intermediate **A** would be cleaved into ketene **B** by hydrolysis.¹² This may be hydrolytically transformed into the enantiomeric intermediates **C1** and **C2** that may undergo a 1,4-Michael addition reaction to yield intermediates **D1** and **D2**. These may be converted into **E1** and **E2** by

(8) Jiang, Z. H.; Wen, X. Y.; Tanaka, T.; Wu, S. Y.; Liu, Z.; Iwata, H.; Hirose, Y.; Wu, S.; Kouno, I. *J. Nat. Prod.* **2008**, *71*, 719–723.

(9) Farina, F.; Maestro, M. C.; Mattin, M. V.; Soria, M. L. *Tetrahedron* **1987**, *43*, 4007–4014.

(10) (a) Okuda, T.; Yoshida, T.; Hatano, T.; Ito, H. In *Chemistry and Biology of Ellagitannins, An Underestimated Class of Bioactive Plant Polyphenols*; Quideau, S., Ed.; World Scientific Publishing: Singapore, 2009; Vol. 1, pp 1–54. (b) Ding, Y.; Li, X. C.; Ferreira, D. *J. Nat. Prod.* **2009**, *72*, 327–335.

(11) Diedrich, C.; Grimme, S. *J. Phys. Chem. A* **2003**, *107*, 2524–2539.

(12) Mayer, W.; Hoffmann, E. H.; Lösch, N.; Wolf, H.; Wolter, B.; Schilling, G. *Liebigs Ann. Chem.* **1984**, *5*, 929–938.

Table 1. α -Glucosidase Inhibitory Activities of Compounds 1–3

no.	IC ₅₀ (μ M) ^a
1	7.15 $\pm$ 0.27
2	12.39 $\pm$ 0.05
3	104.89 $\pm$ 7.73
acarbose ^b	208.60 $\pm$ 3.39

^aIC₅₀ values are shown as mean $\pm$ SD from three independent experiments. ^bPositive control.

trans-esterification. Methyl esterification of **E1** and **E2** would lead to the structural moieties **F1** and **F2** of compounds **1** and **2**.

Given the ethnomedicinal uses of pomegranate flowers for treating diabetes, we evaluated the isolated compounds for biological activities. First, compounds **1–3** and the clinical drug, acarbose, were tested for α -glucosidase inhibitory activities. This enzyme is a carbohydrate hydrolase that breaks down starch and α -linked di-, oligo-, and polysaccharides to glucose leading to elevated blood glucose levels. Thus, inhibition of the enzyme decreases the rate of digestion of carbohydrates and less glucose is absorbed. As shown in Table 1, compounds **1–3** showed potent and concentration-dependent α -glucosidase inhibitory activities (SI Figure S3). Remarkably, puniceatannin A (**1**) was ca. 30 times more potent than acarbose (IC₅₀ = 7.15 vs 208.60 μ M, respectively).

Finally, based on compound availability, we selected compound **1** to evaluate its ability to inhibit lipogenic gene expression. It is known that obesity associated with insulin resistance and type 2 diabetes increases the risk of hepatic lipid accumulation. Lipid accumulation is also associated with increased *de novo* lipogenesis, and high glucose concentrations have been shown to induce lipid accumulation in HepG2 cells.¹³ This is known to occur through the transcription factor, Srebp1c, which along with the lipogenic enzymes Acc-1, Fas, and Scd1 regulate lipogenesis. Furthermore, loss of Scd1 protects against obesity induced by decreased *de novo* lipogenesis,¹⁴ and forced CD36 expression increases fatty acid uptake in HepG2 cells.¹⁵ Here we showed that compound **1** prevents Acc1, Scd1, and CD36 induction in HepG2 cells exposed to high

glucose (Figure 2). This suggests that compound **1** may repress the induction of lipogenic gene expression by high glucose exposure and could aid in the prevention of high glucose induced *de novo* lipogenesis and subsequently inhibit lipid accumulation.




Figure 2. Compound **1** prevents elevation of Acc1, Scd1, and CD36 gene expression in HepG2 cells exposed to high glucose. (*) $P < 0.05$, group of high glucose compared to group of low glucose treatments in absence of compound **1**. (#) $P < 0.05$, group of high glucose compared to group of low glucose treatments in presence of compound **1** (50 μ M). (\$) $P < 0.05$, group of high glucose compared to group of low glucose treatments in presence of compound **1** (100 μ M).

In summary, this work describes the discovery of two new ellagitannins containing the rare 3-oxo-1,3,3a,8b-tetrahydrofuro[3,4-*b*]benzofuran moiety. The abilities of these compounds to inhibit α -glucosidase enzyme activity and lipogenic gene induction could account for the reported antidiabetic properties of pomegranate flowers. Furthermore, the structure identified in these natural compounds could serve as a scaffold for the synthesis of structural analogs with enhanced antidiabetic activities.

Supporting Information Available. General experimental procedures, plant materials, extraction and isolation of compounds, hydrolysis of compound **1** and sugar analysis, α -glucosidase and lipogenic gene expression inhibition assays, calculated ECD analysis, tabulated NMR data and full NMR, MS spectra for compounds **1** and **2**, and ¹H NMR and MS spectra for compound **1a**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The authors declare no competing financial interest.

(13) Zang, M.; Zuccollo, A.; Hou, X.; Nagata, D.; Walsh, K.; Herscovitz, H.; Brecher, P.; Ruderman, N. B.; Cohen, R. A. *J. Biol. Chem.* **2004**, 279, 47898–47905.

(14) Miyazaki, M.; Kim, Y. C.; Ntambi, J. M. *J. Lipid Res.* **2001**, 42, 1018–1024.

(15) Krammer, J.; Digel, M.; Ehehalt, F.; Stremmel, W.; Fullekrug, J.; Ehehalt, R. *Int. J. Med. Sci.* **2011**, 8, 599–614.