

The Intermolecular Pauson-Khand Reaction of Sugar-Derived Azaenynes: A Case of Total Diastereoselectivity


Pilar Areces and Esther Carrasco

Departamento de Química Orgánica, Universidad de Extremadura, Facultad de Ciencias, Badajoz, Spain

Joaquín Plumet

Departamento de Química Orgánica, Universidad Complutense, Facultad de Química, Madrid, Spain

Sugar azaenynes **1a** and **1b**, easily available from D-glucal and D-galactal, respectively, react with norbornene and norbornadiene under Pauson-Khand conditions to give the intermolecular adducts as major or exclusive product. In all cases only one diastereomer was obtained in convenient yield. This reaction constitutes an unprecedented case of totally diastereoselective intermolecular Pauson-Khand reaction using a sugar moiety as chiral auxiliary.



Keywords Intermolecular Pauson-Khand reaction, Cyclopentenones, Carbohydrates


Received November 10, 2005; accepted January 27, 2006.

Address correspondence to Pilar Areces, Departamento de Química Orgánica, Universidad de Extremadura, Facultad de Ciencias, 06071 Badajoz, Spain. E-mail: pareces@unex.es; and to Joaquín Plumet, Departamento de Química Orgánica, Universidad Complutense, Facultad de Química, 28040 Madrid, Spain. E-mail: plumety@quim.ucm.es


The Pauson-Khand (P-K) reaction, a cobalt carbonylative co-cyclization of an alkene and an alkyne, is the most popular organometallic-mediated reaction for the synthesis of cyclopentenones.^[1] In the last few years, many efforts were devoted to the development of the enantioselective version of this reaction and, for the intermolecular process,^[2] several methodologies in which a chiral inductor is covalently bonded to one of both partners have been developed.^[3] Regarding the use of carbohydrates as a source of chirality, these compounds have been considered as chiral substrates in the intramolecular P-K reactions.^[4] Recently we have reported the intramolecular P-K reaction of sugar azaenines **1a** and **1b**, derived from D-glucal and D-galactal, respectively, obtaining the expected cycloadducts **2** and **3** with poor^[5] to moderate^[6] diastereoselectivity (Sch. 1).

To the best of our knowledge, sugar derivatives have never been used as chiral auxiliaries in the intermolecular Pauson-Khand reactions. In this context we thought that compounds **1a** and **1b** may be interesting alkyne partners for the intermolecular Pauson-Khand reactions, assuming that the intermolecular reaction should be conveniently faster than the intramolecular process. Taking into account that the concurrence between intermolecular and intramolecular Pauson-Khand reactions have never been considered in substrates capable to undergo the intramolecular process, we have selected the reactive bicyclic alkenes norbornene **4** and norbornadiene **5** as olefinic counterparts. Norbornene and norbornadiene are very often used in the intermolecular Pauson-Khand reaction because of their high reactivity. In particular, these compounds are excellent reagents for testing the asymmetric version of the reaction. See literature quoted in reference 2, p. 3028. For the use of oxazabicyclic derivatives in Pauson-Khand reaction, see ref.^[7]

Compounds **1a** and **1b** reacted with norbornene and norbornadiene to give the intermolecular *exo* cycloadducts **6** and **7**, respectively (Table 1), as major (entry 1) or only (entries 2–4) reaction products. On the other hand, ¹H and ¹³C NMR spectra of the reaction mixtures showed that each individual



Scheme 1.

Table 1: The intermolecular Pauson-Khand reaction of sugar azaenines **1a** and **1b** with norbornene (**4**) and norbornadiene (**5**).

Entry	Starting materials	Cycloadduct	Isolated yield
1	1a and 4	6a	67% ^a
2	1a and 5	7a	62% ^b
3	1b and 4	6b	80% ^b
4	1b and 5	7b	72% ^b

^aTwenty percent of a mixture of intramolecular cycloadducts **2** and **3** was isolated.^bOnly traces of intramolecular cycloadducts **2** and **3** were observed in the crude reaction mixture.

cyclopentenone was diastereomerically pure with only one set of peaks present in each case. Flash chromatography of the crude reaction mixture through a small column (silica gel, 3 cm high; hexane-ethyl acetate 3:1 → ethyl acetate) followed by HPLC analysis and purification (Zorbax Rx-SIL column, isocratic normal phase separation, hexane-ethyl acetate 50:50) allowed us to isolate compounds **6** and **7** in moderate to good isolated yields and diastereomeric purity >95%. General procedure for compounds **6a**, **6b**, **7a**, and **7b**. A 0.16 M solution of azaenine precursor (**1a** or **1b**) in freshly distilled CH_2Cl_2 was treated with $\text{Co}_2(\text{CO})_8$ (1.1 equiv.) under a CO atmosphere. After 45 min of stirring, the corresponding alkene (**4** or **5**) was added, and the mixture was stirred 10 min more. Then, NMO (6 equiv.) was added portionwise over a 15-min period, and the reaction mixture was cooled to 0°C before each addition and allowed to reach rt before a new one. The mixture was stirred for 45 min at rt; diluted with CH_2Cl_2 ; washed successively with 2 N HCl, a saturated solution of NaHCO_3 , and water; dried (MgSO_4); and concentrated in vacuo. The residue was first eluted through a short column (3 cm long) of SiO_2 (hexane/ethyl acetate 3:1 → ethyl acetate) to remove metal impurities, and the elution was concentrated; flash chromatography (hexane/ethyl acetate 3:1 → 1:1) or HPLC analysis (semipreparative system, Zorbax Rx-SIL column, isocratic normal phase separation, hexane/ethyl acetate 50:50) of the residue gave the cyclopentenone (**6** or **7**).

Compound 6a: ^1H NMR (400 MHz, CDCl_3) δ 7.67 (d, 2H, $J_{3''',2'''} = 8.2$ Hz, H-3''', H-5'''), 7.30 (d, 2H, $J_{5'',6''} = 8.2$ Hz, H-2'', H-6''), 7.35 (s, H-3), 5.52–5.56 (m, 2H, H-2', H-3'), 5.38 (m, H-4'), 5.12 (m, H-5), 4.20–4.00 (m, 2H, H-6'a, H-6'b), 3.95–3.60 (m, 4H, 2H-1', 2H-1''), 2.59 (br s, H-3a), 2.43 (s, 3H, $\text{CH}_3\text{-Ar}$), 2.34 (m, H-7), 2.14 (m, 2H, H-7a, H-4), 2.06 (s, 3H, OAc), 2.05 (s, 3H, OAc), 2.04 (s, 3H, OAc), 1.70–1.50 (m, 2H, H-5), 1.35–1.20 (m, 2H, H-6), 0.97–0.80 (m, 2H, H-8); ^{13}C NMR (100 MHz, CDCl_3) δ 209.5 (C-1), 170.4, 169.9, 169.3 (3O-CO- CH_3), 162.6 (C-3), 144.4 (C-2), 143.5 (C-1'''), 136.5 (C-4'''), 130.0 (C-2'), 129.6 (C-3''' and C-5'''), 128.0 (C-3'), 127.1 (C-2'', C-6''), 71.2 (C-4', C-5'), 61.5 (C-6'), 54.0 (C-7a), 49.8, 49.7 (C-1'), 48.3 (C-3a), 41.9 (C-1''), 38.8 (C-7), 37.7 (C-4), 31.1 (C-8), 28.9 (C-5), 28.2 (C-6), 21.4 (C-7'''), 20.7, 20.5 (3O-CO- CH_3). HRMS calcd for $\text{C}_{30}\text{H}_{37}\text{NO}_9\text{S} + \text{H}$: 588.2267. Found: 588.2267.

Compound 6b: ^1H NMR (400 MHz, CDCl_3) δ 7.66 (d, 2H, $J_{3''',2'''} = 8.2$ Hz, H-3''', H-5'''), 7.35 (m, H-3), 7.30 (d, 2H, $J_{5'',6''} = 8.2$ Hz, H-2'', H-6''), 5.59–5.53 (m, 2H, H-2', H-3'), 5.39 (m, H-4'), 5.13 (m, H-5'), 4.27 (dd, $J_{5',6a'} = 4.0$ Hz, $J_{6b',6a'} = 12.0$ Hz, H-6'a), 3.98–3.93 (m, H-6'b), 3.90–3.73 (m, 4H, 2H-1', 2H-1''), 2.59 (br s, H-3a), 2.43 (s, 3H, $\text{CH}_3\text{-Ar}$), 2.35 (br s, H-7), 2.17–2.14 (m, 2H, H-7a, H-4), 2.07 (s, 3H, OAc), 2.05 (s, 3H, OAc), 2.04 (s, 3H, OAc), 1.67–1.57 (m, 2H, H-5), 1.31–1.25 (m, 2H, H-6), 0.93–0.90 (m, 2H, H-8); ^{13}C NMR (100 MHz, CDCl_3) δ 209.7 (C-1), 170.4, 170.0, 169.5 (3O-CO- CH_3), 162.8 (C-3), 144.5 (C-2), 143.5 (C-1'''), 136.6 (C-4'''), 130.0 (C-3''', C-5'''), 128.2 (C-3'), 127.1 (C-2'', C-6''), 71.1, 71.0 (C-4', C-5'), 61.9 (C-6'), 54.1 (C-7a), 49.7 (C-1'), 48.4 (C-3a), 41.8 (C-1''), 38.9 (C-7), 37.8 (C-4), 31.2 (C-8), 29.0, 28.3 (C-5, C-6), 21.4 (C-7'''), 20.8, 20.7, 20.6 (3O-CO- CH_3); HRMS calcd for $\text{C}_{30}\text{H}_{37}\text{NO}_9\text{S} + \text{H}$: 588.2267. Found 588.2269.

Compound 7a: ^1H NMR (400 MHz, CDCl_3) δ 7.67 (d, 2H, $J_{3''',2'''} = 8.2$ Hz, H-3''', H-5'''), 7.43 (br s, H-3), 7.31 (d, 2H, $J_{5'',6''} = 8.2$ Hz, H-2'', H-6''), 6.29 (dd, $J_{4,5} = 3.0$ Hz, $J_{5,6} = 5.5$ Hz, H-5), 6.19 (dd, $J_{6,7} = 2.9$ Hz, $J_{6,5} = 5.5$ Hz, H-6), 5.64–5.52 (m, 2H, H-2', H-3'), 5.38 (m, H-4'), 5.12 (m, H-5'), 4.18–4.06 (m, 2H, H-6'a, H-6'b), 3.93–3.73 (m, 4H, 2 H-1', 2 H-1''), 2.90 (br s, H-7), 2.73 (br s, H-3a), 2.68 (br s, H-4), 2.43 (s, 3H, $\text{CH}_3\text{-Ar}$), 2.26 (d, $J_{7a,7} = 5.0$ Hz, H-7a), 2.06 (s, 3H, OAc), 2.04 (s, 3H, OAc), 2.03 (s, 3H, OAc), 1.37 (d, $J_{8a,8b} = 9.1$ Hz, H-8b), 1.12 (d, $J_{8a,8b} = 9.1$ Hz, H-8a); ^{13}C NMR (100 MHz, CDCl_3) δ 208.4 (C-1), 170.0, 169.4 (3O-CO- CH_3), 162.7 (C-3), 145.7 (C-2), 143.6 (C-1'''), 138.5 (C-5), 137.0 (C-6), 136.6 (C-4'''), 130.2 (C-2'), 129.8, (C-3''', C-5'''), 128.2 (C-3'), 127.2 (C-2'', C-6''), 71.4, (C-4', C-5'), 61.6 (C-6'), 52.9 (C-7a), 50.1 (C-1'), 48.0 (C-3a), 43.6 (C-7), 42.8 (C-4), 42.0 (C-1''), 41.3 (C-8), 21.5 (C-7'''), 20.8, 20.7 (3O-CO- CH_3); HRMS calcd for $\text{C}_{30}\text{H}_{35}\text{NO}_9\text{S} + \text{H}$: 586.2110. Found: 586.2111.

Compound 7b: ^1H NMR (400 MHz, CDCl_3) δ 7.67 (d, 2H, $J_{2''',3'''} = J_{5''',6'''} = 8.4$ Hz, H-3''', H-5'''), 7.42 (br s, H-3), 7.31 (d, 2H, $J_{5'',6''} = 8.4$ Hz,

H-2''', H-6'''), 6.24 (dd, $J_{4,5} = 3.2$ Hz, $J_{5,6} = 5.2$ Hz, H-5), 6.19 (dd, $J_{6,7} = 2.8$ Hz, $J_{6,5} = 5.2$ Hz, H-6), 5.57 (m, 2H, H-2', H-3'), 5.39 (m, H-4'), 5.13 (m, H-5'), 4.27 (dd, $J_{5',6'} = 4.4$ Hz, $J_{6'a,6'b} = 12.0$ Hz, H-6'a), 3.99–3.73 (m, H-6'b), 3.88–3.72 (m, 4H, 2H-1', 2H-1''), 2.87 (br s, H-7), 2.73 (br s, H-3a), 2.68 (br s, H-4), 2.43 (s, 3H, CH₃-Ar), 2.26 (m, H-7a), 2.07 (s, 3H, OAc), 2.05 (s, 3H, OAc), 2.04 (s, 3H, OAc), 1.37 (d, $J_{8a,8b} = 9.6$ Hz, H-8a), 1.12 (d, $J_{8a,8b} = 9.6$ Hz, H-8a); ¹³C NMR (100 MHz, CDCl₃) δ 209.0 (C-1), 171.0, 170.6, 170.1 (3O-CO-CH₃), 163.4 (C-3), 146.2 (C-2), 142.2 (C-1'''), 139.1 (C-5), 137.6 (C-6), 137.2 (C-4'''), 130.6 (C-2'), 130.4, 130.5 (C-3''', C-5'''), 128.9 (C-3'), 127.8 (C-2'', C-6'''), 71.7, 71.6 (C-4', C-5'), 62.6 (C-6'), 53.5 (C-7a), 50.6 (C-1'), 48.6 (C-3a), 44.2 (C-7), 43.4 (C-4), 42.6 (C-1''), 41.9 (C-8), 22.1 (C-7'''), 21.4, 21.3 (3O-CO-CH₃); HRMS calcd for C₃₀H₃₅NO₉S + H: 586.2110. Found: 586.2113.

The assignment of the relative configuration at the new formed stereocenters in diastereomerically pure compounds **6** and **7** is being carried out and will be reported in due course. On the other hand, the *exo* stereochemistry of the cycloaddition was deduced from the lack of coupling between the *endo* hydrogens and the bridgehead protons in the norbornane framework.^[8]

In summary, in this paper we have showed that the concurrence between inter- and intramolecular Pauson-Khand reactions in substrates such as **1a** and **1b** capable to give both types of reactions was clearly shifted toward the intermolecular process using reactive bicyclic alkenes such as **4** or **5**. Interestingly, only one diastereomer was obtained in these reactions, showing an unprecedented diastereoselective control, which must be exerted by the sugar side chain. Synthetic applications of this process are now under research in our laboratory.

ACKNOWLEDGEMENTS

This work was supported by Ministerio de Educación y Ciencia, Spain (Grant BQU2003-04967-C02-01), and Gobierno de Extremadura, Spain (Grant 2PR02A016). One of us (E.C.) acknowledges a fellowship from the MEC, Spain.

REFERENCES

- [1] (a) Boñaga, L.R.V.; Kraft, M.E. When the Pauson-Khand and Pauson-Khand types reactions go awry: a plethora of unexpected results. *Tetrahedron* **2004**, *60*, 9795–9833; (b) Blanco-Urgoiti, J.; Anorbe, L.; Pérez-Serrano, L.; Domínguez, G.; Pérez-Castell, J. The Pauson-Khand reaction, a powerful synthetic tool for the synthesis of complex molecules. *Chem. Soc. Rev.* **2004**, *33*, 32–42.
- [2] Gibson, S.E.; Mainolfi, M. The intermolecular Pauson-Khand reaction. *Angew. Chem. Int. Ed.* **2005**, *44*, 3022–3037.
- [3] (a) Ingate, S.T.; Marco-Contelles, J.L. The asymmetric Pauson-Khand reaction. A review. *Org. Prep. Proc. Int.* **1998**, *30*, 121–143; (b) Rivero, M.R.; Alonso, I.; Carretero, J.C. Vinyl sulfoxides as stereochemical controllers in intermolecular Pauson-Khand reactions: applications to the enantioselective synthesis of natural

- cyclopentanoids. *Chem. Eur. J.* **2004**, *10*, 5443–5459; (c) Rivero, M.R.; Alonso, I.; Carretero, J.C. Pauson-Khand reactions of alkenyl sulfones and alkenyl sulfoxides: Applications in asymmetric synthesis. *Synlett* **2005**, 26–41.
- [4] (a) Marco-Contelles, J.L.; de Opazo, E.J. Strategies for the synthesis of enantiomerically pure medium-sized carbocycles from carbohydrates. *Carbohydr. Chem.* **2002**, *21*, 201–218; (b) Marco-Contelles, J.L.; Ruiz Caro, J. Synthesis and transformations of new annulated pyranosides using the Pauson-Khand reaction. *Carbohydr. Res.* **2001**, *335*, 71–90; (c) Marco-Contelles, J.L.; Ruiz Caro, J. Iridoids from carbohydrates via Pauson-Khand reaction: Synthesis of advanced highly oxygenated cyclopentane-annulated pyranosides from D-glucal derivatives. *J. Chem.* **1999**, *64*, 8302–8310; (d) Hotha, S.; Maurya, S.K.; Gurjar, M.K. Stereoselective synthesis of spiroannulated cyclopentenones by the Pauson-Khand reaction on carbohydrates derived enynes. *Tetrahedron Lett.* **2005**, *46*, 5329–5332.
- [5] Areces, P.; Durán, M.A.; Plumet, J.; Hursthouse, M.B.; Light, K.E. Synthesis of enantiopure 3-azabicyclo[3.3.0]octen-7-one derivatives via intramolecular Pauson-Khand cycloaddition reaction using tri-*o*-acetyl-D-glucal as starting material. *J. Org. Chem.* **2002**, *67*, 3506–3507.
- [6] Areces, P.; Carrasco, E.; Plumet, J. From D-galactose to enantiopure 3-azabicyclo[3.3.0]octen-7-one derivatives via Pauson-Khand reaction. *Arkivoc* **2005** (ix) 165–174.
- [7] (a) Arjona, O.; Csáky, A.G.; Murcia, C.; Plumet, J. J. Control of the regioselectivity in the Pauson-Khand reaction of 7-oxanorbornene derivatives. *Org. Chem.* **1999**, *64*, 7338–7341; (b) Arjona, O.; Csáky, A.G.; Medel, R.; Plumet, J. First intermolecular Pauson-Khand reaction of 7-azanorbomenes. Control of the regioselectivity by the effect of the substituents attached to the olefinic partner. *Tetrahedron Lett.* **2001**, *42*, 3085–3088; (c) Ahmar, M.; Cazes, B. Pauson-Khand reaction of 7-oxabicyclo [2.2.1] systems. *Tetrahedron Lett.* **2003**, *44*, 5403–5406.
- [8] (a) Anet, F.A.L. Coupling constants in rigid five-membered rings: N.M.R. spectra of the camphene-2,3-diols. *Canad. J. Chem.* **1961**, *39*, 789; (b) Huisgen, R.; Möbius, L.; Müller, G.; Stangl, H.; Szemies, G.; Vernon, J.M. 1,3-Dipolare cycloadditionen, XXII. Zur anlagerung organischer azide an winkelgespannte doppelbindungen. *Chem. Ber.* **1965**, *98*, 3992.

Copyright of Journal of Carbohydrate Chemistry is the property of Taylor & Francis Ltd and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.