

tion products of benzylidenedecalone,3 were undoubtedly the stereoisomers II and III epimeric about the carbon holding the methyl group. One of these isomers  $(\beta^2)$  was converted to estrone by reactions already described<sup>1</sup> and is therefore represented by formula III which corresponds to the most probable configuration for the natural product. The  $\beta^1$  epimer (formula II) yielded estrone "g" which differs from natural estrone only in the configuration at C<sub>13</sub> and therefore represents unequivocally that stereoisomeric form which was postulated by Butenandt and co-workers<sup>4</sup> for lumiestrone. We have now compared the infrared spectra of dl-estrone "g" methyl ether and lumiestrone methyl ether,  ${}^{4}$  m.p. 130–130.6°,  $[\alpha]^{32}$ D –27° in chloroform, and have found them to be identical.<sup>5</sup> These results clearly demonstrate that our estrone "g" instead of Anner and Miescher's estrone "f" is lumiestrone. Moreover, the identity of lumiestrone with synthetic material known to be epimeric with estrone at C13, establishes the configuration postulated by Butenandt for this stereoisomer, and proves that the photoisomerization of estrone involves inversion only at C13.

(3) W. S. Johnson, THIS JOURNAL, 65, 1317 (1943).

(4) A. Butenandt, A. Wolff and P. Karlson, Ber., 74, 1308 (1941);
 A. Butenandt, et al., ibid., 75, 1931 (1942); 77, 392 (1944).

(5) We are indebted to Dr. R. N. Jones of the National Research Council, Canada, for determining the spectra.

LABORATORY OF ORGANIC CHEMISTRY

UNIVERSITY OF WISCONSIN MADISON, WISCONSIN

RECEIVED JULY 23, 1951

## Thiation with Phosphorus Pentasulfide in Pyridine Solution

# By Erwin Klingsberg<sup>1</sup> and Domenick Papa

The conversion of pyridones to thiopyridones by means of phosphorus pentasulfide has hitherto been effected by fusion<sup>2</sup> or by heating in a liquid medium, such as benzene<sup>3</sup> or toluene,<sup>4</sup> with limited solvent power for one or both reactants. These reactions, however, have apparently been limited

(1) American Cyanamid Company, Calco Chemical Division, Bound Brook, New Jersey.

(8) Arndt and Kalischek, Ber., 68, 587 (1930).

(4) Arndt, ibid., 65, 92 (1932).

to comparatively low-melting pyridones. Rath<sup>5</sup> treated 5-iodo-2-pyridone (I) with phosphorus pentasulfide and obtained not the corresponding iodothiopyridones, but 2-thiopyridone, the iodine having been lost in the conversion.

In the present investigation, initial experiments with 3,5-diiodo-2-pyridone (II) were unsuccessful, the diiodopyridone being recovered unchanged after prolonged reflux in benzene with phosphorus pentasulfide. Fusion of the two reactants at various temperatures resulted only in intractable reaction mixtures.

It appeared necessary to find a solvent which would permit the conversion reaction to proceed in solution. Pyridine proved excellent for the purpose, having ample solvent power for the reactants, a suitable boiling temperature; and, by its miscibility in water, providing a simple method of isolation. On heating a mixture of 3,5-diiodo-2pyridone, phosphorus pentasulfide and pyridine for 1.5 hours and pouring the reaction solution into approximately twice its volume of water, an excellent yield of 3,5-diiodo-2-thiopyridone (III) separated in substantially pure form. The phosphorus-containing by-products remained in the aqueous pyridine. Similarly, 3,5-diiodo-4-pyridone (IV) was converted to the thiopyridone (V) in quantitative yield. This method facilitates the



conversion of the high-melting pyridones to the corresponding mercaptans in one step, in contrast to the former two-step procedure of converting the pyridone to the active chlorine derivative and subsequent metathesis with hydrosulfide.<sup>6</sup>

The great versatility of pyridine as an organic solvent suggests its use in a variety of thiations with phosphorus pentasulfide, and in a single exploratory experiment, benzanilide was converted to thiobenzanilide in 73% yield.

#### Experimental

3,5-Diiodo-2-thiopyridone (III).—A solution of 70 g. (0.202 mole) of 3,5-diiodo-2-pyridone and 50 g. (0.225 mole) of phosphorus pentasulfide in 300 ml. of pyridine was refluxed for 1.5 hours and poured into 500 ml. of hot water. Cold water tends to give a gummy product, probably due to incomplete solution of the phosphorus by-products. The slurry was cooled and filtered. On recrystallization from dilute pyridine, there was obtained 65 g. of product (89% of theory), m.p. 200-203°. A second recrystallization gave 52 g. (70%) of product, m.p. 206-206.5°.

Anal. Calcd. for C<sub>5</sub>H<sub>8</sub>NI<sub>2</sub>S: C, 16.54; H, 0.83; N, 3.86. Found: C, 16.08; H, 1.09; N, 3.85.

Dohrn and Diedrich<sup>6</sup> prepared this compound from 2chloro-3,5-diiodopyridine and reported m.p. 181°, but no analysis.

**3,5-Diiodo-4-thiopyridone** (V).---3,5-Diiodo-4-pyridone (140 g., 0.404 mole) and phosphorus pentasulfide (110 g., 0.495 mole) were refluxed in 550 ml. of pyridine for 2 to 2.5 hours and the solution then poured into 800 ml. of hot water. On cooling, filtering, washing and drying, there

<sup>(2)</sup> King and Ware, J. Chem. Soc., 873 (1939).

<sup>(5)</sup> Rath, Ann., 487, 105 (1981).

<sup>(6)</sup> Dohrn and Diedrich, ibid., 494, 284 (1932).

was obtained a quantitative yield of 3,5-diiodo-4-thiopyrisolving in sodium carbonate or ammonia solution, filtering and reprecipitating with acid. The purified product melts at  $204-205^{\circ}$ .<sup>6</sup>

Thiobenzanilide.—Benzanilide (41 g., 0.071 mole) and phosphorus pentasulfide (17 g., 0.076 mole) were refluxed 40 minutes in 70 ml. of pyridine and poured into water. The The minutes in 70 mil. of pyridine and poured into water. The resulting gum turned crystalline when made weakly alkaline with potassium hydroxide. The product can be crystallized from dilute methanol or purified by dissolving in alkali and precipitating with hydrochloric acid. There was obtained 11 g. (73%), m.p. 96–96.5°.<sup>7</sup> Methylation with dimethyl sulfate and alkali gave the S-methyl ether, m.p. 65.5–66°.<sup>8</sup>

(7) Bernthsen, Ber., 11, 503 (1878).

(8) May, J. Chem. Soc., 103, 2272 (1913).

CHEMICAL RESEARCH DEPARTMENT

SCHERING CORPORATION

BLOOMFIELD, NEW JERSEY RECEIVED MAY 17, 1951

## Preparation of Some Mixed Dialkyl Phosphites

## By GENNADY M. KOSOLAPOFF

There is a very considerable body of literature concerning the symmetric dialkyl phosphites, but not a single example of a mixed dialkyl phosphite, containing two unlike radicals, can be found.

In order to secure some information about such compounds, their preparation was examined. Although, theoretically, one may expect to obtain such compounds by the reaction of phosphorus trichloride with a mixture of alcohols, such a reaction yields complex mixtures which contain relatively small amounts of the mixed derivatives. Another possible method is exchange esterification with symmetric dialkyl phosphites. This method gives the desired mixed esters in moderate yields and can be made to proceed without difficulty at reflux conditions without added catalysts.

of MR values when AR for phosphorus is taken as 7.04. Trials of the latter value in calculations involving the dialkyl phosphites leave no doubt that this value cannot be applied at all. This confirms the non-trivalence of phosphorus in these compounds. The use of the 3.75 value in calculations with the compounds described in this paper, along with a number of previously prepared symmetric esters, reveals a need for a new AR value for phosphorus for compounds having the structures of the type found in dialkyl phosphites. The table below gives the results of calculations using the "phosphate" AR of 3.75, and the newly proposed value of 4.44, derived on empirical basis from the calculations with Kabachnik's value. The magnitude of this constant appears to indicate that the phosphorus atom in these esters is much closer to the tetra-(or penta)-valent state of the phosphates, than it is to the trivalent state found in other phosphites in which the trivalency is clearly demonstrable.

#### Experimental

Ethyl Butyl Phosphite.—A mixture of 27.6 g. (0.2 mole) of diethyl phosphite and 14.8 g. (0.2 mole) of *n*-but anol was slowly heated in a distillation apparatus up to bath temperature of 150°, when a brisk reaction started and ethanol began to distil. The bath was kept at 150–155° for 55 minutes, at which time 5.8 ml. of ethanol was collected. Distillation of the mixture under reduced pressure gave 8 g. of unchanged diethyl phosphite, 16 g. of crude ethyl butyl phosphite, b.p. 103-105° at 13 mm., and 3 g. of crude dibutyl phos-phite, b.p. 110-115° at 11 mm.; the residue showed signs of decomposition on attempts at further distillation. Redistillation of the middle cut gave pure ethyl butyl phos-phite (15.0 g., 45%), b.p. 99-100° at 13 mm.,  $n^{27}$ D 1.4139,  $d^{27}_0$  1.0120. Found: P, 18.6, 18.8. Calcd. for (EtO)-(BuO)POH: P, 18.7. Treatment with warm water, acidified with hydrochloric acid, rapidly yielded phosphorous acid which was detected by the calomel test. This showed that no isomerization to a phosphonate took place during the re-action, thus confirming the above formula.

I ADLE 1						
Ester	Ref.	R. <b>I</b> .	Density	MRobsd.	$AR = \frac{MR_{\text{cale}}}{3.75}$	ed. with: AR = 4.44
(EtO)₂POH	<b>2</b>	n <sup>20</sup> D 1.4080	$d^{20}$ , 1.0742	31.69	31.02	31.71
(PrO) <sub>2</sub> POH	2	1.4172	1.0184	41.03	40.25	40.94
(BuO) <sub>2</sub> POH	$^{2}$	1,4240	0.9888	50.10	<b>49.49</b>	50.18
(C <sub>6</sub> H <sub>13</sub> O) <sub>2</sub> POH	<b>2</b>	1.4325	0.9486	68.75	67.96	68.65
$(C_7H_{15}O)_2POH$	<b>2</b>	1.4382	0.9363	77.95	77.19	77.88
(EtO)(BuO)POH		$n^{27}$ d 1.4139	d <sup>27</sup> 1.0120	40.98	40.25	40.94
(EtO)(AmO)POH		1.4189	0. <b>993</b> 5	45.71	44.87	45.56

TADID

In the course of characterization of the products it was desirable to evaluate their molecular refractions. It was found that the values of MRthat are found experimentally are significantly higher than the calculated values, when the reasonable assumption is made that the esters possess the "keto" structure, i.e., (RO)<sub>2</sub>P(O)H. This formulation calls for a value of atomic refraction of phosphorus that is probably close to the value suggested by Kabachnik<sup>1</sup> for the phosphates, *i.e.*, for phosphorus in the higher state of oxidation: AR = 3.75. The use of a value in the vicinity of the above is reasonable since the dialkyl phosphites do not display the chemical properties of trivalent phosphorus. The truly trivalent derivatives, such as (RO), P and ROPX, give very good correlation

(1) Kabachnik, Isvestiya Akad. Nauk S.S.S.R., otdel. khim. nauk, 219 (1948); C. A., 48, 5786# (1948).

Ethyl n-Amyl Phosphite.---The procedure described above, using 27.6 g. of diethyl phosphite and 17.6 g. of n-amyl alcohol, gave 11.5 g. of unreacted diethyl phosphite and 12

alconol, gave 11.5 g. of unreacted diethyl phosphite and 12 g. of pure (after two distillations) ethyl n-amyl phosphite (33%), b.p. 109-110° at 13 mm., n<sup>27</sup>D 1.4189, d<sup>27</sup>0 0.9935.
Found: P, 17.2, 17.25. Calcd. for (EtO)(AmO)POH: P, 17.25. Behavior on hydrolysis was as described above. Reaction with n-Decanol.—When diethyl phosphite (27.6 g.) was heated as above with 31.6 g. of n-decanol, the reaction took place briskly at 150-155° bath temperature with distillation of ethanol. Attempted distillation of his further two gave only 9 g. of unreacted diethyl phosphite further. ture gave only 9 g. of unreacted diethyl phosphite, further distillation leading to considerable decomposition of the mixture even at 1 mm. Thus, the probably formed ethyl decyl phosphite could not be isolated.

# ROSS CHEMICAL LABORATORY

### ALABAMA POLYTECHNIC INSTITUTE Auburn, Ala.

**RECEIVED APRIL 16, 1951** 

(2) Arbuzov and Vinogradova, Doklady Akad. Nauk S.S.S.R., 55, 31 (1947); Investiya Akad. Nauk S.S.S.R., oldel, khim, nauk, 617 (1947); C. A., 48, 5844h (1948).