

THE STRUCTURES OF DIAZAQUINOMYCINS A AND B, NEW ANTIBIOTIC METABOLITES

SATOSHI ŌMURA*, AKIRA NAKAGAWA, HIROSHI AOYAMA, KIYOIZUMI HINOTOZAWA AND HIROSHI SANO

School of Pharmaceutical Sciences, Kitasato University and The Kitasato Institute
Minato-ku, Tokyo 108, Japan

Summary: Novel substituted 1,8-diazaanthraquinone structures of diazaquinomycins A and B were determined by the application of nmr spectroscopy.

Diazaquinomycins A ¹⁾ (1) and B ²⁾ (2), produced by Streptomyces sp. OM-704, exhibit antibacterial activities against Gram-positive bacteria and are antimetabolites of folate metabolism in Streptococcus faecium IF0 3181. This inhibitory activity is reversed by thymidine, leucovorin, and dihydrofolic acid. In this paper, we wish to report a novel 1,8-diazaanthraquinone structure (1) for diazaquinomycin A by the application of nmr spectroscopy.

The antibiotic (1), deep red needles, mp 291-295°C, EI Mass: $M^+ m/z$ 354.157 (Calcd. for $C_{20}H_{22}N_2O_4$, 354.157), showed a characteristic UV absorption, λ_{max}^{MeOH} nm (ϵ) 250 (11800, sh), 260 (13600, sh), 278 (20100, sh), 286 (21700), 309 (9760), 321 (8950), 367 (4130), and 490 (1150), and IR absorption, ν_{max}^{KBr} 1670, 1625 cm^{-1} (carbonyl) for an anthraquinone like structure.²⁾ The ^{13}C -nmr spectrum (100 MHz, $CDCl_3 + CD_3OD$, Table 1) showed eleven carbon signals, δ_C 182.9 and 173.9 (quinone carbonyls), δ_C 162.2 (two amide carbonyls), δ_C 151.2, 136.8, 135.0, and 117.3 (each two substituted aromatic carbons), δ_C 32.4 and 22.9 (each two methylene carbons), and δ_C 14.6 and 12.9 (each two methyl carbons), indicating the existence of a symmetric structural unit in the molecule. In the 1H -nmr spectrum (400 MHz, $CDCl_3 + CD_3OD$), the appearance of the signals at δ_H 2.27 (6H s), 1.01 (6H t, $J=7.2$ Hz), 1.57 (4H m) and 3.03 (4H t, $J=7.8$ Hz) and spin decoupling experiments indicated the existence of each two methyl and propyl groups linked to double bonds. Further nmr spectral investigation was carried out with methyl derivatives of 1 because of extremely low solubility of 1 in most nmr solvents.


Methylation of 1 with methyl iodide-silver oxide in N,N-dimethylformamide at room temperature afforded two products; N,N'-dimethyl derivative (3), red needles, mp 216-218°C, EI Mass: $M^+ m/z$ 382.190 (Calcd. for $C_{22}H_{26}N_2O_4$, 382.189), UV: λ_{max}^{MeOH} nm (ϵ) 250 (17900, sh), 260 (22700, sh), 284 (32500), 312 (20600), 323 (22500), 352 (7450), and 452 (1340), and N,O-dimethyl derivative (4), orange needles, mp 138-140°C, EI Mass: $M^+ m/z$ 382.187 (Calcd. for $C_{22}H_{26}N_2O_4$,

382.189), UV: $\lambda_{\text{max}}^{\text{MeOH}}$ nm (ϵ) 246 (13400, sh), 252 (15300, sh), 278 (33000), 309 (15900), 318 (15700), 340 (6300), and 425 (1530). The nmr spectrum of 3 showed an N-methyl signal (δ_{H} 3.74 and δ_{C} 34.4). On the other hand, the nmr [an N-methyl (δ_{H} 3.91 and δ_{C} 34.7) and a methoxy (δ_{H} 4.12 and δ_{C} 54.5)] and UV spectral data of 4 indicated the change from the symmetric structure to an asymmetric one by methylation. Methylation of 1 in chloroform gave an O,O'-dimethyl derivative (5), yellow needles, mp 136-138°C, EI Mass: $M^+ m/z$ 382.190 (Calcd. for $\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_4$, 382.189), UV: $\lambda_{\text{max}}^{\text{MeOH}}$ nm (ϵ) 245 (11900, sh), 254 (19100, sh), 274 (55400), 305 (17600), and 380 (1430), which showed a methoxy signal (δ_{H} 4.15 and δ_{C} 54.4) in its ^1H and ^{13}C nmr spectra. The methyl derivatives 3 and 5 possess a symmetrical structure, however, compound 4 does not, because of appearance of twenty two signals in the ^{13}C nmr spectrum of 4.

C-H Long range selective proton decoupling experiment (LSPD) is a good tool to determine bonding situation between carbons in substituted aromatic ring systems. In LSPD of 3 (Fig. 1), multiplets at δ_{C} 134.0 (C-3) and 162.1 (C-2) collapsed to a triplet (J_{CH} 4.6 Hz) and a quartet, respectively, and a multiplet at δ_{C} 147.2 (C-4) was sharpened upon irradiation of the signal at δ_{H} 2.23 assignable to C-methyl group. Upon irradiation of methylene proton (δ_{H} 2.89) of the propyl group bonded to the aromatic ring, multiplets at δ_{C} 147.2 (C-4) and 134.0 (C-3) collapsed to quartets (J_{CH} 3.8 and 5.3 Hz, respectively) and a triplet at δ_{C} 118.2 (C-4a) collapsed to a singlet. Irradiation of the N-methyl proton (δ_{H} 3.74) gave a singlet and a quartet (J_{CH} 3.8 Hz) from quartet (δ_{C} 139.7, C-9a) and multiplet (δ_{C} 162.1, C-2), respectively, and enhanced the intensity (60%) of a singlet assignable to a quinone carbonyl carbon (δ_{C} 178.6, C-9) resulted from $^{13}\text{C}\{-^1\text{H}\}$ NOE.³⁾ LSPD experiment in 5 also showed a similar change of signal patterns (Fig. 1); a multiplet (δ_{C} 125.9, C-3) to a triplet, a multiplet (δ_{C} 164.1, C-2) to a quartet and increase of the signal intensity of a multiplet (δ_{C} 152.8, C-4) upon irradiation of δ_{H} 2.28 (C-CH₃); multiplet (δ_{C} 152.8, C-4) to a quartet, triplet (δ_{C} 126.2, C-4a) to a singlet, multiplet (δ_{C} 125.9, C-3) to a quartet on irradiation of δ_{H} 3.09 (=C-CH₂CH₂CH₃); multiplet (δ_{C} 164.1, C-2) to a quartet on irradiation of δ_{H} 4.15 (OCH₃). These facts indicated the existence of a substituted six membered ring structure and thus symmetric structures of 3 and 5 were determined. Thus, substituted, 1,8-diazaanthraquinone structure (1) for diazaquinomycin A has been established from the structures of methylated derivatives 3, 4 and 5, in conjunction with its spectroscopic data.


Diazaquinomycin B (2), a minor component, was isolated from slow moving portion of silica

gel column chromatography as colorless needles, mp $>300^{\circ}\text{C}$, UV: $\lambda_{\text{max}}^{\text{MeOH}}$ nm (ϵ) 277 (20100), 310 (14800, sh), 325 (11750, sh), 356 (6230), and 373 (6050). The ^1H -nmr spectrum of 2 showed the existence of two methyl [δ_{H} 2.26 (6H s)] and two propyl groups [δ_{H} 1.13 (6H t, $J=7.6$ Hz, $=\text{C}-\text{CH}_2\text{CH}_2\text{CH}_3$), 1.72 (4H m, $=\text{C}-\text{CH}_2\text{CH}_2\text{CH}_3$), and 2.97 (4H t, $J=8.1$ Hz, $=\text{C}-\text{CH}_2\text{CH}_2\text{CH}_3$)]. Further, the air oxidation product of 2 was identical with 1 in every respect, indicating that diazaquinomycin B is a 9,10-dihydro derivative of 1.




1: R = H (diazaquinomycin A)


3: R = CH₃



2 (diazaquinomycin B)



4



5




Fig. 1. Long range C-H coupling in 3 and 5

Table 1. ^{13}C -nmr chemical shifts of diazaquinomycin A (1) and its derivatives 3, 4 and 5.

Carbon No.	<u>1</u>	<u>3</u>	<u>4</u>	<u>5</u>
2, 7	162.2	162.1	162.6 164.4	164.1
3, 6	135.0	134.0	134.0 124.6	125.9
4, 5	151.2	147.2	147.7 152.1	152.8
4a, 10a	117.3	118.2	120.8 126.3	162.2
8a, 9a	136.8	139.7	145.1 139.8	145.7
9	173.9	178.6	180.6	182.1
10	182.9	183.3	185.5	187.3
11, 18	12.9	13.6	13.8 11.8	11.8
12, 15	32.4	32.1	31.4 32.5	31.7
13, 16	22.9	23.0	23.0 23.2	23.1
14, 17	14.6	14.5	14.5 14.5	14.6
N-CH ₃	-	34.4	34.7	-
O-CH ₃	-	-	54.5	54.4

1 was measured in $\text{CDCl}_3 + \text{CD}_3\text{OD}$ and 3, 4, 5 in CDCl_3 .

References

- 1) S. Omura, Y. Iwai, K. Hinotozawa, H. Tanaka, Y. Takahashi and A. Nakagawa, *J. Antibiotics*, **35**, 1425 (1982).
- 2) A. I. Scott, "Ultraviolet spectra of natural products", Pergamon Press, London, p. 125 (1964).
- 3) J. Uzawa and S. Takeuchi, *Org. Mag. Res.*, **11**, 502 (1978).

Acknowledgement

The authors wish to thank Mrs. M. Yoshida, Tokyo Research Laboratory, Kyowa Hakko Kogyo Co., Ltd., for ^1H -nmr spectroscopy.

(Received in Japan 17 May 1983)