

286. The Chemistry of Simple Heterocyclic Systems. Part III Basic Centres of 4-Substituted Quinazoline Derivatives.


By J. S. MORLEY and J. C. E. SIMPSON.

The preparation from 4-hydroxyquinazoline of 4-phenoxy- and 4-amino-quinazoline is described.

Fusion of 4-phenoxyquinazoline with methyl toluene-*p*-sulphonate yields 1-methyl-4-quinazoline *via* an unstable quaternary salt, thus disclosing $N_{(1)}$ as the basic centre in the phenoxy-compound; however, similar treatment of 4-methoxyquinazoline attacks both $N_{(1)}$ and $N_{(3)}$ with formation of the quaternary salt (IV) or (IVa).

Attempted conversion of cinnoline methiodide into 1-methyl-4-cinnolone results in dealkylation and formation of cinnoline in 50% yield.

It has recently been shown (Morley and Simpson, *J.*, 1948, 360) that 6- and 7-nitro-4-acetamido- and -4-phenoxy-quinazoline are converted by quaternary-salt formation and subsequent fission into 6- and 7-nitro-1-methyl-4-quinazoline, and therefore that $N_{(1)}$ is the proton-accepting centre in these quinazolines. On the other hand, the centre of basicity, as deduced from quaternary-salt formation and degradation, is $N_{(3)}$ in quinazoline itself (Gabriel and Colman, *Ber.*, 1904, **37**, 3643). It was therefore desirable to ascertain whether $N_{(1)}$ or $N_{(3)}$ is the basic centre in other 4-substituted quinazolines, and we now record our results with 4-phenoxyquinazoline (I; R = Ph) and 4-methoxyquinazoline (I; R = Me).



4-Phenoxyquinazoline was readily prepared from 4-chloroquinazoline by reaction with phenol and potassium hydroxide. Treatment of the chloro-compound with phenol and ammonium carbonate likewise yielded (I; R = Ph) together with some 4-aminoquinazoline (II), which was also obtained from 4-chloroquinazoline and concentrated aqueous ammonia at room temperature (cf. Tomisek and Christensen, *J. Amer. Chem. Soc.*, 1945, **67**, 2112) and, almost quantitatively, from the phenoxy-compound and ammonium acetate at 190°. The properties of compounds previously designated as 4-aminoquinazoline are not in good agreement with those of our material; thus Dewar (*J.*, 1944, 619) states that 4-aminoquinazoline, m. p. 259—260°, is almost non-basic and fails to react with acid chlorides under a variety of conditions, and Tomisek and Christensen (*loc. cit.*) state that it melts at 245—260° (decomp.) and yields (apparently with considerable difficulty) an acetyl derivative, m. p. 172°. In our hands, 4-aminoquinazoline prepared by each of the three methods mentioned has m. p. 267° (decomp.), readily yields 4-acetamidoquinazoline, m. p. 174—175°, and has pronounced basic properties (see Part IV, following paper).

Fusion of 4-phenoxyquinazoline with methyl toluene-*p*-sulphonate proceeded with elimination of phenol, and subsequent treatment with water yielded 1-methyl-4-quinazoline (III), isolated as the *picrate*. The structure of this base was proved by alkaline fission to *o*-methylaminobenzamide (*N*-methylanthranilamide) and also by nitration to 6-nitro-1-methyl-4-quinazoline (Morley and Simpson, *loc. cit.*). As 4-hydroxyquinazoline yields 3-methyl-4-quinazoline under ordinary conditions of alkylation (Knape, *J. pr. chem.*, 1891, **43**, 209; Bogert and Geiger, *J. Amer. Chem. Soc.*, 1912, **34**, 524), these results indicate that 4-phenoxyquinazoline is converted by reaction at $N_{(1)}$ into an unstable quaternary salt, which then undergoes decomposition to 1-methyl-4-quinazoline.

Bogert and Geiger (*J. Amer. Chem. Soc.*, 1912, **34**, 683) found that 4-methoxyquinazoline, 4-hydroxyquinazoline, and 3-methyl-4-quinazoline reacted with methyl iodide to give the same iodide, C₁₀H₁₁ON₂I, which must therefore be represented by one or other of the structures (IV) or (IVa) (X = I); its formation is thus of no value for the purpose of deciding whether $N_{(1)}$ or $N_{(3)}$

is the basic centre in 4-methoxyquinazoline. In the hope that the use of a different reagent might produce the desired result of quaternisation without further alkylation, 4-methoxyquinazoline was fused with methyl toluene-*p*-sulphonate (1 mol.), but, after conversion into the iodide, the sole product was, again, the dialkylated salt (IV or IVa; X = I). Treatment of this salt with cold alkali yielded an oil, the properties of which suggested that it might be the *pseudo*-base (V); however, the oil gave a *picrate*, m. p. 189°, which differed from the quaternary *picrate* (IV or IVa; X = C₆H₂O₇N₃), m. p. 198°, prepared from the iodide.

No decision, therefore, can be made as to the position of the basic centre in 4-methoxyquinazoline. The results described in this and an earlier paper (J., 1948, 360), however, establish that the introduction of a phenoxy- or an acetamido-group into the quinazoline nucleus at C₄ produces a shift of the basic centre from N₍₃₎ to N₍₁₎. Particular interest thus attaches to the determination of the centre of basicity of the isomeric cinnoline (VI), inasmuch as N₍₁₎ has been shown to function in this capacity in 4-methyl- and 4-amino-cinnolines (Atkinson and Simpson, J., 1947, 808; Simpson, J., 1947, 1653). Preliminary attempts in this direction have been unsuccessful; oxidation of cinnoline methiodide with alkaline ferricyanide gave 50% of cinnoline, and no 1-methyl-4-cinnolone (Schofield and Simpson, J., 1945, 512) could be detected; and treatment of the methiodide with alkali in an atmosphere of nitrogen gave no definite product.

EXPERIMENTAL.

Melting points are uncorrected.

4-Phenoxyquinazoline.—(a) 4-Chloroquinazoline (3.3 g.) was added during 10 minutes to a solution of potassium hydroxide (1.3 g.) in warm phenol (40 g.; temperature > 50°). After 1 hour at 50° the melt was digested with cold 2N-sodium hydroxide and the suspension extracted with ether. The product (4.01 g.; 90%) from the washed and dried extract was crystallised from ligroin (b. p. 60–80°), yielding 4-*phenoxyquinazoline* as colourless plates, m. p. 78–79° (Found: C, 75.6; H, 4.6. C₁₄H₁₀ON₂ requires C, 75.65; H, 4.5%).

(b) 4-Chloroquinazoline (1 g.) was added with powdered ammonium carbonate (4 g.) to phenol (10 g.) at 90°. After $\frac{3}{4}$ hour the product was worked up as in (a). It was not wholly soluble in ether, and the insoluble fraction, after dissolution in 0.5N-hydrochloric acid and reprecipitation with ammonia, yielded 4-aminoquinazoline (0.17 g.), m. p. 265–267° (decomp.) alone and mixed with the material described below. The ether-soluble portion of the reaction-product gave 4-phenoxyquinazoline (0.7 g.).

4-Aminoquinazoline.—(a) 4-Phenoxyquinazoline (1 g.) was added to freshly-fused ammonium acetate (5 g.) at 160°; the temperature of the mixture was then raised to 190° and kept there for 2 minutes. When cold, the mass was dissolved in water (10 c.c.) and made alkaline with 2N-sodium hydroxide. Pure 4-*aminoquinazoline* (0.64 g.; 98%) separated in colourless needles, m. p. 267–268° (decomp.), unchanged by crystallisation from alcohol; it could not be obtained anhydrous even after prolonged drying at 80° in a vacuum (Found: C, 63.8; H, 4.85; N, 28.4. C₈H₇N₃· $\frac{1}{2}$ H₂O requires C, 63.6; H, 5.1; N, 27.8%).

(b) 4-Chloroquinazoline (1 g.) and aqueous ammonia (50 c.c.; *d* 0.88) after 10 days at room temperature gave a crude product (0.8 g., m. p. 202–215°) which, after dissolution in warm 0.5N-hydrochloric acid, filtration, and reprecipitation, yielded fairly pure 4-aminoquinazoline (0.59 g.; 67%), m. p. 245–251° (decomp.). This (0.5 g.) was suspended in water (80 c.c.), treated with sufficient acetic acid to effect dissolution, and basified with ammonia; the solution, after filtration, in the hot, from a little gelatinous material, deposited glittering needles of pure 4-aminoquinazoline, m. p. 265–267° (decomp.).

4-Acetamidoquinazoline (60 mg.) was prepared by heating the base (100 mg.) with acetic anhydride (1 c.c.) under reflux for 2 hours, and then adding dry ether (5 c.c.) to the cooled solution; it formed fine, colourless needles, m. p. 174–175° (Found: C, 64.0; H, 4.7; N, 22.5. Calc. for C₁₀H₉ON₂: C, 64.15; H, 4.85; N, 22.45%).

Reaction of 4-Phenoxyquinazoline with Methyl Toluene-*p*-sulphonate.—A mixture of 4-phenoxyquinazoline (3.52 g.) and methyl toluene-*p*-sulphonate (3.2 g.) was fused at 150° (bath temperature) for $\frac{1}{2}$ hour; a strong odour of phenol developed and a clear orange-red melt was formed. When the mixture had cooled, water (25 c.c.) was added, and the mixture extracted with ether; the ether-soluble material was identified by conversion into tribromophenol (4.4 g.; 84%), m. p. 94–95°. Treatment of the aqueous solution with picric acid (4 g.) in hot ethanol (80 c.c.) gave 1-methyl-4-quinazoline *picrate* (5.37 g.; 87%), which formed yellow needles, m. p. 249–250° (decomp.), from methanol, in which it was sparingly soluble (Found: C, 46.5; H, 3.05; N, 18.0. C₁₅H₁₁O₈N₅ requires C, 46.3; H, 2.85; N, 18.0%).

The free base (crude, 1.25 g.), obtained from the *picrate* (5.3 g.) by decomposition with ice-cold ammonia in the presence of chloroform, formed a hydrate, m. p. 65–66°, converted into the hygroscopic anhydrous base (long colourless needles, m. p. 141–142°, from chloroform-ligroin) by drying at 80–90° for 3 hours. The ammoniacal solution, after the removal of chloroform, was treated with 20% aqueous sodium hydroxide and kept for 24 hours at room temperature; extraction with benzene then yielded *o*-methylaminobenzamide (0.36 g.) (see below). Decomposition of the *picrate* with lithium hydroxide or sodium hydroxide gave poorer yields of the quinazoline. When a solution of the quinazoline (0.15 g.) in cold water (2 c.c.) was boiled for 2 minutes with 2N-sodium hydroxide (2 c.c.), some ammonia was evolved, and *o*-methylaminobenzamide (40 mg.) separated on cooling, m. p. 162–163° (Knape, J. *pr. Chem.*, 1891, 43, 209, gives m. p. 159–160°), after crystallisation from aqueous alcohol (Found: C, 63.8;

1356 *Keneford, Morley, Simpson, and Wright: The Chemistry of*

H, 6.5. Calc. for $C_8H_{10}ON_2$: C, 64.0; H, 6.7%. The quinazalone (100 mg.) was added in portions to a mixture of nitric acid (0.5 c.c.; d 1.50) and concentrated sulphuric acid (0.5 c.c.) at 25–30°. After 20 minutes the solution was poured on ice, made just alkaline with solid potassium carbonate, and filtered, and the filtrate was evaporated under reduced pressure, yielding 6-nitro-1-methyl-4-quinazalone (30 mg.), m. p. 270–272° alone and mixed with an authentic specimen (*J.*, 1948, 360).

3-Methyl-4-quinazalone.—The method of Bogert and Geiger (*J. Amer. Chem. Soc.*, 1912, **34**, 524) gave the base in 85% yield, but the yield was somewhat lower when methyl sulphate and excess of alkali were used. The anhydrous base, m. p. 105–106° (Bogert and Geiger give m. p. 105°), readily reverted to the hydrate, m. p. 71–72° (Bogert and Geiger give m. p. 70–71°); both forms gave large depressions in m. p. when mixed with the corresponding specimens of 1-methyl-4-quinazalone. The *picrate*, m. p. 215–216° (depressed by 1-methyl-4-quinazalone *picrate*), formed soft, yellow needles from alcohol, in which it was much more soluble than the 1-methyl isomer (Found: C, 46.6; H, 3.05; N, 18.1. $C_{15}H_{11}O_8N_5$ requires C, 46.3; H, 2.85; N, 18.0%).

Reaction of 4-Methoxyquinazoline with Methyl Toluene-*p*-sulphonate.—4-Methoxyquinazoline (0.64 g.) and methyl toluene-*p*-sulphonate (0.8 g.) were heated together for a few minutes at 160°. The colourless crystalline mass obtained on cooling was dissolved in warm water (2 c.c.) and treated with a solution of potassium iodide (1 g.) in water (2 c.c.), yielding the iodide (1.01 g.; 83%) as colourless needles, m. p. 275–277° after sintering at 265° [Bogert and Geiger, *loc. cit.*, give m. p. 275° (corr.)] (Found: C, 39.75; H, 3.7; N, 9.7. Calc. for $C_{10}H_{11}ON_2I$: C, 39.75; H, 3.7; N, 9.3%). The corresponding *picrate*, obtained from the iodide and picric acid in aqueous solution, crystallised from water in soft yellow plates, m. p. 197–198° (Found: C, 47.65; H, 3.35; N, 17.35. $C_{16}H_{13}O_8N_5$ requires C, 47.65; H, 3.25; N, 17.4%). A solution of the iodide (0.2 g.) in water (5 c.c.) was treated with 2N-sodium hydroxide (5 c.c.) and left overnight. A colourless oil (insoluble in water, readily soluble in dilute acids and in ether, benzene, or chloroform) had then separated; this (0.13 g.) was isolated by extraction with ether and converted into a *picrate* (0.25 g.), which separated from alcohol in yellow needles, m. p. 188–189° (153–155° when mixed with the *picrate*, m. p. 197–198°, described above) (Found: C, 46.4; H, 3.85; N, 18.2, 17.9, 18.3. $C_{16}H_{13}O_8N_5 \cdot \frac{1}{2}H_2O$ requires C, 46.6; H, 3.4; N, 17.0%).

Alkaline Decomposition of Cinnoline Methiodide.—(a) N-Sodium hydroxide (5 c.c.) and a solution of cinnoline methiodide (0.4 g.) in water (10 c.c.) were added alternately, in small portions, to a solution of potassium ferricyanide (1.2 g.) in water (10 c.c.) during 5 minutes; the deep emerald-green solution thus formed was kept at room temperature for $\frac{1}{2}$ hour with occasional shaking. Addition of excess of 20% sodium hydroxide and extraction with chloroform gave cinnoline, isolated as its *picrate* (0.26 g.; 49%), m. p. and mixed m. p. 196–197° (decomp.), and further authenticated by reversion into the methiodide [m. p. and mixed m. p. 171–173° (decomp.)] with warm methanolic methyl iodide.

(b) Cinnoline methiodide (0.2 g.) and 0.5N-sodium hydroxide (5 c.c.) were set aside under nitrogen for $\frac{1}{2}$ hour at room temperature. The deep-blue suspension was filtered into a separating funnel under nitrogen, giving a residue (A); the filtrate was treated with excess of 20% sodium hydroxide and extracted with benzene, but only a trace of oil was obtained on evaporation. The blue-green residue (A) (0.1 g., m. p. 120–125° (decomp.)) was almost insoluble in water and in benzene, but dissolved in dilute mineral acids giving pale red solutions from which it was reprecipitated by ammonia or caustic alkalis; it showed no tendency to form a *picrate*.

We are indebted to the Medical Research Council for a Research Studentship (J. S. M.).

WARRINGTON YORKE DEPARTMENT OF CHEMOTHERAPY,
LIVERPOOL SCHOOL OF TROPICAL MEDICINE.

[Received, August 27th, 1948.]