and nine-tenths grams of 2-amino-4-methyl-5-bromothiazole was suspended in 100 cc. of water. Two grams of thiourea was added and the mixture shaken until all solids had dissolved. The solution was made very alkaline by adding solid sodium hydroxide. In a few seconds, the characteristic glistening crystals of bis-(2-amino-4methyl-5-thiazolyl) sulfide were thrown out. The yield was 0.3 g. With some bis-(2-amino-4-methyl-5-thiazolyl) sulfide previously prepared there was no melting point depression.

FRICK CHEMICAL LABORATORY PRINCETON UNIVERSITY PRINCETON, N. J. RECEIVED JUNE 17, 1949

The Hydroxymethylation of Kojic Acid

Br L. L. Woods

The reaction of formaldehyde with kojic acid introduces one hydroxymethyl group in either the 3 or the 6 position of the pyrone ring. An earlier report¹ indicates that both of these positions are reactive so that a definite structure cannot be assigned to the derivative herein described. However, the compound is probably 2,6-bis-(hydroxymethyl)-5-hydroxy-4-pyrone since base catalyzed reactions² presumably require a mechanism which depends upon the nucleophilic character of the phenolic ion, thus causing the primary substitution to take place at a position ortho to the phenolic hydroxyl, in this instance position 6.

Treatment of the above product with thionyl chloride, followed by recrystallization from boiling water, gives a substance containing only one chlorine atom. The 2-chloromethyl group is assumed to remain intact since Yabuta³ has shown that 2-chloromethyl-5-hydroxy-4-pyrone resists hydrolysis by hot water.

Attempts to introduce two hydroxymethyl groups into kojic acid under forcing conditions always gave mixtures whose composition approached that of 2,3,6-tri-(hydroxymethyl)-5-hydroxy-4-pyrone, but the analysis of the benzoate indicated that it was contaminated with condensation products with similar physical properties.

Experimental⁴

Two grams of paraformaldehyde was added to a boiling mixture of 10 g. of kojic acid in 100 ml. of absolute ethanol, the solution was cooled slightly and 1 g. of potassium bicarbonate or anhydrous sodium carbonate was added with stirring. After standing for two hours, the solid was filtered and dried in a vacuum desiccator over sulfuric acid. Recrystallization of the crude product (9.8 g.) from absolute ethanol gave very pale yellow needles which melted at 155-156°, following an apparent change in structure above 132°, and which gave a red-purple coloration with ferric chloride solution.

Anal. Calcd. for $C_7H_8O_5$: C, 48.83; H, 4.65. Found: C, 49.20, 49.02; H, 4.37, 4.42.

Two grams of the above compound was refluxed with 7

(2) Price, "Reactions at Carbon-Carbon Double Bond," Interscience Publishers, New York, N. Y., 1946, p. 46.

(3) Yabuta, J. Chem. Soc., 125, 575 (1924).

ml. of thionyl chloride. The product, weighing 0.5 g. after recrystallization from 25 ml. of boiling water, separated either as needles or stout prisms which changed to needles between 120 and 140° and melted at 165°.

Anal. Caled. for C₇H₇O₄C1: C, 44.38; H, 3.16. Found: C, 44.71, 44.60, 44.68; H, 3.24, 3.08, 3.18.

Benzoylation of 2 g. of the compound by the usual Schotten-Baumann method produced a white solid which was recrystallized three times from absolute ethanol; yield $2.2 \text{ g., m. p. } 134.5-135.5^{\circ}$.

Anal. Calcd. for $C_{28}H_{20}O_8$: C, 69.42; H, 4.13. Found: C, 69.04; H, 4.08.

In an attempt to introduce two hydroxymethyl groups into the pyrone nucleus, a mixture of 10 g. of kojic acid, 3.8 g. of paraformaldehyde, 1 g. of potassium bicarbonate and 50 ml. of absolute ethanol was heated for seventeen hours at 75°. Two ml. of concentrated hydrochloric acid was then added, the mixture was refluxed for thirty minutes, treated with Norite and filtered. After standing in the refrigerator for two days, the solution deposited 5.4 g. of a brown solid which was not appreciably soluble in alcohol or other organic solvents. Recrystallization from water, followed by digestion with absolute ethanol, gave a yellow powder which decomposed above 233° and gave a red coloration with ferric chloride.

Anal. Caled. for $C_8H_{10}O_6\colon$ C, 47.52; H, 4.95. Found: C, 47.99; H, 4.70.

Benzoylation of 1.8 g. of the above powder with 12 ml. of benzoyl chloride at 100–110° for one hour, followed by the action of 10% sodium hydroxide, gave a solid which was recrystallized once from benzene and four times from ethanol. The product, a yellow powder melting at 99–101°, failed to give the proper analysis, an indication that the substance from which the benzoate was prepared probably contained significant amounts of condensation products similar to 2,3,6-tri-(hydroxymethyl)-5-hydroxy-4-pyrone.

Anal. Caled. for $C_{36}H_{26}O_{19}$: C, 69.90; H, 4.20. Found: C, 68.71; H, 4.00.

DEPARTMENT OF CHEMISTRY

TEXAS STATE UNIVERSITY FOR NEGROES

HOUSTON, TEXAS RECEIVED DECEMBER 5, 1949

Calcium Hydride as a Basic Catalyst in the Perkin Reaction

By Saverio Zuffanti and John J. Sardella

Gibb^{1,2} and others³⁻⁶ have investigated the use of calcium hydride in the condensations of aldehydes, ketones and esters.

This note reports data on the use of calcium hydride in the Perkin reaction. Benzaldehyde has been treated with acetic, propionic, and *n*-butyric anhydrides.

Experimental

Materials.—The calcium hydride was the commercial gray grade of about 95% purity. The benzaldehyde was purified by keeping it in contact with calcium hydride for several days and then collecting the fraction distilling at 178–180°. The acetic, propionic and *n*-butyric anhydrides were purified by allowing each to stand over metallic so-dium for a few days, distilling, then allowing to stand over calcium hydride for a few days, and distilling again before use.

(3) Taboury and Godchot, Compt. rend., 169, 62-64 (1919).

(1925).

(6) Packendorff, Ber., 64B, 948-949 (1931).

⁽¹⁾ Woods, This Journal, 68, 2744 (1946).

⁽⁴⁾ The kojic acid was obtained from the Corn Products Sales Company. Analyses were by Dr. Carl Tiedcke. The Fisher-Johns apparatus was used for all melting points

⁽¹⁾ Gibb and Mann, M. I. T., B.S. Thesis, 1946.

⁽²⁾ Gibb and Horne, M. I. T., B.S. Thesis, 1946.

⁽⁴⁾ Porlezza and Gatti, Gazz. chim. ital., 54, 491 (1924); 55, 224

⁽⁵⁾ Perkin and Pratt, J. Chem. Soc., 95, 159 (1909).