

HIGHLY ENANTIOSELECTIVE REDUCTION OF δ -KETO ACIDS WITH FERMENTING BAKER'S YEAST.
A FACILE SYNTHESIS OF OPTICALLY PURE (R)-(+)-5-HEXADECANOLIDE


Masanori UTAKA, Hisashi WATABU, and Akira TAKEDA*

Department of Synthetic Chemistry, School of Engineering,
Okayama University, Tsushima, Okayama 700

Optically pure (R)-(+)-hexadecanolide, (+)-5-tridecanolide, and (+)-5-nonanolide were prepared from the corresponding δ -keto acids by reduction with fermenting baker's yeast.

5-Hexadecanolide (1a), a pheromone isolated from the oriental hornet (*vespa orientalis*) in 1969,¹⁾ has induced ten attempts to synthesize the optically active form.²⁾ Here we wish to report the eleventh synthesis of 1a that involves the most effective method of obtaining the optically pure (R)-(+)-enantiomer of 1a by using fermenting baker's yeast as shown in Scheme 1. Use of baker's yeast in the reduction of δ -keto acid 4a was once reported to be unsatisfactory because the optical yield was only 39%.^{2d,h)} We have not yet clarified why our reduction was highly enantioselective.

2-Undecylcyclohexanone (2a)³⁾ was converted via its 2,6-dibromo derivative to 3-undecyl-1,2-cyclohexanedione (3a) in 57% yield.⁴⁾ The dione 3a reacted readily with oxygen by catalysis of CuCl_2 to afford δ -keto acid 4a in 81% yield.⁵⁾ A suspension of baker's yeast (2.76 g, Nisshin Seihun Co.), glucose (3.10 g), KH_2PO_4 (7 mg), MgSO_4 (4 mg) in boiled water (9 ml) was stirred for 30 min at 25 °C. To the fermenting mixture was added 4a (135 mg, 0.50 mmol) dissolved in water by addition of 1 M KOH and the resulting mixture was adjusted to pH 7 with 1 M KOH.⁶⁾ After stirring for 5 h at 25 °C, the glucose added initially was consumed as checked by a test paper (Diastix-II, Miles-Sankyo Co.). Then 6.2 g of glucose was added to the mixture. After 17 h, when the vigorous fermentation appeared to cease, the equal amount of the initially used yeast-glucose suspension was added and stirring was continued at 25 °C until the total elapse of fermentation reached 48 h. The mixture was stirred with Celite (8.2 g) for 30 min and filtered. The Celite was washed with water and then with ether three times. The filtrate and washings were combined, acidified to pH 2 with 10% HCl, and extracted with ether five times. The combined ethereal solution, after drying (MgSO_4) and evaporation,



Scheme 1.

gave a crude oil (305 mg) which was refluxed in benzene for 1 h with *p*-TsOH. The crude product (293 mg) was purified by passing through a silica gel column (hexane-acetone 30:1) to give 129 mg of a semisolid. This was further purified by HPLC (silica gel, hexane-AcOEt 5:1) to afford 51 mg (40%) of (R)-(+)-1a as colorless crystals, mp 37.5–38 °C (lit.^{2h}) 37 °C), $[\alpha]_D^{23} +39.5^\circ$ (c 1.74, THF) (lit.^{2h}) $[\alpha]_D^{21.5} +40.2^\circ$ (c 1.76, THF). In order to determine the optical purity accurately, 1a was converted to the diol with MeLi,⁷⁾ followed by ¹H FT NMR examination of the diol in the presence of Eu(hfc)₃. The two methyl groups attached to the quaternary carbon atom appeared as two singlets without any trace of paired signals,⁸⁾ indicating that the optical purity was >98% ee.

In similar ways, the use of fermenting baker's yeast in the reduction of 5-oxotridecanoic and 5-oxononanoic acids (4b,c) afforded the corresponding (+)-5-tridecanolide (1b) and (+)-5-nonanolide (1c) in 54% yield (>98% ee, $[\alpha]_D^{26} +45.2^\circ$ (c 1.58, THF)) and 67% yield (>98% ee, $[\alpha]_D^{27} +58.1^\circ$ (c 3.66, THF)) (lit.⁹⁾ $[\alpha]_D^{20} +58.2^\circ$ (MeOH)), respectively.

In conclusion, we have found that the δ -keto group can be reduced by fermenting baker's yeast with high enantioselectivity. This methodology can be applied to the synthesis of optically pure δ -lactones effectively.

References

- 1) R. Ikan, R. Gottlieb, E. D. Bergmann, and J. Ishay, J. Insect Physiol., **15**, 1709 (1969).
- 2) a) J. L. Coke and A. B. Richon, J. Org. Chem., **41**, 3516 (1976); b) W. H. Pirkle and P. E. Adams, *ibid.*, **44**, 2169 (1979); c) G. Solladie and F. Matloubi-Moghadam, *ibid.*, **47**, 91 (1982); d) N. Naoshima, H. Ozawa, H. Kondo, and S. Hayashi, Agric. Biol. Chem., **47**, 1431 (1983); e) S. Servi, Tetrahedron Lett., **24**, 2023 (1983); f) M. Larcheveque and J. Lalande, Tetrahedron, **40**, 1061 (1984); g) T. Kikukawa and A. Tai, Chem. Lett., **1984**, 1935; h) K. Mori and T. Otsuka, Tetrahedron, **41**, 547 (1985); i) H. Kosugi, H. Konta, and H. Uda, J. Chem. Soc., Chem. Commun., **1985**, 211; j) T. Fujisawa, T. Itoh, M. Nakai, and T. Sato, Tetrahedron Lett., **26**, 771 (1985).
- 3) Prepared from cyclohexylidenecyclohexylamine, EtMgBr, and n-C₁₁H₂₃Br.
- 4) M. Utaka, S. Matsushita, and A. Takeda, Chem. Lett., 1980, 779.
- 5) M. Utaka, M. Hojo, Y. Fujii, and A. Takeda, Chem. Lett., **1984**, 635.
- 6) We used the procedure reported by M. Hirama et al. with modification. See: M. Hirama, M. Shimizu, and M. Iwashita, J. Chem. Soc., Chem. Commun., **1983**, 599.
- 7) I. J. Jakovac and J. B. Jones, J. Org. Chem., **44**, 2165 (1979).
- 8) The diol derived from an enantiomeric mixture of 1a showed two pairs of singlets in the ¹H NMR (100 MHz) spectrum in the presence of Eu(hfc)₃.
- 9) G. T. Muys, B. Van der Ven, and A. P. De Jonge, Appl. Microbiol., **11**, 389 (1963).

(Received July 16, 1985)