

Flory⁴ for monomers in which the reacting double bond is not in resonance.

(4) Flory, *THIS JOURNAL*, **59**, 241 (1937).


RESEARCH LABORATORIES
EASTMAN KODAK COMPANY
KODAK PARK WORKS
ROCHESTER 4, N. Y.

RECEIVED JULY 2, 1947

Dihydropyran Addition Products


BY G. FORREST WOODS AND DAVID N. KRAMER

2,3-Dihydropyran is the dehydration product of a hemiacetal and as such has been shown to add water¹ readily in the presence of a trace of mineral acid yielding an equilibrium mixture of 5-hydroxypentanal and 2-hydroxytetrahydropyran. Paul^{1a} observed that methyl alcohol




added to dihydropyran and from this reaction 2-methoxytetrahydropyran was obtained.

We have prepared a number of acetals from 2,3-dihydropyran according to the equation below modifying the procedure of Paul



These acetals are stable in a basic medium and are therefore easily isolated after destruction of the acid catalyst with a strong inorganic base.

pyran the yellow 2,4-dinitrophenylhydrazone of 5-hydroxypentanal, the hydrolysis product of dihydropyran, has been used. We have found that 3,5-dinitrobenzoic acid when mixed with an excess of dihydropyran forms an addition product which is excellent for characterization. This product does not form a semicarbazone when treated with free semicarbazide in methanol. Therefore, the structure (I) is preferred to that of (II) for this substance.



Experimental

Acetal Formation.—Equimolar quantities of alcohol and dihydropyran in the presence of a trace of concentrated hydrochloric acid were shaken and allowed to stand for three hours. The reaction is exothermic. A few pellets of sodium hydroxide were then added to the reaction mixture to destroy the acid, and the product isolated directly by distillation. The pure material was obtained by redistillation in the presence of a few pellets of sodium hydroxide.

The 2,4-dinitrophenylhydrazone of 5-hydroxypentanal from these acetals were prepared in the usual manner. The yellow crystalline material obtained melted at 107–109° and gave no depression in a mixed melting point determination with an authentic sample.^{1c}

The 3,5-Dinitrobenzoate of 2-Hydroxytetrahydropyran.—Five grams of 3,5-dinitrobenzoic acid was dissolved with warming in a 50% excess of dihydropyran. Upon cooling, 5 cc. of ether was added and the product slowly crystal-

TABLE I
ACETALS FROM DIHYDROPYRAN

R =	B. p., °C.	<i>n</i> _D	Yield, %	C	H	Calcd., %	C	H	Found, %
Methyl ^a	125	1.4260	85	62.07	10.35		62.06	10.01	
Ethyl ^b	146	1.4248	93	64.62	10.76		65.04	10.58	
<i>n</i> -Propyl-	165	1.4280	91	66.67	11.11		66.27	11.00	
Allyl-	126	1.4440	70	67.53	9.86		67.48	10.07	
<i>n</i> -Butyl-	183	1.4312	75	68.39	11.33		68.45	11.36	
Phenyl-	103 (4 mm.)	1.5290	37	74.16	7.87		73.96	7.92	
Benzyl-	107 (3 mm.)	1.5128	41	75.00	8.33		74.63	8.40	
Furfuryl-	124 (24 mm.)	1.4828	34	65.92	7.78		65.96	8.06	
2,2'-(<i>sym</i> -Ethylenedioxy-ditetrahydropyran)	164 (32 mm.)	1.4610	40	62.53	9.56		62.54	9.74	

^a Cf. ref. 1a. ^b This substance has been prepared by catalytic hydrogenation of 2-ethoxy-Δ³-dihydropyran [Woods and Sanders, *THIS JOURNAL*, **68**, 2483 (1946)]. ^c In the addition of equimolar quantities of ethylene glycol to dihydropyran, mixtures of di-tetrahydropyran and mono-tetrahydropyran addition products are obtained. Ratios of two moles of ethylene glycol to one mole of dihydropyran were used to favor the preparation of the monomer which could not be obtained in a pure state. The dimer was obtained by adding two moles of dihydropyran to one mole of ethylene glycol. In each case, the two products could be separated from the initial reaction mixture by fractional distillation.

Hydrolysis of these acetals by acids in an aqueous medium in the presence of 2,4-dinitrophenylhydrazine yielded 5-hydroxypentanal-2,4-dinitrophenylhydrazone.

For purposes of identification of 2,3-dihydro-

lized as needles. After a second recrystallization from 80% dihydropyran-ether, a pale yellow solid was obtained which melted sharply at 103°. The yield was essentially quantitative.

Anal. Calcd. for C₁₂H₁₂O₇N₂: C, 48.65; H, 4.05. Found: C, 49.05; H, 4.46.

CHEMISTRY DEPARTMENT
UNIVERSITY OF MARYLAND
COLLEGE PARK, MARYLAND

RECEIVED APRIL 1, 1947

(1a) R. Paul, *Bull. soc. chim.*, [5] **1**, 973 (1934).

(1b) Schniepp and Geller, *THIS JOURNAL*, **68**, 1646 (1946).

(1c) Woods and Sanders, *ibid.*, **68**, 2111 (1946).