



Synthesis of 1,2,5,6- and 1,4,5,8-anthracenetetrone: Building blocks for π -conjugated small molecules and polymers

Florian Glöcklhofer, Berthold Stöger & Johannes Fröhlich

To cite this article: Florian Glöcklhofer, Berthold Stöger & Johannes Fröhlich (2018): Synthesis of 1,2,5,6- and 1,4,5,8-anthracenetetrone: Building blocks for π -conjugated small molecules and polymers, Synthetic Communications, DOI: [10.1080/00397911.2018.1483027](https://doi.org/10.1080/00397911.2018.1483027)

To link to this article: <https://doi.org/10.1080/00397911.2018.1483027>



© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



[View supplementary material](#)



Published online: 17 Aug 2018.



[Submit your article to this journal](#)



Article views: 200



[View Crossmark data](#)

Synthesis of 1,2,5,6- and 1,4,5,8-anthracenetetrone: Building blocks for π -conjugated small molecules and polymers


Florian Glöcklhofer^a , Berthold Stöger^b  and Johannes Fröhlich^a 

^aInstitute of Applied Synthetic Chemistry, TU Wien, Vienna, Austria; ^bX-Ray Centre, TU Wien, Vienna, Austria

ABSTRACT

Reliable reactions for the synthesis of two interesting anthracenetetrone derivatives have been identified and optimized. Both syntheses start from dihydroxy-9,10-anthraquinones and were selected for maximized efficiency and minimized workload. Work-up of all reactions can be achieved without column chromatography, which facilitates further scale-up. So far, both target compounds are considerably underexplored despite their promising molecular structure for use in devices and in organic synthesis, especially as building blocks for π -conjugated compounds. The crystal structure of 1,4,5,8-anthracenetetrone is reported.

GRAPHICAL ABSTRACT



ARTICLE HISTORY

Received 8 May 2018

KEYWORDS

Anthracenetetrone;
anthraquinone; building
block; crystal
structure; tetrone

Introduction

Anthracenetetrone derivatives are interesting compounds for technological applications and further conversion. For example, 1,4,9,10-anthracenetetrone (Scheme 1, top), which is currently the best investigated anthracenetetrone, was used to modify electrodes for supercapacitors,^[1] was applied as acceptor in charge-transfer complexes,^[2–4] it served as a precursor in the synthesis of antitumor compounds,^[5] and afforded push-pull chromophores in cycloaddition reactions with electron-rich alkynes.^[6] Investigations of this anthracenetetrone are facilitated by its facile synthesis in a single high-yielding reaction that uses


CONTACT Florian Glöcklhofer  florian.gloecklhofer@tuwien.ac.at  Institute of Applied Synthetic Chemistry, TU Wien, Getreidemarkt 9/163, Vienna 1060, Austria.

 Supplemental data for this article can be accessed on the [publisher's website](#).

Color versions of one or more of the figures in the article can be online at www.tandfonline.com/lsyc.

© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Scheme 1. Facile synthesis of 1,4,9,10-anthracenetrone starting from 1,4-dihydroxy-9,10-anthraquinone (top), molecular structure of 1,2,5,6- and 1,4,5,8-anthracenetetrone (bottom).


low-cost 1,4-dihydroxy-9,10-anthraquinone as the starting material (Scheme 1, top).^[7] Different to other anthracenetetrone, also the crystal structure of this compound has been reported.^[8]

In contrast, the synthesis of 1,2,5,6- and 1,4,5,8-anthracenetetrone (Scheme 1, bottom) is more challenging.^[9–12] This may explain why these compounds are less investigated despite the large number of available reactions for converting quinones, including our new conversion into cyanoarenes (demonstrated for 5,7,12,14-pentacenetetrone and a range of other para- and ortho-quinones).^[13–16] Beside other syntheses, the two tetrone are particularly interesting for the synthesis of π -conjugated small molecules and polymers, but they have been used in only a few reactions so far.

1,2,5,6-Anthracenetetrone was used for condensation with aromatic ortho-diamines, a reaction first demonstrated by Boldt.^[9] This reaction yields fully aromatic compounds by connecting the building blocks via pyrazine units. It has later been shown that the corresponding polycondensation with tetraamines yields fully conjugated ladder polymers.^[17] The polymers have not yet been tested in devices, but Bunz et al. recently optimized the condensation for larger ortho-diamines and tested the resulting compounds in organic light-emitting diodes.^[18] Regarding applications, environmentally benign methods for (poly)condensations, which afford highly crystalline materials, open up new opportunities.^[19,20]

1,4,5,8-Anthracenetetrone, on the other hand, was used as a dienophile in double Diels–Alder cycloaddition reactions, which – depending on the diene – can yield macrocyclic compounds,^[21] substituted 5,7,12,14-pentacenetetrone,^[22] and pentacene precursors.^[23] The absorption and electrochemical properties of this tetrone, which is the most stable anthracenetetrone according to a computational study,^[24] were reported decades ago.^[11,25] In a more recent study, the binding of this tetrone on graphene was computationally predicted to be the strongest of several electroactive compounds for green battery electrodes.^[26] Such strong binding benefits the cycling performance of batteries.

It was our aim to facilitate the use and investigation of 1,2,5,6- and 1,4,5,8-anthracenetetrone beyond computational studies by identifying and evaluating reliable synthetic



Scheme 2. (i) Reduction of 2,6-dihydroxy-9,10-anthraquinone (**1a**) to 2,6-dihydroxyanthracene (**2a**). Conditions: NaBH_4 , 1 M Na_2CO_3 solution, under argon, RT, overnight. (ii) Oxidation to 1,2,5,6-anthracenetetrone (**3a**). Conditions: benzeneseleninic acid anhydride, dry THF, under argon, 50°C , 3 h. (iii) Conversion of 1,8-dihydroxy-9,10-anthraquinone (**1b**) into 1,4,5,8-tetramethoxyanthracene (**2b**). Conditions: see Ref. [28], and (iv) Oxidation to 1,4,5,8-anthracenetetrone **3b**. Conditions: silica-supported ceric ammonium nitrate (CAN), CH_2Cl_2 , RT, 1 h.

routes to these tetrone, starting from dihydroxy-9,10-anthraquinones (as for the synthesis of 1,4,9,10-anthracenetetrone). The syntheses should be optimized regarding workload and efficiency.

Results and discussion

For the synthesis of 1,2,5,6-anthracenetetrone **3a**, oxidation of 2,6-dihydroxyanthracene **2a** (Scheme 2, top) using benzeneseleninic acid anhydride was reported to be superior to other reactions.^[10] Fortunately, the required precursor **2a** can be efficiently prepared from 2,6-dihydroxy-9,10-anthraquinone **1a**. Screening the literature for reliable protocols to minimize the workload and maximize the yields, we discovered a protocol reported by Guo et al.^[27] Following this protocol, we immediately achieved 86% yield (just 2% less than reported). There is no need for column chromatography or recrystallization to achieve high purity, which significantly reduces the workload.

In contrast to the synthesis of **2a**, the subsequent oxidation to **3a** required considerable optimization. Initially, we followed a protocol reported by Boldt et al.,^[10] which relies on crystallization of **3a** from the reaction medium at room temperature and additionally requires extraction and crystallization steps to improve the yield to 71%. Unfortunately, we did not observe any crystallization of **3a** at room temperature (also storing at -18°C overnight just afforded small amounts of precipitate). As we wanted to simplify the protocol anyways, we decided to directly evaporate the reaction solvent as well as volatile by-products instead of storing the reaction for crystallization. The resulting solid was then recrystallized from 1,4-dioxane affording tetrone **3a** as red needles. Rapid weathering prevented analysis of the crystal structure, but NMR spectroscopy confirmed the purity, except for some 1,4-dioxane, which disappears during storage or when heating the solid, as was observed before.^[9] Very good yields of 85% were obtained.

Our other target compound, 1,4,5,8-anthracenetetrone **3b**, can be prepared by oxidation of 1,4,5,8-tetrahydroxyanthracene using freshly prepared silver oxide,^[29] but the yields of this reaction (and the five reactions needed to prepare the tetrahydroxy anthracene) were reported to be unsatisfactory by Miller et al.^[11] Instead, they suggest to oxidize 5,8-dimethoxy-1,4-anthraquinone, which they prepared in a sequence of seven reactions. Obviously, both routes do not comply with our aims of high yields and low workload.

Fortunately, tetrone **3b** can also be prepared by oxidation of 1,4,5,8-tetramethoxyanthracene **2b** (Scheme 2, bottom) using ceric ammonium nitrate (CAN),^[12] a reaction that seemed more promising to us despite the reported moderate yield of 45% and the not reported experimental details and characterization results. The required tetramethoxy anthracene **2b** can be prepared from 1-bromo-2,5-dimethoxybenzene.^[30] The easy accessibility of this precursor certainly is an advantage of this reaction, but the low yield of 32% as well as the need for lithium 2,2,6,6-tetramethylpiperidide (LTMP) as the reagent are disadvantages that have to be considered. We decided to follow a different approach starting from low-cost 1,8-dihydroxy-9,10-anthraquinone **1b** (Scheme 2, bottom), as this approach was reported to be useful also on a larger scale:

Starting from **1b**, an overall yield of 68% was reported for the synthesis of **2b** by methylation of the hydroxyl groups, subsequent reduction to 1,8-dimethoxy anthracene, 2-fold bromination to 1,8-dibromo-4,5-dimethoxy anthracene, and final replacement of the bromo substituents by methoxy groups.^[28] Despite carrying out this reaction sequence on a 50% larger scale, we achieved exactly the same overall yield at the first try (lower yield for the methylation, higher yield for the reduction, similar yields for the two other steps). We followed the published protocol with slight adaptations: the crude methylation product was concentrated *in vacuo* and redissolved in CH₂Cl₂ before passing through a pad of silica, 2 N NaOH was used for washing the crude bromination product instead of 5% HCl, and **2b** was triturated in boiling CH₂Cl₂ instead of room temperature. In contrast to the alternative synthesis of **2b**, which uses 1-bromo-2,5-dimethoxybenzene as a precursor, none of the reactions required purification by column chromatography. Nevertheless, the synthesis using 1-bromo-2,5-dimethoxybenzene may be more convenient, if only small amounts of **2b** are required.

For the final oxidation to target compound **3b**, we developed a protocol based on a work using silica-supported CAN for the synthesis of benzoquinones.^[31] Besides other advantages, silica-supported CAN can be easily removed by filtration after the reaction. Furthermore, it allows for using CH₂Cl₂ as the solvent, which dissolves **2b** better than the usual aqueous acetonitrile. The reaction was carried out simply by adding CH₂Cl₂ and **2b** to freshly prepared silica-supported CAN and stirring for 1 h at room temperature. Facile work-up afforded crude **3b** in quantitative yields. For removing impurities, the solid was recrystallized from 1,4-dioxane, which reduced the yield to 64%. Despite the loss during recrystallization, this is still significantly higher than reported before without experimental details.

In contrast to **3a**, needles of **3b** were suitable for structural characterization (Figure 1 and Supplementary Material). In the solid state, **3b** is essentially flat (largest distances from least-squares plane defined by the C-atoms: O2, 0.0385(9) Å and C3, 0.0183(10) Å). It crystallizes with one molecule of 1,4-dioxane in the space group *P* $\bar{1}$. The asymmetric unit contains half of a formula unit (*Z'* = 1/2), whereby **3b** and 1,4-dioxane




Figure 1. Crystal structure of **3b**. 1,4-dioxane viewed down (a) [100] and (b) [010]. (c) The molecular structure of **3b**. C (grey) and O (red) atoms are represented by ellipsoids drawn at the 50% probability levels; H atoms by white spheres of arbitrary radius.

are located each on a center of inversion. No significant π - π stacking is observed. In addition to these results, we were also able to determine the crystal structure of the precursor 1,8-dibromo-4,5-dimethoxyanthracene, which has not yet been reported (in contrast to the crystal structure of the direct precursor **2b**) and is provided as [Supplementary Material](#).

Conclusions

1,2,5,6- and 1,4,5,8-Anthracenetetrone **3a** and **3b** can be prepared in efficient reactions starting from dihydroxy-9,10-anthraquinones by following the presented synthetic routes. None of the reactions requires column chromatography for purification and the workload is generally low. For the synthesis of the precursor **2b**, an alternative procedure is available, which may be preferred on a smaller scale. Both **3a** and **3b** form solvates when recrystallized from 1,4-dioxane, but the solvent can be removed easily. The developed synthetic routes greatly facilitate the exploration of tetrones **3a** and **3b** in devices and in the synthesis of π -conjugated small molecules and polymers.

Experimental

Reagents and solvents were purchased from commercial suppliers and used without further purification. Anhydrous solvents were prepared by filtration through drying columns (methanol, THF) or used as received from commercial suppliers (acetone). Reactions were monitored by thin-layer chromatography (Merck, silica gel 60 F₂₅₄).

2,6-Dihydroxyanthracene **2a** was prepared from 2,6-dihydroxy-9,10-anthraquinone **1a** following a published protocol.^[27] The synthesis of 1,4,5,8-tetramethoxyanthracene **2b** from 1,8-dihydroxy-9,10-anthraquinone **1b** was also achieved following a published procedure, but with minor changes described above.^[28]

NMR spectra were recorded at 600 MHz for ¹H and 151 MHz for ¹³C on a Bruker Avance III HD spectrometer. ¹H NMR signals were assigned as suggested by MestReNova 12.0.1-20560 (Supplementary Material). For ¹³C NMR signals, the number of attached protons (C, CH) was assigned by attached proton test (APT) experiments.

Synthesis of 1,2,5,6-anthracenetetrone **3a**

2,6-Dihydroxyanthracene **2a** (263 mg, 1.25 mmol, 1.00 equiv.) was dissolved in 25 mL dry THF and purged with argon. The solution was then added in portions to a stirred solution of benzeneseleninic acid anhydride (900 mg, 2.50 mmol, 2.00 equiv.) in 50 mL dry THF under argon (in a three-necked round-bottom flask equipped with a reflux condenser with argon balloon and a thermometer). The reaction was heated to 50 °C for 3 h and was then allowed to cool to room temperature. After storing overnight, the solvent and volatile by-products were removed *in vacuo* using a rotary evaporator, which was placed in a fume hood to prevent inhalation of volatile selenium by-products. Recrystallization of the resulting solid from 1,4-dioxane afforded red needles (1,4-dioxane solvates), which were dried to afford pure tetrone **3a** (252 mg, 1.06 mmol) in yields of 85%. ¹H NMR (600 MHz, THF-d₈): δ = 8.10 (s, 2H), 7.64 (d, *J* = 10.2 Hz, 2H), 6.44 (d, *J* = 10.2 Hz, 2H) ppm (in accordance with the literature^[18]). ¹H NMR (600 MHz, CDCl₃): δ = 8.13 (s, 2H), 7.57 (d, *J* = 10.2 Hz, 2H), 6.62 (d, *J* = 10.2 Hz, 2H) ppm. ¹³C NMR (151 MHz, CDCl₃): δ = 179.5 (C), 177.9 (C), 143.3 (CH), 136.4 (C), 135.8 (C), 130.9 (CH), 130.1 (CH) ppm.

Synthesis of 1,4,5,8-anthracenetetrone **3b**

Ceric ammonium nitrate (CAN) (6.03 g, 11.0 mmol, 5.50 equiv.) was dissolved in 6 mL water and added dropwise to 13 g silica (in a stirred 100 mL round-bottom flask equipped with a septum). The solid was stirred until a free-flowing yellow solid was obtained (about 30 min). 25 mL of CH₂Cl₂ was then added, followed by the addition of 1,4,5,8-tetramethoxyanthracene **2b** (597 mg, 2.00 mmol, 1.00 equiv.) suspended in 35 mL CH₂Cl₂. Another 10 mL CH₂Cl₂ was used to transfer residues of **2b**. The reaction stirred at room temperature for 1 h and was then filtered through a sintered glass funnel to remove the silica. The silica was washed with 200 mL CH₂Cl₂ and the combined solutions were washed with water and dried over sodium sulfate. The solvent was removed *in vacuo* and the solid residue was recrystallized from 1,4-dioxane, which afforded red needles (1,4-dioxane solvates). After drying, tetrone **3b** (303 mg, 1.27 mmol) was obtained in yields of 64%. ¹H NMR (600 MHz, CDCl₃): δ = 8.82 (s, 2H), 7.14 (s, 4H) ppm (in accordance with the literature apart from reversed integrals^[11]). ¹³C NMR (151 MHz, CDCl₃): δ = 183.3 (C), 139.4 (CH), 135.1 (C), 125.8 (CH) ppm.

Acknowledgments

We thank Markus Lunzer and Andreas J. Morawietz for contributing to synthetic experiments. Brigitte Holzer and Christian Hametner are acknowledged for NMR measurements. The authors acknowledge the TU Wien University Library for financial support through its Open Access Funding Program.

Funding

This study was supported by the Technische Universität Wien.

ORCID

Florian Glöcklhofer  <http://orcid.org/0000-0002-6911-8563>

Berthold Stöger  <http://orcid.org/0000-0002-0087-474X>

Johannes Fröhlich  <http://orcid.org/0000-0003-0840-2314>

References

- [1] Isikli, S.; Díaz, R. *J. Power Sources* **2012**, *206*, 53–58. DOI:10.1016/j.jpowsour.2012.01.088.
- [2] Kaul, B. B.; Yee, G. T. *Polyhedron* **2001**, *20*, 1757–1759. DOI:10.1016/S0277-5387(01)00685-4.
- [3] Asahi, H.; Inabe, T. *Synth. Met.* **1995**, *70*, 1117–1118. DOI:10.1016/0379-6779(94)02780-3.
- [4] Asahi, H.; Inabe, T. *Chem. Mater.* **1994**, *6*, 1875–1879. DOI:10.1021/cm00046a050.
- [5] Schenck, L. W.; Kuna, K.; Frank, W.; Albert, A.; Asche, C.; Kucklaender, U. *Bioorg. Med. Chem.* **2006**, *14*, 3599–3614. DOI:10.1016/j.bmc.2006.01.026.
- [6] Dengiz, C.; Swager, T. M. *Synlett* **2017**, *28*, 1427–1431. DOI:10.1055/s-0036-1588771.
- [7] Cassis, R.; Valderrama, J. A. *Synth. Commun.* **1983**, *13*, 347–356. DOI:10.1080/00397918308066988.
- [8] Kitamura, C.; Kawase, T. *Acta Crystallogr., Sect. E* **2013**, *69*, o1597. DOI:10.1107/S1600536813026342.
- [9] Boldt, P. *Chem. Ber.* **1966**, *99*, 2322–2336. DOI:10.1002/cber.19660990731.
- [10] Zippel, S.; Boldt, P. *Synthesis* **1997**, *1997*, 173–175. DOI:10.1055/s-1997-1171.
- [11] Almlöf, J. E.; Feyereisen, M. W.; Jozefiak, T. H.; Miller, L. L. *J. Am. Chem. Soc.* **1990**, *112*, 1206–1214. DOI:10.1021/ja00159a049.
- [12] Cory, R. M.; McPhail, C. L.; Dikmans, A. J. *Tetrahedron Lett.* **1993**, *34*, 7533–7536. DOI:10.1016/S0040-4039(00)60392-1.
- [13] Glöcklhofer, F.; Lunzer, M.; Stöger, B.; Fröhlich, J. *Chem. - Eur. J.* **2016**, *22*, 5173–5180. DOI:10.1002/chem.201600004.
- [14] Glöcklhofer, F.; Kautny, P.; Fritz, P.; Stöger, B.; Fröhlich, J. *ChemPhotoChem* **2017**, *1*, 51–55. DOI:10.1002/cptc.201600018.
- [15] Glöcklhofer, F.; Morawietz, A. J.; Stöger, B.; Unterlass, M. M.; Fröhlich, J. *ACS Omega* **2017**, *2*, 1594–1600. DOI:10.1021/acsomega.7b00245.
- [16] Glöcklhofer, F.; Petritz, A.; Karner, E.; Bojdys, M. J.; Stadlober, B.; Fröhlich, J.; Unterlass, M. M. *J. Mater. Chem. C* **2017**, *5*, 2603–2610. DOI:10.1039/c7tc00143f.
- [17] Imai, K.; Kurihara, M.; Mathias, L.; Wittmann, J.; Alston, W. B.; Stille, J. K. *Macromolecules* **1973**, *6*, 158–162. DOI:10.1021/ma60032a002.
- [18] Hahn, S.; Koser, S.; Hodecker, M.; Tverskoy, O.; Rominger, F.; Dreuw, A.; Bunz, U. H. F. *Chem. - Eur. J.* **2017**, *23*, 8148–8151. DOI:10.1002/chem.201701304.
- [19] Baumgartner, B.; Bojdys, M. J.; Unterlass, M. M. *Polym. Chem.* **2014**, *5*, 3771–3776. DOI:10.1039/C4PY00263F.

- [20] Baumgartner, B.; Svirikova, A.; Binting, J.; Hametner, C.; Marchetti-Deschmann, M.; Unterlass, M. M. *Chem. Commun.* **2017**, 53, 1229–1232. DOI:[10.1039/c6cc06567h](https://doi.org/10.1039/c6cc06567h).
- [21] Cory, R. M.; McPhail, C. L.; Dikmans, A. J.; Vittal, J. J. *Tetrahedron Lett.* **1996**, 37, 1983–1986. DOI:[10.1016/0040-4039\(96\)00263-8](https://doi.org/10.1016/0040-4039(96)00263-8).
- [22] Bénard, C. P.; Geng, Z.; Heuft, M. A.; VanCrey, K.; Fallis, A. G. *J. Org. Chem.* **2007**, 72, 7229–7236. DOI:[10.1021/jo0709807](https://doi.org/10.1021/jo0709807).
- [23] Chow, T. J.; Wu, C.-C.; Chuang, T.-H.; Hsieh, H.-H.; Huang, H.-H.; Synthesis and applications of soluble pentacene precursors and related compounds. U.S. Patent 8,277,903 B2, 2012.
- [24] Nourmohammadian, F.; Salami, S. *J. Mol. Struct.: THEOCHEM* **2008**, 861, 147–148. DOI:[10.1016/j.theochem.2008.04.026](https://doi.org/10.1016/j.theochem.2008.04.026).
- [25] Fukuda, M.; Tajiri, A.; Oda, M.; Hatano, M. *Bull. Chem. Soc. Jpn.* **1983**, 56, 592–596. DOI:[10.1246/bcsj.56.592](https://doi.org/10.1246/bcsj.56.592).
- [26] Yu, Y.-X. *ACS Appl. Mater. Interfaces* **2014**, 6, 16267–16275. DOI:[10.1021/am504452a](https://doi.org/10.1021/am504452a).
- [27] Weidman, J. R.; Luo, S.; Zhang, Q.; Guo, R. *Ind. Eng. Chem. Res.* **2017**, 56, 1868–1879. DOI:[10.1021/acs.iecr.6b04946](https://doi.org/10.1021/acs.iecr.6b04946).
- [28] Navale, T. S.; Rathore, R. *Synthesis* **2012**, 44, 805–809. DOI:[10.1055/s-0031-1289695](https://doi.org/10.1055/s-0031-1289695).
- [29] Boldt, P.; Vardakis, F. *Angew. Chem., Int. Ed.* **1965**, 4, 1078–1078. DOI:[10.1002/anie.196510782](https://doi.org/10.1002/anie.196510782).
- [30] Fitzgerald, J. J.; Drysdale, N. E.; Olofson, R. A. *Synth. Commun.* **1992**, 22, 1807–1812. DOI:[10.1080/00397919208020501](https://doi.org/10.1080/00397919208020501).
- [31] Hashmat Ali, M.; Niedbalski, M.; Bohnert, G.; Bryant, D. *Synth. Commun.* **2006**, 36, 1751–1759. DOI:[10.1080/00397910600616859](https://doi.org/10.1080/00397910600616859).