

Preparation of $n\text{-Bu}_3\text{SnCH}_2\text{CH}=\text{CH}(\text{CH}_3)$ and $n\text{-Bu}_2(\text{Cl})\text{SnCH}_2\text{CH}=\text{CH}(\text{CH}_3)$ by Elimination Reaction of Alkoxides of the Type $n\text{-Bu}_2(\text{X})\text{Sn}-\text{O}-\text{C}(\text{CH}_3)(i\text{-C}_3\text{H}_7)\text{CH}(\text{CH}_3)\text{CH}=\text{CH}_2$ ($\text{X} = n\text{-Bu}$ and Cl)

VALERIO PERUZZO, G. TAGLIAVINI


Istituto di Chimica Analitica, Università di Padova, Via Marzolo, 1, I-35100, Padua, Italy

and ALESSANDRO GAMBARO

Istituto di Chimica Fisica, Università di Padova, Via Loredan, 1, I-35100, Padua, Italy



Received February 23, 1979

Recently it has been shown that di-*n*-butylallyltin chloride allows facile allylstannation of carbonyl compounds [1–3]. In addition it has been pointed out that organotin alkoxides of the type:



which are the products of the addition reaction, can reversibly form $n\text{-Bu}_2(\text{X})\text{SnCH}_2\text{CH}=\text{CH}_2$ by elimination of the unsaturated organic compound $\text{R}'\text{COR}''$ [2, 3].

As a consequence of these findings we should expect that organotin alkoxides II and III,



having an α -substituted allyl group joined to the tertiary carbon atom, could undergo an elimination reaction to give the corresponding ketone $\text{R}'\text{COR}''$ and the mixed 2-butenyl-butylns, $n\text{-Bu}_3\text{SnCH}_2\text{CH}=\text{CHR}'''$ and $n\text{-Bu}_2(\text{Cl})\text{SnCH}_2\text{CH}=\text{CHR}'''$, respectively. Thus we have prepared by transalkoxilation reaction between 2,3,4-trimethyl-5-hexen-3-ol and tri-*n*-butyltin- and di-*n*-butylchlorotin-methoxide the two alkoxides II and III having $\text{R}'' = (\text{CH}_3)_2\text{CH}$ and $\text{R}' = \text{R}''' = \text{CH}_3$. These, on heating, furnish the two mixed 2-butenyl-butylns, $n\text{-Bu}_3\text{SnCH}_2\text{CH}=\text{CHCH}_3$ (IV) and $n\text{-Bu}_2(\text{Cl})\text{SnCH}_2\text{CH}=\text{CHCH}_3$ (V) as *cis* and *trans*-isomer mixtures.


It is to be pointed out that this work deals with a simple route to new mixed substituted allyl-tins and shows the organometallic synthetic rôle of the found 'reversible allylstannation'.

Experimental

Preparation of 2,3,4-Trimethyl-5-hexen-3-ol

This compound was prepared by reacting crotyl bromide (100 g, 0.75 mol) and magnesium turnings (27.3 g, 1.12 mol) under reflux of diethyl ether as previously described [4]. The obtained crude liquid distilled twice under reduced pressure gave a final sample of 20.6 g (26% yield) of the title compound, b.p. 69–70 °C/16 mm (Lit., 63 °C/15 mm [4]).

The IR spectrum registered as a thin film (KBr optics) shows the following relevant bands: ν_{OH} at 3480 cm^{-1} and $\nu_{\text{C}=\text{C}}$ at 1635 cm^{-1} . The ^1H NMR spectrum of the carbinol,



has been registered by using a Bruker WH 90 operating in FT mode using TMS as internal standard. It clearly shows an ABC pattern in the range 6.02–4.91 ppm typical of the terminal ${}^5\text{CH}={}^6\text{CH}_2$ group, a multiplet centered at 2.39 ppm of the proton in the position 4, and another multiplet in the range 1.92–1.65 ppm for the *i*-propyl proton 2. The singlet of the OH proton is centered at 1.25 ppm, while all CH_3 protons resonate in the interval 1.14–0.84 ppm.

*Preparation of 2-Butenyl-tri-*n*-butyltin from Tri-*n*-butyltin Methoxide and 2,3,4-Trimethyl-5-hexen-3-ol*

In a three necked flask (50 ml) equipped with condenser, thermometer and separating funnel, 9.49 g (29.5 mmol) of $n\text{-Bu}_3\text{SnOMe}$ were mixed with 4.2 g (29.5 mmol) of the carbinol under stirring. The temperature was raised to 190–200 °C. Over 22 hours a mixture of methanol and 2-methyl-buten-3-one (2.2 g) was condensed and collected. The residue in the reaction flask was distilled under reduced pressure to give 6.5 g (63% yield) of $n\text{-Bu}_3\text{SnCH}_2\text{CH}=\text{CH}(\text{CH}_3)$ boiling at 98 °C/0.1 mm.

The IR spectrum registered on a thin film (KBr optics) shows two medium bands centered at 1655 and 1640 cm^{-1} attributable to two $\text{C}=\text{C}$ stretching vibrations dealing with a *cis*- and *trans*-isomeric mixture. The presence of *cis*-2-butenyltributyltin together with *trans*-2-butenyltributyltin has been confirmed by both ^1H and ^{13}C NMR spectra registered in CDCl_2 using TMS as internal standard.




Fig. 1. Schematic diagram of the proton decoupled ^{13}C NMR spectra of $\text{CH}_3\text{-(2)CH}=\text{(3)CH-CH}_2\text{SnBu}_2\text{X}$ (a, X = Cl; b, X = Bu) in the range of the olefinic carbon resonance.

behaviour may be ascribed to the electron withdrawing ability of the chlorine substituent which can modify the charge distribution of the double bond without affecting the geometrical arrangement of the olefinic moiety. The constant separation of the couple peaks of the *cis*- and *trans*-isomer in both compounds supports this view.

These results are promising for future development to new allyl-substituted tin compounds and work is now in progress in this laboratory.

Acknowledgments

We thank the CNR, Rome, for financial support and Mr D. Marton for technical assistance.

References

- 1 G. Tagliavini, V. Peruzzo, G. Plazzogna and D. Marton, *Inorg. Chim. Acta*, **24**, L47 (1977).
- 2 G. Tagliavini, V. Peruzzo and D. Marton, *Inorg. Chim. Acta*, **26**, L41 (1978).
- 3 V. Peruzzo and G. Tagliavini, *J. Organometal. Chem.*, **162**, 37 (1978).
- 4 P. Miginiac, *Bull. Soc. Chim. France*, 1077 (1970).
- 5 J. A. Verdone, J. A. Mangravite, N. M. Scarpa and H. G. Kuivila, *J. Am. Chem. Soc.*, **97**, 843 (1975).
- 6 A. G. Davies and P. G. Harrison, *J. Chem. Soc.*, 298 (1976).
- 7 J. A. Mangravite, J. A. Verdone and H. G. Kuivila, *J. Organometal. Chem.*, **104**, 303 (1976).
- 8 G. E. Maciel, P. D. Ellis, J. J. Matterstad and G. B. Savitsky, *J. Magn. Res.*, **1**, 589 (1969).
- 9 E. Lippmaa, S. Rang, O. Eisen and T. Pehk, *Eesti NVS. Tead. Akad. Toim. Keem. Geol.*, **16**, 351 (1967).
- 10 S. Rang, E. Lippmaa, T. Pehk and O. Eisen, *Eesti NSV Tead. Akad. Toim. Keem. Geol.*, **17**, 294 (1968).