[1959]

Canavan and Eaborn.

3751

Part XXI.* Organosilicon Compounds. 759. Some Compounds Containing Phosphorus.

By A. E. CANAVAN and C. EABORN.

Organosilicon compounds containing phosphorus have been prepared, including phosphonates [e.g., PhMe₂Si·CH₂·P(O)(OBuⁿ)₂], phosphine oxides $[e.g., (p-MeC_6H_4Me_2\cdot Si\cdot CH_2)_3PO]$, phosphates $[e.g., (p-Me_3Si\cdot C_6H_4\cdot O)_3PO]$, and a phosphinic acid $(p-Cl\cdot C_6H_4Me_2\cdot Si\cdot CH_2)_2P(O)\cdot OH$.

INTEREST attaches to organosilicon compounds containing phosphorus because of the possibility of modifying the properties of Silicones (particularly lubricating power and flameresistance) by introduction of phosphorus-containing groups. We have prepared several compounds containing silicon attached to phosphorus through carbon or carbon and oxygen. A few such compounds have been reported previously.¹⁻⁵

Interaction of the appropriate organosilylmethyl halides with sodium salts of the type $NaP(O)(OR)_2$ has been used to give the phosphonates $Me_3Si \cdot CH_2 \cdot P(O)(OR)_2$ (R = $n-C_4H_9$, $n-C_5H_{11}$), $ArMe_2Si-CH_2\cdot P(O)(OBu^n)_2$ ($Ar = Ph, p-Me-C_6H_4$, and $p-Cl-C_6H_4$), and $Me_3Si \cdot O \cdot Si Me_2 \cdot CH_2 \cdot P(O)(OBu^n)_2$. The products are liquids, stable enough to be fractionally distilled at reduced pressure.

The solid phosphine oxides $(Me_3Si \cdot CH_2)_3PO$ and $(p - Me \cdot C_6H_4Me_2 \cdot Si \cdot CH_2)_3PO$ have been prepared from the appropriate Si CH2 MgCl compound and phosphorus oxychloride in ether. The same reaction involving the Grignard reagent p-Cl·C₆H₄Me₂·Si·CH₂·MgCl gave only the phosphinic acid, $(p-Cl \cdot C_6 H_4 Me_2 \cdot Si \cdot CH_2)_2 P(O) \cdot OH$.

The phosphates (Me₃Si·CH₂·O)₃PO (a liquid stable to distillation at reduced pressure) and $(p-Me_3Si+C_6H_4)O_3PO$ (a solid) were prepared from phosphorus oxychloride and the

* Part XX, J., 1957, 955.

¹ Frisch and Lyons, J. Amer. Chem. Soc., 1953, 75, 4078.

² Gilbert and Precopio, Abs. Papers, 125th Meeting of the American Chemical Society, 1954, p. 16N; Gilbert, U.S.P. 2,768,193/1956; Chem. Abs., 1957, 51, 5816.
³ Keeber and Post, J. Org. Chem., 1956, 21, 509.
⁴ Chernyshev and Petrov, Doklady Akad. Nauk S.S.S.R., 1955, 105, 282; Chem. Abs., 1956, 50,

11,283.

⁵ Seyferth, J. Amer. Chem. Soc., 1958, 80, 1136. 6 G

sodium salts of trimethylsilylmethanol and p-trimethylsilylphenol in ether. Use of the latter salt with dimethylphosphorochloridate similarly gave the compound p-Me₃Si·C₆H₄·OP·(O)(OMe)₂.

Tristrimethylsilylmethylphosphine oxide decomposes in nitrogen at 270°, hexamethyldisiloxane probably being a product. Di-n-butyl trimethylsilylmethylphosphonate decomposes with polymerization on boiling at atmospheric pressure.

EXPERIMENTAL

General.—In fractional distillations special precision-made Vigreux columns (ca. 15 theoretical plates) were used.

(Aryldimethylsilyl)methyl Chlorides.—These were made by boiling the appropriate ArMgBr compounds with chloro(chloromethyl)dimethylsilane in ether for 12 hr., and had the following properties: (Ar =) Ph, b. p. 134—137°/48 mm.; p-Me·C₆H₄, b. p. 120—122°/12 mm., 136—138°/26 mm. (previously ⁶ reported incorrectly as 138—140°/12 mm.); p-Cl·C₆H₄, b. p. 102—104°/3 mm. (previously ⁶ reported incorrectly as 133°/3 mm.), $n_{\rm D}^{20}$ 1·5342.

Phosphonates.—Bromomethyltrimethylsilane (1 mol.) was boiled for 12 hr. with a light petroleum (b. p. 60—80°) solution of sodium (1 mol.) in dialkyl hydrogen phosphite (1 mol.). Washing with water followed by fractionation gave di-n-butyl trimethylsilylmethylphosphonate 3 (25%), b. p. 154—158°/17 mm., or *di-n-pentyl trimethylsilylmethylphosphonate* (36%), b. p. 186—188°/18 mm., $n_{\rm D}^{20}$ 1·4418 (Found: C, 55·0; H, 11·0. $C_{14}H_{33}O_3$ PSi requires C, 54·5; H, 10·8%). Similarly from (chloromethyl)pentamethyldisiloxane, with 23 hr. of refluxing (but with filtration to remove sodium chloride instead of washing), was obtained *di-n-butyl pentamethyldisiloxanylmethylphosphonate* (23%), b. p. 150—152°/10 mm., $n_{\rm D}^{20}$ 1·4251 (Found: C, 46·9; H, 9·7. $C_{14}H_{35}O_4$ PSi₂ requires C, 47·4; H, 9·9%).

Similarly from aryl(chloromethyl)dimethylsilanes were prepared di-n-butyl aryldimethylsilylmethylphosphonates as follows: (aryl =) *phenyl* (40%), b. p. 200–202°/10 mm., $n_{\rm p}^{20}$ 1.4864 (Found: C, 59.6; H, 9.5. C₁₇H₃₁O₃PSi requires C, 59.6; H, 9.1%); p-chlorophenyl (11%), b. p. 186.5–188°/2 mm., $n_{\rm p}^{20}$ 1.4942 (Found: C, 54.5; H, 8.4. C₁₇H₃₀O₃ClPSi requires C, 54.2; H, 8.0%); p-tolyl (32%, in this case refluxing was for 36 hr.), b. p. 187–189°/2.6 mm., $n_{\rm p}^{20}$ 1.4879 (Found: C, 60.3; H, 9.2. C₁₈H₃₃O₃PSi requires C, 60.6; H, 9.3%).

Diethyl trimethylsilylmethylphosphonate,² b. p. 118·5—121°/22 mm., was obtained in 25% yield by refluxing together equimolecular quantities of bromomethyltrimethylsilane and triethyl phosphite until the temperature rose from 96° to 120° (5 days), and then fractionating. Diethyl pentamethyldisiloxanylmethylphosphonate ² (41%), b. p. 154—156°/36 mm., was prepared similarly from chloromethylpentamethyldisiloxane by refluxing for 42 hr., during which the temperature of the mixture rose from 153° to 195°.

Phosphine Oxides.—Phosphorus oxychloride (14 g., 0.092 mole) in ether (50 ml.) was added slowly, with cooling to -5° , to the Grignard reagent from magnesium (6.7 g., 0.28 g.-atom) and bromomethyltrimethylsilane (4.6 g., 0.27 mole) in ether (150 ml.), and the mixture was boiled for 13 hr. After the usual hydrolysis procedure the ether layer was evaporated to dryness and the residue was recrystallised from ethanol, to give tristrimethylsilylmethylphosphine oxide (14.5 g., 51%), m. p. 182° (Found: C, 47.0; H, 10.9. Calc. for C₁₂H₃₃OPSi₃: C, 46.7; H, 10.8%). (This compound was recently prepared by a different route.⁵)

Similarly from chloromethyldimethyl-p-tolylsilane and p-bromophenyltrimethylsilane, respectively, but with 18 hours' refluxing, were obtained tris(dimethyl-p-tolylsilylmethyl)phosphine oxide, m. p. 111.5° (from ethanol) (Found: C, 67.3; H, 8.6. $C_{30}H_{45}$ OPSi₃ requires C, 67.1; H, 8.45%), and tris-(p-trimethylsilylphenyl)phosphine oxide, m. p. 259° (from ethanol) (cf. ref. 1). A similar procedure involving chloromethyl-p-chlorophenyldimethylsilane gave only bis-(p-chlorophenyldimethylsilylmethyl)phosphinic acid, m. p. 95° (from ethanol) (Found: C, 50.4; H, 5.9. $C_{18}H_{25}O_2Cl_2PSi_2$ requires C, 50.1; H, 5.85%).

An attempt to prepare tris(pentamethyldisiloxanylmethyl)phosphine oxide by this method was unsuccessful, an unidentified solid of m. p. 140° being obtained (Found: C, 29.0; H, 10.2%).

Phosphates.—Sodium (3.5 g., 0.15 g.-atom.) was dissolved in trimethylsilylmethanol (15 g., 0.15 mole) in ether (100 ml.), phosphorus oxychloride (7.7 g., 0.05 mole) in ether (50 ml.) was added, and the mixture was boiled for 16 hr. After filtration, the residue was distilled and the

⁶ Eaborn and Jeffrey, J., 1954, 4266.

Published on 01 January 1959. Downloaded by University of California - San Diego on 06/04/2017 12:14:45

distillate fractionated to give tristrimethylsilylmethyl phosphate (3.2 g., 18%), b. p. 134–136°/5 mm., $n_{\rm p}^{20}$ 1.4331 (Found: C, 40.7; H, 9.4. $C_{12}H_{33}O_4PSi_3$ requires C, 40.4; H, 9.3%).

Similarly from *p*-trimethylsilylphenol (but without distillation of the product) was obtained tris-p-trimethylsilylphenyl phosphate (64%), m. p. 99° (from ethanol) (Found: C, 59·8; H, 7·4. $C_{27}H_{39}O_4PSi_3$ requires C, 59·7; H, 7·25%), and from *p*-trimethylsilylphenol (0·1 mole) and dimethyl phosphorochloridate (0·1 mole) was obtained dimethyl p-trimethylsilylphenyl phosphate (44%), b. p. 152—154°/4 mm., n_p^{20} 1·4862 (Found: C, 47·8; H, 6·9. $C_{11}H_{19}O_4PSi$ requires C, 48·1; H, 7·0%).

Thermal Decomposition.—When tristrimethylsilylmethylphosphine oxide was refluxed in a slow stream of nitrogen, with continuous removal of volatile material from the top of the condenser, the temperature of the boiling liquid fell from 270° to 220° during 3 hr., and a little liquid of b. p. 95—103°, $n_{\rm D}^{20}$ 1·3771, was collected. This gave silica on burning, and is believed to be hexamethyldisiloxane (b. p. 101°, $n_{\rm D}^{20}$ 1·3774).

When di-n-butyl trimethylsilylmethylphosphonate was boiled under similar conditions for 6 hr., a little liquid of b. p. $102-104^{\circ}$, $n_{\rm D}^{20}$ 1·387, was collected and the residue became dark brown. Some porous pot was added and the heating continued for 1 hr.; the residue was transformed into a brittle white solid insoluble in alcohol or hydrocarbons.

We thank Messrs. Albright and Wilson (Mfg.) Limited for the gift of several organophosphorus compounds.

THE UNIVERSITY, LEICESTER.

[Received, June 19th, 1959.]