

PRELIMINARY COMMUNICATION


AN INTRAMOLECULAR ADDITION TO A BENZYNE*: THE SYNTHESIS OF MONOSUBSTITUTED HEPTAFLUOROBIPHENYLENES

S.C. COHEN, D. MOORE, R. PRICE and A.G. MASSEY


Department of Chemistry, Queen Mary College, Mile End Road, London E1 (Great Britain)

(Received March 27th, 1968)


As a continuation of our studies² on the thermal decomposition of polyfluoro-aryllithium reagents we allowed 2,2'-dilithiooctafluorobiphenyl³ to decompose at room temperature in the presence of furan and obtained two isomeric difuran benzyne adducts:



However, if the 2,2'-dilithiooctafluorobiphenyl is held at 0° for 30 minutes in the absence of a trapping reagent, the loss of one mole of lithium fluoride appears to be followed by an intramolecular addition of the intact half of the molecule to the benzyne to give 1-lithioheptafluorobiphenylene:



which can be converted, via either hydrolysis or bromination, to the pale yellow compounds 1-hydro- or 1-bromo-heptafluorobiphenylene:



* Part XVIII of a series; for Part XVII see ref. 1.

The reaction is a remarkably clean one in that if sufficient time is allowed for the total decomposition of 2,2'-dilithiooctafluorobiphenyl (e.g. about 30 minutes for 1.5 g quantities) the only impurities in the biphenylenes are polymeric species of much lower volatility. The biphenylenes (II) and (III) can thus be sublimed from the reaction mixtures in a state of high purity.

The proton NMR spectrum of (II) is centred at 6.53 ppm (cf. biphenylene⁴) whilst the ¹⁹F NMR spectrum consists of the expected seven groups of lines. The mass spectrum of (II) shows (as the main points of interest) a parent ion at $m/e = 278$, an ion at $m/e = 277$, a doubly-charged parent ion at $m/e = 139$ and three metastable peaks corresponding to the loss of CF, CF₂ and CF₃ from the parent ion.

REFERENCES

- 1 S.C. Cohen and A.G. Massey, *Chem. Ind. (London)*, (1968) 252.
 - 2 D.E. Fenton, A.J. Park, D. Shaw and A.G. Massey, *J. Organometal. Chem.*, 2 (1964) 437; D.E. Fenton and A.G. Massey, *Tetrahedron*, 21 (1965) 3009; S.C. Cohen, D.E. Fenton, D. Shaw and A.G. Massey, *J. Organometal. Chem.*, 8 (1967) 1; A.J. Tomlinson and A.G. Massey, *J. Organometal. Chem.*, 8 (1967) 321; S.C. Cohen, A.J. Tomlinson, M.R. Wiles and A.G. Massey, *Chem. Ind. (London)*, (1967) 877; S.C. Cohen, A.J. Tomlinson, M.R. Wiles and A.G. Massey, *J. Organometal. Chem.*, 11 (1968) 385.
 - 3 S.C. Cohen and A.G. Massey, *J. Organometal. Chem.*, 10 (1967) 471
 - 4 G. Fraenkel, Y. Asahi, M.J. Mitchell and M.P. Cava, *Tetrahedron*, 20 (1964) 1179.
- J. Organometal. Chem.*, 12 (1968) P37-P38