Nucleotides

DOI: 10.1002/anie.200501591

Expeditious, Potentially Primordial, Aminoacylation of Nucleotides**

Jean-Philippe Biron, Alastair L. Parkes, Robert Pascal,* and John D. Sutherland*

In contemporary biochemistry, enzymatically synthesized, activated aminoacyl-tRNA esters (aa-tRNA) serve as substrates for coded peptide synthesis by ribosomes,^[1] and a major goal is to understand the evolutionary path through which this process arose. As a first step towards this goal, we have been looking to find a prebiotically plausible means for the aminoacylation of ribonucleotide 2'-/3'-hydroxy groups. A number of simple, potentially prebiotic, activated amino acid derivatives have been reported, most notably *N*-carboxyan-hydrides (NCAs).^[2] The case for NCAs in prebiotic evolution has recently been strengthened by the finding that they can be produced from amino acids through the action of the simple volcanic gas carbonyl sulfide.^[3]

We therefore decided to investigate the aminoacylation of nucleotides by NCAs. Initially we were concerned that the nucleobase amino groups of adenine and cytosine would prove more nucleophilic than the 2'-/3'-hydroxy groups; however, recent findings show that NCAs react with inorganic phosphate to give aminoacyl phosphates.^[4] This suggests that it might be possible to generate nucleotide aminoacyl esters by initial aminoacylation of the phosphate monoester, followed by intramolecular aminoacyl transfer. Nucleotide aminoacylation by way of intermediate carboxylic phosphoric anhydrides would be analogous to the chemistry of aminoacyl-tRNA synthetase (aa-RS) enzymes. These enzymes

[*] J.-P. Biron, Dr. R. Pascal Dynamique des Systèmes Biomoléculaires Complexes UMR 5073-CC 017 Département de Chimie, Université Montpellier 2 Place Eugène Bataillon, 34095 Montpellier (France) Fax: (+33) 46-763-1046 E-mail: rpascal@univ-montp2.fr Dr. A. L. Parkes, Prof. Dr. J. D. Sutherland School of Chemistry, The University of Manchester Oxford Road, Manchester M139PL (UK) Fax: (+44) 161–275–4939 E-mail: john.sutherland@manchester.ac.uk
[***] This work was carried out within working group D27/001/02 of the EU COST action "Prebiotic Chemistry and Early Evolution", and was funded by COST and the Engineering and Physical Sciences

Research Council. We thank Dr Jo Kirkpatrick (Oxford University) for the LC/ESI MS analysis and Jean-Pascal Gimeno for experimental assistance in the chemistry of the mixed anhydride **8**. We gratefully acknowledge Peter Strazewski (Université Claude Bernard, Lyon 1) for his helpful comments on the manuscript and Auguste Commeyras (Université Montpellier 2) for facilitating the initiation of this collaborative project.

Supporting information for this article is available on the WWW under http://www.angewandte.org or from the author.

generate 5'-aminoacyladenylates (5'-aa-AMP) **1** (A = adenine) through an initial nucleophilic attack of an amino acid carboxylate on ATP. The aa-RS then catalyzes the intermolecular aminoacyl transfer from **1** to the 2'/3'-terminus of a cognate tRNA to give adenosine-5'-monophosphate (5'-AMP) (**2**) and an equilibrating mixture of 3'- and 2'-aa-tRNAs **3** and **4** (Scheme 1).



Scheme 1. The biochemistry and chemistry of 5'-aa-AMP 1: biochemically (upper), 1 is used by aa-RS enzymes for the aminoacylation of cognate tRNA. In the absence of enzymes (lower), 1 can undergo intramolecular aminoacyl transfer and hydrolysis reactions.

When separated from the protective environment of an aa-RS, 5'-aa-AMPs are highly unstable and undergo hydrolysis and isomerization in aqueous solution.^[5,6] The isomerization involves a slow initial intramolecular aminoacyl transfer from the 5'-phosphate to the 3'-hydroxy group via an eight-membered transition state to give the 3'-aminoacyl ester **5**, followed by rapid equilibration of the latter with a 2'-aminoacyl ester. If it were possible to generate 5'-aa-AMP **1** from 5'-AMP **2** by reaction with an NCA, then this intramolecular transfer would hopefully result in the sought-after formation of aminoacyl esters.

We first investigated whether the aminoacylation of a simple model phosphate monoester with an NCA was possible. We chose methyl phosphate **6** as the model phosphate monoester and the valyl derivative **7** to allow comparison with the previous work on inorganic phosphate^[4] and because **7** can be easily prepared and stored (Scheme 2).^[7] In a general sense, the reaction of a phosphate monoester with any electrophile is faster at pH values above 7 when the phosphate is in its dianionic state, (p K_a for the monoanionic state $\approx 6-7$ (Supporting Information)); however, NCA hydrolysis/polymerization is also favored at high pH values.

Communications



Scheme 2. Aminoacylation of model phosphate monoester by Val-NCA 7.

To slow such degradative reactions while still allowing reaction of the NCA with the phosphate group of **6**, we conducted our experiments under very slightly acidic conditions. In this case, although **6** was predominantly monoanionic, some of the reactive dianionic form still existed. Using ¹H NMR spectroscopy for analysis, we observed a low-level conversion of **6** and **7** into L-valyl methyl phosphate **8** (not isolated, according to ¹H NMR integration ≈ 23 % based on **7**, see Supporting Information). The conversion was only transient, however, and **8** was gradually converted back into the starting methyl phosphate **6** along with valine and oligovaline.

These observations imply that the reaction of 6 and 7 to give 8 is an equilibrium that is rapidly established and lies in favor of 6 and 7. The equilibrium is eventually dissipated through the hydrolysis of 7 and 8 to valine, and the subsequent reaction of the valine with further 7 and 8 giving oligovaline. The existence of the proposed equilibrium was proven when we found that 8, as prepared by conventional synthesis, was almost completely converted into 6 and 7 in the presence of bicarbonate. Once again, hydrolysis/polymerization of the activated valine derivatives subsequently yielded the free amino acid and oligomers.

We next turned our attention to the reaction of nucleotides with Val-NCA 7. 5'-AMP 2 showed similar behavior to methyl phosphate 6, and we observed an equilibrium production of 5'-Val-AMP 1 (R = iPr). However, the equilibrium was dissipated by NCA hydrolysis/polymerisation before the rearrangement of 1 into 5 occurred (see Supporting Information).^[5,8] We next studied the reaction of other regioisomeric nucleotides with Val-NCA 7. We recognized that if a nucleoside 3'/2'-valyl phosphate 9/10 could be transiently formed from the corresponding 3'/2'-nucleotide 11/12 and 7, then intramolecular aminoacyl transfer to the 2'/3'-OH group, via a seven-membered transition state^[9] might now be faster than hydrolysis/polymerization of the NCA or the intermediate mixed anhydrides (Scheme 3). We reasoned, therefore, that even an unfavorable equilibrium situation might lead to subsequent intramolecular aminoacyl transfer, and that this transfer should then displace the equilibrium according to Le Chatelier's principle. There was a major concern with the 3'/2'-nucleotides, however, as we foresaw cyclization of the intermediate 3'/2'-aminoacyl phosphates 9/10 to give the 2',3'-cyclic phosphate 15 as a likely competing reaction (Scheme 3). Indeed, 2'/3'-phosphates can be cyclized with great ease, to 2',3'-cyclic phosphates via fivemembered transition states, by almost any form of phosphate activation.[10]



Scheme 3. Aminoacylation/cyclization of 3'- and 2'-nucleotides by Val-NCA 7.

Undeterred by this concern, we initially studied the reaction of cytidine-3'-monophosphate (3'-CMP) **11** (base (B) = cytosine (C)) with Val-NCA **7**, again under very slightly acidic conditions. After 55 min, no new signals consistent with **9** (B = C) were observed in the ¹H NMR spectrum. However, new signals were observed which were tentatively assigned to the aminoacyl-transfer product **13** (B = C) (Figure 1a)). In



Figure 1. ¹H NMR spectra (500 MHz, D_2O) of the products of the reaction of 3'-CMP **11** (B=C) with Val-NCA **7**. a) Spectrum acquired after 1 h. b) Spectrum acquired after spiking the sample with authentic **15** (B=C) (0.5 mg).

particular, a diagnostic triplet between $\delta = 5.4$ and 5.5 ppm was shown by COSY analysis to correspond to 2'-H of the new species, and the downfield shift of this signal was consistent with a 2'-ester. Although **13** was not isolated,^[11] integration of the ¹H NMR spectrum of the reaction mixture suggested that it had been formed in $\approx 9\%$ overall yield (Table 1).

Table 1: Yields of the products observed in the reactions of 3'/2'-nucleotides 11/12 with Val-NCA 7.

Nucleotide	t [min]	Starting material [%]	Valyl ester (13 or 14) [%]	Divalyl ester [%]	15 [%]
11 (B=C)	55	90.6	(13) 8.7	trace	0.7
11 $(B = A)$	30	82.2	(13) 13.8	3.9	trace
	90	79.5	(13) 13.4	5.6	1.5
12 (B = C)	45	94.9	(14) 3.0	0	2.1
12 (B=A)	90	99.0	(14) 0.5	0	0.5

As our experiment was initiated at a pD value close to the pKa of the monoanionic form of 11, the yield of 13, based upon the reactive dianionic form of 11, is probably considerably higher. This suggests that isomerization of the intermediate 9 into 13 is highly efficient relative to the fast backwards reaction of 9 with CO₂. After further time had passed (>1 h), ¹H NMR spectroscopic analysis suggested that 13 underwent very slow reversion to 11, presumably by a combination of hydrolysis and nucleophilic attack by the amino group of valine and valyl derivatives, including 13. The minor signals in the ¹H NMR spectrum (accompanying the signals of 13) in the reaction of 11 and 7 were shown, by spiking the reaction with an authentic sample, to be due to the 2',3'-cyclic phosphate **15** (B = C) (Figure 1b)). However, this anticipated by-product was formed in much lower amounts (13/15 > 10:1), and we have not been able to prove the structure of the second minor product definitively, but MS and HMBC data strongly suggest that it is the 2'-divalyl analogue of 13.

We next investigated the effect of the nucleobase on this remarkable conversion and examined the reaction of 3'-AMP 11 (B = A) with Val-NCA 7. Again, we could not detect the intermediate aminoacyl phosphate 9 (B = A) but observed the aminoacyl-transfer product 13 (B = A) in a maximal overall yield of $\approx 14\%$ after 30 min (Table 1). In this experiment the putative 2'-divalvl analogue of 13 was also formed in a greater quantity and, after 90 min, had accumulated to the level of ≈ 6 %. On the basis of the amount of the reactive dianionic starting material, the combined synthesis efficiency of 13 (B = A) and the divalyl analogue is again, probably considerably higher. Close examination of the ¹H NMR spectra revealed the presence of the 2',3'-cyclic phosphate 15 (B = A) in low yield (13/15 > 8:1). It thus appeared that the nucleobase had a relatively small effect on the aminoacylation of 3'-nucleotides with Val-NCA, and as such, we switched our attention to the isomeric 2'-nucleotides with the expectation of equally efficient aminoacylation.

Through the use of the same conditions employed for the 3'-isomer, we found that cytidine-2'-monophosphate (2'-CMP) **12** (B = C), in the presence of Val-NCA **7**, gave the 3'-ester **14** (B = C) in only \approx 3% yield after 45 min (Table 1). Not only was the aminoacyl-transfer product formed in low yield, but it was accompanied by a comparable (\approx 2%) amount of the 2',3'-cyclic phosphate **15** (B = C) (**14**/**15** < 2:1). The lower yield of the ester product and the higher yield of **15** suggest that aminoacyl transfer from **10** is less efficient than it is from the 3'-phosphoryl isomer **9**. This conclusion was

supported by experiments with 2'-AMP 12 (B = A) and Val-NCA 7 in which we found that both the 3'-ester 14 (B = A)and the cyclic phosphate 15 (B = A) were formed in only trace amounts. This marked contrast in aminoacylation behavior between 3'- and 2'-nucleotides was not anticipated. It now seems likely that the efficiency of the aminoacylation reaction depends on the pK_a value of the phosphate and 2'/3'-OH groups, and possibly on conformational effects such as the furanose ring pucker. The pK_a value of the phosphate presumably influences the efficiency of the intramolecular aminoacyl-transfer step through a correlation with the leaving-group ability. Although the absolute pK_a values reported in the literature for the individual nucleotides differ (Supporting Information),^[12-14] there is a consensus that the 3'phosphates are more acidic than the 2'-phosphates by approximately 0.2 pK_a units and this is consistent with the more efficient conversion of 11 into 13. Within the 3'phosphate series, the greater yield of 13 (B = A) relative to 13(B = C) is possibly related to the fact that aden-9-yl is a better stabilizer for the 2'-oxyanion than any other aglycons.^[15]

The results described herein indicate that uncoded aminoacylation of 3'-nucleotides by NCAs is a prebiotically plausible reaction. The chemical mechanism of this process suggests that it should also be possible for 3'-phosphoryl oligoribonucleotides to undergo aminoacylation by NCAs in the same way, but with potential assistance by the ribonucleic acid chain through stereocomplementary binding. It is therefore possible that ribozymes, capable of coded aminoacylation by NCAs might have evolved and could have been key intermediates in the emergence of translation.^[16] The use of 5'-aa-AMPs as intermediates in coded aminoacylation of RNA is most likely a later evolutionary development.

Received: May 10, 2005 Revised: June 27, 2005 Published online: September 27, 2005

Keywords: amino acids · cyclization · *N*-carboxyanhydrides · ribonucleotides · RNA

- [1] P. Schimmel, Annu. Rev. Biochem. 1987, 56, 125.
- [2] K. Wen, L. E. Orgel, Origins Life Evol. Biosphere 2001, 31, 241.
- [3] L. Leman, L. Orgel, M. R. Ghadiri, Science 2004 306, 283.
- [4] J.-P. Biron, R. Pascal, J. Am. Chem. Soc. 2004, 126, 9198.
- [5] N. S. M. D. Wickramasinghe, M. P. Staves, J. C. Lacey, *Biochemistry* 1991, 30, 2768.
- [6] N. S. M. D. Wickramasinghe, J. C. Lacey, Origins Life Evol. Biosphere 1992, 22, 361.
- [7] In preliminary work, we found that other NCAs (e.g. Ile-NCA) behave in a manner similar to 7 and we are now extending the study to NCAs that are most conveniently generated in situ in water (e.g. Gly-NCA).
- [8] J. C. Lacey, N. Senaratne, D. W. Mullins, Origins Life 1984, 15, 45.
- [9] D. H. Rammler, Y. Lapidot, H. G. Khorana, J. Am. Chem. Soc. 1963, 85, 1989.
- [10] L. E. Orgel, R. Lohrmann, Acc. Chem. Res. 1974, 7, 368.
- [11] Our early attempts to isolate 13 were thwarted by its instability in the presence of (oligo)valine at the pH value of the reaction and over the time period required for HPLC. The successful characterization of 13 by LC–MS (see the Supporting Informa-

Communications

tion) involved an acidic mobile phase, and we are now attempting preparative separations under similar conditions.

- [12] D. D. Perrin, Dissociation Constants of Organic Bases in Aqueous Solution, Butterworths, London, 1965.
- [13] S. C. Gupta, N. B. Islam, D. L. Whalen, H. Yagi, D. M. Jerina, J. Org. Chem. 1987, 52, 3812.
- [14] L. F. Cavalieri, J. Am. Chem. Soc. 1953, 75, 5268.
- [15] I. Velikyan, S. Acharya, A. Trifonova, A. Földesi, J. Chattopadhyaya, J. Am. Chem. Soc. 2001, 123, 2893.
- [16] V. Borsenberger, M. A. Crowe, J. Lehbauer, J. Raftery, M. Helliwell, K. Bhutia, T. Cox, J. D. Sutherland, *Chem. Biodiversity* **2004**, *1*, 203.