

The pyrrole- and furan-derived bisadducts (Table I) could be aromatized in various ways. For example, decamethylantracene

(11) The ratio of the two isomers of **24** depended on the precursor, being 62:38 from **22** and 71:29 from **23** (determined by NMR integration).

$$\begin{array}{ccc}
 \mathbf{1} & \xrightarrow[\mathbf{6, PhMe}]{\text{BuLi (1 equiv)}} & \mathbf{22 (95\%)} \\
 \downarrow \text{BuLi (1 equiv)} & & \downarrow \text{BuLi (1 equiv)} \\
 \text{2 (R = CH}_3\text{), PhMe} & & \text{2 (R = CH}_3\text{), PhMe} \\
 & & \text{(42\%)} \\
 \downarrow \text{BuLi (1 equiv)} & & \\
 \text{6, PhMe (86\%)} & & \\
 \mathbf{23 (88\%)} & & \mathbf{24}
 \end{array}$$

iv

(18) Hoffman, R. W. "Dehydrobenzene and Cycloalkynes", Academic Press: New York, 1967.

Received June 2, 1980


Considerable interest has been shown in the chemistry of tetrathiafulvalenes (TTF) since some of these π -donors react with

Table I. Experimental Data for 3, 4, 7, 9, and 11

compd	mp, °C	color (solvent) ^a	spectra ^b
3, R = CN	196 dec	magenta (CH ₃ CN)	(IR) 2210, 2205 (CN), 1675 (C=O)
3, R = CF ₃	146-147	yellow (hexane)	(IR) 1680, 1665 (C=O), 1610, 1280, 1155 (CF ₃); (¹⁹ F NMR) 49.6
4, R = CN	290 dec	black (CHCl ₃ , continuous extraction)	(IR) 2210 (CN)
4, R = CF ₃	233-235	orange (benzene)	(IR) 1600, 1260, 1140 (CF ₃)
7	252 dec	red (chlorobenzene)	(IR) 1605, 1270, 1165 (CF ₃)
9, R = CO ₂ CH ₃	204 dec	olive (nitrobenzene)	(IR) 1740, 1720, 1710 (ester), 1680 (C=O); (¹ H NMR) 3.76, 3.80
11, R = CO ₂ CH ₃	88-90	maroon (hexane)	(IR) 1725, 1710 (ester); (¹ H NMR) 2.32, 3.72


^a Recrystallization solvent. ^b IR: KBr, cm⁻¹. ¹⁹F NMR: CHCl₃, ppm relative to CCl₃F. ¹H NMR: ppm relative to Me₄Si.

acceptors to form the most conducting organic solids presently known.¹ The recent synthesis of 1,3,5,7-tetrathiapentalene-2,6-dione^{2,3} (1, thiapendione) provided the possibility of elaborating a wide variety of novel TTF derivatives through cross-coupling reactions⁴ with 1,3-dithioles (2) (eq 1).



The cross-coupling reaction depends on the nature of the R and X substituents in dithiole 2. Mono and bis capping of thiapen to give products 3 and 4, respectively, occurs when R = CO₂CH₃ or CF₃ and X = S and when R = CN and X = O. No cross-coupling was observed for R = alkyl and/or X = Se. In a typical reaction, a 2-4 mol excess of dithiole derivative to thiapendione was refluxed with excess trimethyl phosphite in dry benzene under N₂ for several hours. The products (3-6) were isolated by chromatography and by their solubility differences. Identification of the products was based on their spectral properties, mass spectral data (except for 4) and elemental analysis. Table I summarizes some of this data.⁵

Selective high-yield preparations of either 3 or 4 could be achieved by modifying the cross-coupling reaction conditions. At 60 °C in neat triethyl phosphite, thiapendione (1) does not undergo self-coupling to dithiapendione (6) whereas dithioles 2 readily self-couple to their respective TTF derivatives 5. Because of solubility differences, addition of dithiole to a 60 °C solution of thiapendione in triethyl phosphite selectively precipitates in good yield (50-75%, based on thiapen) only mono-capped product 3, when R = CO₂CH₃, and only bis-capped product 4, when R =



7

(1) For reviews, see: (a) A. F. Garito and A. J. Heeger, *Acc. Chem. Res.*, **7**, 232 (1974); (b) A. N. Bloch, D. O. Cowan, and T. O. Poehler in "Energy and Charge Transfer in Organic Semiconductors", K. Masuda and M. Silver, Eds., Plenum Press, New York, 1974; (c) A. J. Heeger and A. F. Garito in "Low Dimensional Cooperative Phenomena", H. J. Keller, Ed., Plenum Press, New York and London, 1975; (d) E. M. Engler, *CHEMTECH*, **6**, 274 (1976); (e) M. Narita and C. U. Pittman, Jr., *Synthesis*, 489 (1976).

(2) R. R. Schumaker and E. M. Engler, *J. Am. Chem. Soc.*, **99**, 5521 (1977).

(3) For an alternate synthesis of the thiapen ring system, see: W. P. Krug, A. N. Bloch, and D. O. Cowan, *J. Chem. Soc., Chem. Commun.*, 180 (1979).


(4) E. M. Engler, V. V. Patel, and R. R. Schumaker, *J. Chem. Soc., Chem. Commun.*, 516 (1979).

(5) For data on 3 and 4, R = CO₂CH₃, see ref 4.

(6) H. K. Spencer, M. V. Lakshminathan, M. P. Cava, and A. F. Garito, *J. Chem. Soc., Chem. Commun.*, 867 (1975).


CF₃ or CN. In refluxing triethyl phosphite, mono-capped product 3 (R = CF₃) self-couples to provide, in about 10% yield, the first example of a trisubstituted TTF derivative, bis(dithiolide)dithiapen (7).

Extension of the cross-coupling reaction to 1,3-diselenoles (8, R = CO₂CH₃) failed when X = Se and led to a rearranged



mono-capped product 9 when X = S. As noted earlier⁷ in the self-coupling of diselenole thiones, an interchange of a ring selenium with the carbonyl sulfur occurs to give a mono-capped product (9) containing only one selenium.

Mono-capped products can be converted in strong base (e.g., NaOMe) to the corresponding TTF dithiolate intermediates (10)



which can then be treated with methyl iodide to provide unsymmetrically substituted TTF derivatives (11, R = CO₂CH₃). Alternately, treatment of 10 with transition-metal salts yielded a wide variety of novel metal dithiolene derivatives (12) which incorporate TTF as conjugated ligands.⁸

(7) E. M. Engler, V. V. Patel, and R. R. Schumaker, *J. Chem. Soc., Chem. Commun.*, 835 (1977).

(8) For experimental details, see: N. Martinez Rivera, E. M. Engler, and R. R. Schumaker, *J. Chem. Soc., Chem. Commun.*, 184 (1979).

Robert R. Schumaker*

IBM Research Laboratory
San Jose, California 95193

Edward M. Engler

IBM Watson Research Center
Yorktown Heights, New York 10598

Received April 11, 1980

Chemistry of Exciplexes. 10. Role of Excimers in Photocycloadditions of Arenes to 1,3-Dienes

Sir:

Photoexcited anthracenes and their catacondensed homologues form excimers,¹ and many are known to form exciplexes with