

DIRECTIVE EFFECTS IN THE HYDROBORATION OF ISOMERIC METHOXY-2-HEXYNES

GEORGE W. KABALKA* and SUZANNE SLAYDEN

Department of Chemistry, University of Tennessee, Knoxville, Tennessee 37916 (U.S.A.)

(Received January 6th, 1975)

Summary

A series of isomeric methoxy-2-alkynes were hydroborated to determine the effect of the methoxy group on the direction of the hydroboration. The effect of the methoxy group is to direct the boron atom to the side of the triple bond nearest to it. The magnitude of the directive effect increases as the methoxy group approaches the site of unsaturation. However, steric interactions become significant when the substituent is adjacent to the triple bond.

Introduction

The hydroboration reaction has become increasingly important in recent years [1]. The importance of hydroboration as a synthetic technique is enhanced by the fact that a wide variety of functionalities are unaffected by the hydroborating reagents [2]. Functional substituents may, however, affect the regio-specificity of the hydroboration reaction and considerable attention has been directed toward studies of the hydroboration of functionally substituted olefins [3-5].

In contrast to the alkene studies, little is known concerning substituent effects in the hydroboration of substituted alkynes [6-9]. It would appear that inductive effects could operate in the hydroboration of alkynes as they do in the case of alkenes.





We wish to report the results of a systematic study of the substituent effects on the regiospecificity of the monohydroboration of alkynes.

Results

A series of isomeric methoxyhexynes was synthesized and hydroborated. The methoxy group was chosen because it exhibits a strong inductive effect and because alkoxyorganoboranes are not prone to elimination reactions [10]. Dicyclohexylborane was used as the hydroborating agent. Dialkylboranes react

cleanly with alkynes to yield vinyl dialkylboranes which may be readily oxidized to the corresponding ketones [11]. When borane ($\text{BH}_3 \cdot \text{THF}$) was employed as the hydroborating reagent, material balances were poor due to dihydroboration [12], and consequently the observed ketone isomer distribution might not reflect the actual distribution of boron in the intermediates*.

The compounds hydroborated were the 6- (I), 5- (II), 4- (III), and 1-methoxy-2-hexyne (IV).



The resultant organoboranes were oxidized to the corresponding ketones and the products analyzed by NMR and GLPC. The overall sequence is illustrated for 6-methoxy-2-hexyne (Scheme 1). The results are summarized in Table 1.

SCHEME 1




TABLE I

THE HYDROBORATION OF A SERIES OF METHOXY-2-HEXYNES WITH DICYCLOHEXYLBORANE

Alkyne	Yield of ketones ^a (%)	2-Ketone ^{a,b} (%)	3-Ketone ^{a,b} (%)
2-Hexyne	98	67	33
6-Methoxy-2-hexyne	96	61	39
5-Methoxy-2-hexyne	98	39	61
4-Methoxy-2-hexyne	99	72	28
1-Methoxy-2-hexyne	85	80	20

^a Yields and percentages were determined by NMR and GLPC analyses using appropriate internal standards.

^b Products were isolated by preparative GLPC. They were characterized as the corresponding 2,4-dinitrophenylhydrazones. All exhibited satisfactory spectral and elemental analyses.

* It is interesting to note that we observed ketone isomer distributions for $\text{BH}_3 \cdot \text{THF}$ hydroborations parallel to those for dicyclohexylborane, except in hydroboration of 4-methoxy-2-hexyne where steric interactions for dicyclohexylborane would be expected to be larger than for borane itself.

Experimental

Reagents. 1-Pentyn-5-ol, 2-hexyn-1-ol, and 2-hexyn-5-ol were purchased from Farchan Chem. Co. and used as received.

Analyses. NMR analyses were performed on a Varian Associates Model A-60 spectrometer. Analytical GLPC analyses were performed on Varian Aerograph 1700 and 90-P instruments using a 10' column, 6% Carbowax on Chromosorb W. Preparative GLPC was performed on Varian Aerograph 711 using a 20' column, 30% FFAP on Chromosorb W. Elemental analyses were performed by Galbraith Laboratories, Knoxville, Tennessee and Microanalytical Laboratories, Oxford, England.

Alkynes

6-Methoxy-2-hexyne (I). The compound was prepared by the reaction of two equivalents of sodamide with 1-pentyn-5-ol, followed by addition of two equivalents of methyl iodide [13]. Yield, 31 g (0.28 mol, 55%); b.p. 146°C/739 Torr. NMR (neat): δ 3.17 (2H, t, $J = 6$ Hz, $-\text{CH}_2-\text{OCH}_3$), 3.07 (3H, s, $\text{CH}_3\text{O}-$), 1.9 (2H, m, $-\text{C}\equiv\text{C}-\text{CH}_2-$), 1.52 (3H, t, $J = 2.5$ Hz, $\text{CH}_3\text{C}\equiv\text{C}-$), 1.45 (2H, m, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$). Anal. Found: C, 74.76; H, 10.69. $\text{C}_7\text{H}_{12}\text{O}$ calcd.: C, 74.94; H, 10.80%.

5-Methoxy-2-hexyne (II). The compound was prepared from 2-hexyn-5-ol 49 g (0.5 mol) and methylated by the method of Brandsma [14]. Yield, 44 g (0.39 mol, 78%); b.p. 135-136°C/740 Torr. NMR (CCl_4): δ 3.11 (1H, m, HC), 3.10 (3H, s, $\text{CH}_3\text{O}-$), 2.05 (2H, m, $-\text{C}\equiv\text{C}-\text{CH}_2-$), 1.57 (3H, t, $J = 2$ Hz, $\text{CH}_3-\text{C}\equiv\text{C}-$), 1.00 (3H, d, $J = 6$ Hz, $\text{CH}_3-\text{CH}-$). Anal. Found: C, 74.90; H, 10.71. $\text{C}_7\text{H}_{12}\text{O}$ calcd.: C, 74.94; H, 10.80%.

4-Methoxy-2-hexyne (III). 4-Hydroxy-2-hexyne was prepared according to published procedure [15]. The methyl ether derivative was then prepared by the method of Brandsma [14]. B.p. 124-125°C/740 Torr. NMR (CCl_4): δ 3.67 (1H, m, $-\text{C}\equiv\text{C}-\text{CH}$), 3.24 (3H, s, $\text{CH}_3\text{O}-$), 1.65 (3H, d, $J = 2.5$ Hz, $\text{CH}_3\text{C}\equiv\text{C}-$), 1.47 (2H, m, CH_3-CH_2-), 0.87 (3H, t, $J = 7$ Hz, CH_2CH_3). Anal. Found: C, 74.86; H, 10.88. $\text{C}_7\text{H}_{12}\text{O}$ calcd.: C, 74.94; H, 10.80%.

1-Methoxy-2-hexyne (IV). The compound was prepared from 2-hexyn-1-ol by the method of Brandsma [14]. Yield, 48 g (0.43 mol, 86%); b.p. 139.5-140.5°C/740 Torr. NMR (CCl_4): δ 3.82 (2H, t, $J = 2$ Hz, $\text{CH}_3\text{O}-\text{CH}_2-$), 3.08 (3H, s, $\text{CH}_3\text{O}-$), 1.95 (2H, m, $-\text{C}\equiv\text{C}-\text{CH}_2\text{CH}_2-$), 1.3 (2H, m, CH_3CH_2-), 0.77 (3H, t, $J = 6.5$ Hz, CH_2CH_3). Anal. Found: C, 74.78; H, 10.82. $\text{C}_7\text{H}_{12}\text{O}$ calcd.: C, 74.94; H, 10.80%.

Monohydroboration of methoxyalkynes

The following procedure for the hydroboration of the alkynes with dicyclohexylborane is representative. To a suspension of dicyclohexylborane (11 mmol) in tetrahydrofuran (THF) was added at -10.0°C the appropriate alkyne (10 mmol). The reaction mixture was stirred at -10.0°C for 1 h and at room temperature for 1 h. The resulting vinylborane was oxidized by adding 3.65 ml of 3 N sodium hydroxide and 3.30 ml of 30% hydrogen peroxide to the reaction mixture. The cyclohexanol and ketone products formed were extracted with diethyl ether and dried over anhydrous magnesium sulfate. The distribution of

ketones was determined by GLPC (except IV). The yield of ketone was determined by NMR using benzene as an internal reference. The ketones were isolated by preparative GLPC.

Products from the hydroboration of 6-methoxy-2-hexyne (I)

6-Methoxyhexan-2-one. NMR (CCl_4): δ 3.22 (2H, t, $J = 5.5$ Hz, CH_3OCH_2-), 3.17 (3H, s, $\text{CH}_3\text{O}-$), 2.3 (2H, t, $J = 6.5$ Hz, $-\text{CH}_2\text{C}=\text{O}$), 1.97 (3H, s, $\text{CH}_3\text{C}=\text{O}$), 1.42 (4H, m, $-\text{CH}_2-\text{CH}_2\text{CH}_2-\text{CH}_2-$); 2,4-DNPH m.p. 68-69°C [lit. [16] 60-70°].

6-Methoxyhexan-3-one. NMR (CCl_4): δ 3.27 (2H, t, $J = 5.5$ Hz, CH_3OCH_2-), 3.21 (3H, s, $\text{CH}_3\text{O}-$), 2.57-2.10 (4H, m, $-\text{CH}_2\text{COCH}_2-$), 1.68 (2H, m, $\text{CH}_2-\text{CH}_2-\text{CH}_2-$), 0.95 (3H, t, $J = 7.5$ Hz, CH_3CH_2-). The pure 2,4-DNPH derivative of the ketone could not be obtained. An analysis of the isomeric ketone derivative mixture and the composite NMR spectrum are consistent with this product [16]. Anal. Found: C, 50.16; H, 5.77; N, 18.06. $\text{C}_{13}\text{H}_{18}\text{N}_4\text{O}_5$ calcd.: C, 50.31; H, 5.86; N, 18.05%.

Products from the hydroboration of 5-methoxy-2-hexyne (II)

5-Methoxyhexan-2-one. NMR (CCl_4): δ 3.22 (1H, m, HC), 3.22 (3H, s, $\text{CH}_3\text{O}-$), 2.4 (2H, t, $J = 7$ Hz, $-\text{CH}_2\text{C}=\text{O}$), 2.05 (3H, s, $\text{CH}_3\text{C}=\text{O}$), 1.6 (2H, m, $\text{CH}_2-\text{CH}_2-\text{CH}$), 1.05 (3H, d, $J = 6$ Hz, CH_3CH); 2,4-DNPH m.p. 67-68°C. Anal. Found: C, 49.99; H, 5.77; N, 18.33. $\text{C}_{13}\text{H}_{18}\text{N}_4\text{O}_5$ calcd.: C, 50.31; H, 5.86; N, 18.05%.

5-Methoxyhexan-3-one. NMR (CCl_4): δ 3.68 (1H, m, $\text{HC}-\text{OCH}_3$), 3.23 (3H, s, $\text{CH}_3\text{O}-$), 2.41 (4H, m, $-\text{CH}_2\text{COCH}_2-$), 1.07 (3H, d, $J = 6$ Hz, CH_3CH), 0.97 (3H, t, $J = 7.5$ Hz, CH_3CH_2-); 2,4-DNPH derivative isolated as the α,β -unsaturated hydrazone [17], m.p. 112-112.5°C [lit. [18] m.p. 121°]. NMR (CDCl_3): δ 6.63-6.27 (2H, vinyl), 2.62 (2H, q, $J = 7.5$ Hz, $-\text{CH}_2-\text{C}=\text{N}$), 2.0 (3H, m, $\text{CH}_3-\text{CH}=\text{CH}-$), 1.23 (3H, t, $J = 7.5$ Hz, CH_3CH_2-).

Products from the hydroboration of 4-methoxy-2-hexyne (III)

4-Methoxyhexan-2-one. NMR (CCl_4): δ 3.4 (1H, m, HC), 3.13 (3H, s, $\text{CH}_3\text{O}-$), 2.34 (2H, m, $-\text{CH}_2\text{C}=\text{O}$), 1.97 (3H, s, $\text{CH}_3\text{C}=\text{O}$), 1.33 (2H, m, CH_3-CH_2-), 0.72 (3H, t, $J = 6.5$ Hz, CH_3CH_2-). 2,4-DNPH derivative was isolated as the α,β -unsaturated hydrazone [17], m.p. 166-167°C [lit. [19] m.p. 162-163.5°]. NMR (CDCl_3): δ 6.48 (2H, vinyl), 2.65-2.05 (2H, m, $-\text{CH}_2-$), 2.22 (3H, s, $\text{CH}_3-\text{C}=\text{N}$), 1.15 (3H, t, $J = 7.5$ Hz, CH_3CH_2-).

4-Methoxyhexan-3-one. NMR (CCl_4): δ 3.33 (1H, t, $J = 6$ Hz, CH), 3.25 (3H, s, $\text{CH}_3\text{O}-$), 2.38 (2H, q, $J = 7$ Hz, $-\text{CH}_2-\text{CH}_3$), 1.52 (2H, m, $\text{CH}_3-\text{CH}_2-\text{CH}$), 0.9 (6H, m, 2 CH_3). The pure 2,4-DNPH- α,β -unsaturated hydrazone derivative could not be isolated. The NMR spectrum of the isomeric hydrazone mixture was consistent with this product.

Products from the hydroboration of 1-methoxy-2-hexyne (IV)

1-Methoxyhexan-2-one. NMR (CCl_4) δ 3.8 (2H, s, CH_3OCH_2-), 3.35 (3H, s, $\text{CH}_3\text{O}-$), 2.37 (~2H, m, CH_2CO), 1.33 (~4H, m, $\text{CH}_3\text{CH}_2\text{CH}_2-$), 0.83 (~3H, CH_3-CH_2-); 2,4-DNPH m.p. 91-92°C [lit. [20] m.p. 93°]. NMR (CDCl_3): δ 4.36 (2H, s, CH_3OCH_2-), 3.60 (3H, s, $\text{CH}_3\text{O}-$), 2.36 (2H, t, $J = 7$ Hz, $-\text{CH}_2-\text{C}=\text{N}$), 1.53 (4H, m, $\text{CH}_3\text{CH}_2\text{CH}_2-$), 0.95 (3H, t, $J = 6$ Hz, CH_3-CH_2-).

1-Methoxyhexan-3-one was observed as an impurity in the 1-methoxyhexan-2-one material. Repeated attempts to isolate this material were fruitless. The isomers were inseparable by GLPC. The NMR spectrum of the 2-ketone indicated a maximum of 20% of the 3-ketone as an impurity. NMR (CCl_4): δ 3.5 (2H, t, $J = 6.5$ Hz, $\text{CH}_3\text{O}-\text{CH}_2-$), 3.22 (3H, s, $\text{CH}_3\text{O}-$), 2.37 ($\sim 4\text{H}$, $-\text{CH}_2\text{CO}-\text{CH}_2-$), 1.33 ($\sim 2\text{H}$, CH_3CH_2-), 0.83 ($\sim 3\text{H}$, CH_3CH_2-).

Acknowledgement

The authors wish to thank Research Corporation for support of this study.

References

- 1 H.C. Brown, *Boranes in Organic Chemistry*, Cornell University Press, 1972, p. 372.
- 2 H.C. Brown, *Hydroboration*, Benjamin, 1962, p. 265.
- 3 H.C. Brown and M.K. Unni, *J. Amer. Chem. Soc.*, **90** (1968) 2902.
- 4 H.C. Brown and R.M. Gallivan, *J. Amer. Chem. Soc.*, **90** (1968) 2906.
- 5 D.J. Pasto and J. Hickman, *J. Amer. Chem. Soc.*, **90** (1968) 4445.
- 6 (a) J. Plamondon, J.T. Snow and G. Zweifel, *Organometal. Chem. Syn.*, **1** (1971) 249; (b) G. Zweifel, A. Horng and J. Snow, *J. Amer. Chem. Soc.*, **92** (1970) 1427.
- 7 G. Zweifel, H. Arzoumanian and C.C. Whitney, *J. Amer. Chem. Soc.*, **89** (1967) 3652.
- 8 E. Negishi and T. Yoshida, *J. Amer. Chem. Soc.*, **95** (1973) 6837.
- 9 E. Negishi and T. Yoshida, *Chem. Commun.*, (1973) 606.
- 10 H.C. Brown and E. Knights, *J. Amer. Chem. Soc.*, **90** (1968) 4439.
- 11 G. Zweifel, G.M. Clark and N.L. Polston, *J. Amer. Chem. Soc.*, **93** (1971) 3395.
- 12 G. Zweifel and H. Arzoumanian, *J. Amer. Chem. Soc.*, **89** (1967) 291.
- 13 N. Rabjohn, *Coll. Org. Syn.*, **IV**, 117 (1963).
- 14 L. Brandsma, *Preparative Acetylenic Chemistry*, Elsevier, 1971, p. 172.
- 15 E.A. Braude and J.A. Coles, *J. Chem. Soc.*, (1951) 2083.
- 16 R.C. Elderfield, B.M. Pitt and I. Wempen, *J. Amer. Chem. Soc.*, **72** (1950) 1334.
- 17 I.N. Nazarov and I.I. Zaretskaya, *Bull. Acad. Sci. URSS Classe Sci. Chem.*, (1942) 200.
- 18 H. Normant and G. Martin, *Bull. Soc. Chim. Fr.*, (1957) 429.
- 19 Zh.A. Krasnaya and V.F. Kuchеров, *Izv. Akad. Nauk SSSR Ser. Khim.*, **6** (1965) 1070.
- 20 S. Maruyama, *Sci. Papers Inst. Phys. Chem. Research (Tokyo)*, **20** (1933) 53.