among saturated chiral deuterium compounds.²⁹ We have observed a similarly large [α] for the corresponding dimethylamino compound, synthesized either by direct displacement on tosylate **8** or by reaction of **8** with NaN₃ followed by LiAlH₄ reduction and Eschweiler–Clarke methylation; because the two routes gave amine with the same [α], it is reasonable to conclude that the reaction of the tosylate with a secondary amine, like the reaction with azide ion,^{29c,d} occurs with complete inversion of configuration.

- (29) (a) L. Verbit, *Prog. Phys. Org. Chem.*, 7, 51 (1970); (b) D. Arigoni and E. L. Eliel, *Top. Stereochem.*, 4, 127 (1969); (c) B. Stephenson, G. Sołladić, and H. S. Mosher, *J. Am. Chem. Soc.*, 94, 4184 (1972); (d) P. H. Anderson, B. Stephenson, and H. S. Mosher, *ibid.*, 96, 3171 (1974); (e) W. Meister, R. D. Guthrie, J. L. Maxwell, D. A. Jaeger, and D. J. Cram, *ibid.*, 91, 4452 (1969).
- (30) C. K. İngold, "Structure and Mechanism in Organic Chemistry", 2nd ed, Cornell University Press, Ithaca, N.Y., 1969, pp 853–861.

Ronald M. Magid,* O. Stanley Fruchey

Department of Chemistry, The University of Tennessee Knoxville, Tennessee 37916 Received September 9, 1977

A New Approach to the Synthesis of Large-Ring Phosphorus Compounds¹

Sir:

Very few monocyclic phosphorus compounds with ring size greater than six have been prepared, and virtually nothing is known about the stereochemical properties of the especially interesting systems where the ring contains about seven-nine members.² On the other hand, a great deal of work has been done on five- and six-membered rings, and research continues at a fast rate on these relatively accessible systems. In addition, studies focused on possible aromaticity in the five- and sixmembered rings are numerous, but, save for one case, where cyclic delocalization was considered for 1-phenylphosphepin 1-oxide,³ no attention has been given to delocalization phenomena among phosphorus compounds with larger rings. We have devised a simple synthetic approach for phosphorus compounds containing seven or more ring members that should lead to many new compounds permitting the exploration of stereochemical and delocalization phenomena. Our approach provides diketo derivatives of the ring systems, an extremely desirable feature since the keto functionality permits synthetic maneuverability at important ring positions.

Our method basically comprises two steps from a nonphosphorus-containing starting material. We first employ the family of 1,2-dimethylenecycloalkanes (1) as dienes in the



McCormack cycloaddition⁴ with phosphorus(III) halides, obtaining in good yield under conventional conditions (standing

for several days at room temperature) a series of bicyclic phospholene derivatives (2). These compounds have been fully characterized by analysis and by ¹H, ¹³C, and ³¹P NMR spectroscopy, details of which will be published elsewhere. The phospholene products are then subjected to ring-opening ozonolysis at -78 °C in methanol. Following removal of any excess ozone by nitrogen bubbling, trimethyl phosphite is added to the mixture to reduce the ozonide and the temperature allowed to reach that of the room. Evaporation of the solution leaves a crystalline residue of the diketones 3, which is broken up in ether for washing and collection on a filter.

We have so far used the dienes⁵ based on the cyclobutane, cyclopentane, and cyclohexane systems in this new synthesis, and from methylphosphonous dichloride have obtained derivatives of the phosphepane (4), phosphocane (5), and phos-



phonane (6) systems, respectively. These products are easily recrystallized (6, benzene; 4 and 5, acetonitrile) stable solids. They are mildly hygroscopic. Some important physical properties are given in Table I; all of the compounds gave correct C, H, and P analyses. The spectra do not reveal the presence of any enol form, in contrast to the behavior of a β -keto derivative of the five-membered phospholane system.⁶ However, the four α protons of 6 were found to be rapidly and completely exchanged when D₂O was added to a Me₂SO solution.

Phosphorus trihalides also have been used in our two-step process, giving diketophosphinic acids. Thus, starting with 1,2-dimethylenecyclohexane and PBr_3 we have formed phosphinic acids 7 and 8. Compound 8 had the intriguing property



of undergoing internal aldol condensation on storage as the solid at room temperature for several weeks. The same reaction occurred within 2.5 h simply on refluxing in benzene with water removal (Dean-Stark). The product is a new bicyclic compound 9, mp 150-152 °C, whose structure became evident



Table I. Properties of the 1-Substituted β , β' -Diketophosphacycloalkane 1-Oxides.

					¹³ C NMR, δ^b				
Compd	Yield, %	Mp, °C	$\nu_{\rm C=0}, {\rm cm}^{-1}$	³¹ P NMR, δ^a		C=0	Cγ	Cδ	P-CH ₃
4	90	182-184	1690¢	+29.8	49.2 (50.1)	202.8 (4.9)	38.5 <i>d</i>		16.2 (70.8)
5	83	204-207	1690 <i>°</i>	+29.0	45.3 (48.8)	204.5 (4.9)	45.6 ^d	19.0 <i>d</i>	14.7 (70.8)
6	94	130-132	1700 <i>°</i>	+32.0	46.1 (54.7)	206.0 (4.9)	44.5 <i>d</i>	23.4 ^d	19.2 (72.3)
8	84	102-104	1690 ^f	+26.5	47.4 (77.2)	205.8 (2.9)	42.5 <i>d</i>	23.0 ^d	

^{*a*} In Me₂SO-*d*₆; downfield from 85% H₃PO₄ as reference. Taken on a Bruker HFX-10 spectrometer with proton decoupling. ^{*b*} Compounds **4**, **5**, and **8** in Me₂SO-*d*₆; **6** in CDCl₃. Values in parentheses are ¹³C-³¹P coupling constants, in hertz. Taken on a JEOL FX-60 spectrometer with proton decoupling. ^{*c*} Nujol mull. ^{*d*} No ¹³C-³¹P coupling observed. ^{*e*} CHCl₃ solution. ^{*f*} KBr disk.

from analysis and spectral features for an α,β -unsaturated ketone (¹³C NMR δ 188.9 (C=O), 140.0 (=C_{α}), 154.7 $(=C_{\beta})$; IR $\nu_{C=O}$ 1650, $\nu_{C=C}$ 1610 cm⁻¹) in a monomeric system (single ³¹P signal at δ +46.8). The phosphine oxide counterpart of 8 (6) also could be made to undergo the internal aldol condensation, but required the presence of a trace of added acid (p-toluenesulfonic) catalyst in the refluxing benzene medium. The product (10) had mp 97-98 °C and similar spectral features to those of 9. These bicyclic compounds are members of a new series and may prove to have value as intermediates for construction of multicyclic phosphorus compounds, another area under intense development.⁷

References and Notes

- (1) Supported in part by Public Health Service Grant CA-05507 from the National Cancer Institute
- (2)M. J. Gallagher in "Stereochemistry of Heterocyclic Compounds", Vol. II, W. L. F. Armarego, Ed., Wiley-Interscience, New York, N.Y., 1977, Chapter
- G. Märkl and H. Schubert, Tetrahedron Lett., 1273 (1970)
- W. B. McCormack, U. S. Patents 2 663 636 and 2 663 637 (1953). This work has been reviewed: L. D. Quin in ''1,4-Cycloaddition Reactions'', J. Hamer, (4) Ed., Academic Press, New York, N.Y., 1967, Chapter 3
- The dienes were prepared by pyrolysis of the N,N'-dioxide of the bis(dimethylaminomethyl) derivatives using the procedure of A. C. Cope and E. Ciganek, Org. Syn., 39, 40 (1959), as a guide. The trans-dicarboxylic acids were the source of the amines by the sequence (1) SOCi₂, (2) Me₂NH, (3) LiAIH4. The commercially available anhydride of cis-cyclohexanedicarboxylic acid was converted to the dimethyl ester, epimerized by methanolic sodium methoxide, and hydrolyzed by aqueous KOH to the acid. Cyclobutane and cyclopentanedicarboxylic acids were obtained from diethyl α , α' -dibromoadipate and α, α' -dibromopimelate, respectively, by cyclization with Moalpate and *a*,*a* anothrophinetate, respectively, by cyclication man NaCN, followed by HCI hydrolysis (see R. C. Fuson and T. Y. Kao, *J. Am. Chem. Soc.*, **51**, 1536 (1929); R. C. Fuson and W. Cole, *ibid.*, **60**, 1237 (1938); R. C. Fuson, C. L. Fleming, P. F. Warfield, and D. E. Wolf, *J. Org. Chem.*, **10**, 121 (1945)). The dienes had the expected ¹H NMR spectra (P. D. Bartlett, A. S. Wingrove, and R. Owyang, J. Am. Chem. Soc., 90, 6067 (1968)).
- (6)
- L. D. Quin and R. C. Stocks, *J. Org. Chem.*, **39**, 686 (1974). S. D. Venkataramu, G. D. Macdonell, W. R. Purdum, M. El-Deek, and K. D. Berlin, Chem. Rev., 77, 121 (1977).

Louis D. Quin,* Eric D. Middlemas

Gross Chemical Laboratory, Duke University Durham, North Carolina 27706 Received August 29, 1977

A Comparison of Allyl and Nitrosyl **Coordination in the 18-Electron Complex** AllyInitrosylbis(triphenylphosphine)ruthenium and Its CO Adduct

Sir:

The dual nature of both allyl and nitrosyl ligands as 1e⁻ and 3e⁻ donors in transition metal complexes is well established,^{1,2} and recent attention has focused on the catalytic potential which this duality provides.³⁻⁶ Sorely lacking, however, are examples of stable, mixed allyl nitrosyl complexes, and the comparative studies of these ligands and their bonding propensities which such species will allow. To date, only two systems of this type have been reported,^{7,8} the most extensively studied being $Fe(NO)(\eta^3-C_3H_5)L_2$ where L = CO, tertiary phosphine.⁸ We wish to report the synthesis and crystal structure of the first platinum group metal complex of this type, and evidence for its reaction with CO to form a five-coordinate species containing a bent nitrosyl.

Reaction of RuCl(NO)(PPh₃)₂⁹ in benzene or THF with tetraallyltin at room temperature gives the red, air-sensitive $Ru(NO)(\eta^3-C_3H_5)(PPh_3)_2$, 1, in 70% yield. Recrystallization from THF/Et₂O affords analytically pure¹⁰ crystals which are stable in air for several weeks. The complex is soluble in C_6H_6 , THF, and CH_2Cl_2 , but decomposes rapidly in $CHCl_3$ and CCl_4 to form $RuCl_3(NO)(PPh_3)_2$. The IR spectrum of 1 (KBr) shows a nitrosyl stretch at 1620 cm⁻¹ (1640 cm⁻¹ in



Figure 1. ¹H NMR spectra of the allyl-nitrosyl complexes in benzene- d_6 : (A) $Ru(\eta^3-C_3H_5)(NO)(PPh_3)_2$; (B) computer simulation of spectrum A; (C) $Ru(\eta^3-C_3H_5)(NO)(CO)(PPh_3)_2$; (D) ³¹P-decoupled spectrum of the carbonyl adduct above; (E) computer simulation of spectrum C. Resonances marked with an asterisk (*) are due to traces of H₂O in solvent.

 CH_2Cl_2), while the ¹H NMR spectrum exhibits a pattern characteristic of essentially symmetric η^3 -allyl coordination with phosphorus coupling to the anti protons. The ¹H NMR spectrum of 1 is shown in Figure 1A and a computer simulation¹¹ is given in Figure 1B (central proton, δ 4.70; anti protons, 1.00; syn protons, 2.40 and 2.42 (${}^{3}J_{syn} = 4.0 \text{ Hz}$; ${}^{3}J_{anti} = 10.0$ $Hz; J_{PH_{anti}} = 6.0 Hz)).$

The coordination geometry of **1** and the mode of nitrosyl coordination were unambigously established by a single-crystal x-ray analysis. The complex crystallizes in the monoclinic space group $P2_1$ in a cell of dimensions a = 9.04(1), b = 17.47(3), $c = 11.51 (1) \text{ Å}; \beta = 115.39 (5)^{\circ} \text{ with } Z = 2. \text{ The structure}$ was solved by standard heavy-atom methods and refined by least-squares procedures to final agreement factors R and R'of 0.0299 and 0.0383 for 2074 reflections having $F_0 \ge 3\sigma(F_0)$ and 159 variables.¹² Tables of final structural parameters and intramolecular distances and angles are available as supplementary material (see paragraph at the end of the paper).

Figure 2 presents a perspective view of the structure of 1