

The azo acid obtained by either method above was reprecipitated from dilute bicarbonate solution before weighing. It melted over 10° ranges above 75°; further purification gave material whose melting point, neutralization equivalent and behavior on sublimation were identical with those described previously.² The diphenic acid obtained melted above 210°. Previous experience has shown that this material is at least 90% pure. Typical results are recorded in Table I.

TABLE I

Run	Moles, R ₁ /R ₂ ^a	Azobenzene, %	Azobenzene-2-carboxylic acid, %	Diphenic acid, %
1	1/1	24	16	17
2	1/1	25	15	29
3	3/1	32	40	9
4	3/1	32	47	14
5	3/1	32	34	11
6	1/3	10	23	38
7	1/3	12	36	46
8	1/3	6	38	45

^a R₁ = diazotized aniline, R₂ = diazotized anthranilic acid.

To demonstrate the absence of biphenyl-2-carboxylic acid in the products the acidic products from a typical run were dissolved in concentrated sulfuric acid and the solution heated to 140° for five minutes. Under these conditions any biphenyl-2-carboxylic acid present would be quantitatively converted to fluorenone. The solution was cooled, poured on ice, and the resulting precipitate examined. The non-acidic portion of the product consisted of ether-insoluble tars, from which no fluorenone was isolated; these tars were shown in a separate experiment to be derived from azobenzene-2-carboxylic acid. The acidic portion of the product consisted of fluorenone-4-carboxylic acid, derived from diphenic acid.

(2) Paal and Krecke, *Ber.*, **24**, 3060 (1891).

DEPARTMENT OF CHEMISTRY
UNIVERSITY OF NEW HAMPSHIRE
DURHAM, N. H.

RECEIVED AUGUST 24, 1949

An Improved Synthesis of 2,4-Diacetoxymercurianiline

BY THOMAS C. BRUCE

In the course of a study of aromatic mercury derivatives, it became desirable to prepare large quantities of 2,4-diacetoxymercurianiline. The synthesis of this compound has been previously reported by Vecchiotti in 24% yield.¹ On attempting to repeat this earlier work, the author was unable to obtain the dimercuri compound in yields above 16%. A careful study of the best conditions for obtaining the diacetoxymercurianiline showed that control of pH was a primary factor in permitting isolation of the product in high yield. In a series of runs, carried out in buffered solutions of varying pH, it was found that at pH 4.25 a maximum amount (93.5%) of diacetoxymercurianiline could be isolated; and that deviations of even one pH unit from this optimum condition resulted in very decided lowering of yields. The solution of the preparative problem in the case of aniline is considered important, since the same problem will no doubt be encountered in the

(1) L. Vecchiotti, *Gazz. chim. ital.*, **44**, II, 34-38 (1914).

mercuration of other aromatic amines with mercuric acetate.

Experimental

To 479 g. (7.99 moles) of glacial acetic acid, stirred at room temperature, there was added 251 g. (2.99 moles) of sodium bicarbonate. After the foaming had subsided, 3.2 liters of water was added in such a manner that foaming was kept to a minimum. The pH of the resulting solution was found to be 4.30. One mole (318.6 g.) of C. P. mercuric acetate was added, and the pH was then found to be 4.25. With constant stirring, there was then added one mole (93 g.) of aniline, and the mixture was set aside in the dark for forty-eight hours to permit complete precipitation of the product. If the pH of the solution was carefully adjusted, as above, it was found that washing the precipitate freely with several portions of hot water and then drying in a vacuum desiccator over sodium hydroxide led to 285.3 g. (93.5%) of an excellent product, which melted at 209°, with decomposition. The value recorded in the literature is 208°. The product was converted to 2,4-dibromoacetanilide (m.p. 146°) to confirm its identity.

*Anal.*² Calcd. for C₁₀H₁₁O₄NHg₂: C, 19.68; H, 1.82; Hg, 65.73. Found: C, 19.85; H, 1.88; Hg, 65.54.

If the adjustment of pH in the above preparation was not carefully made, or if different proportions of reagents than those described above were used, it was found that the yields of 2,4-diacetoxymercurianiline were sharply cut. With mercurating solutions of pH above 4.25 the product was contaminated with considerable proportions of *p*-acetoxymercurianiline, which could be removed to some extent by extracting with chloroform. At pH values below 4.25, the yield of the desired product was lowered by formation of the soluble 2,4-diacetoxymercurianilinium acetate.

(2) Analysis by Dr. A. Elek of the Elek Micro-analytical Laboratory.

DEPARTMENT OF CHEMISTRY
UNIVERSITY OF SOUTHERN CALIFORNIA
LOS ANGELES 7, CALIFORNIA RECEIVED AUGUST 27, 1949

X-Ray Diffraction Studies of the System: Zn₂TiO₄-NiTiO₃

BY H. BIRNBAUM AND R. K. SCOTT

By thermal combination, zinc oxide and oxides of elements of the fourth group of the periodic system, such as titanium, silicon, tin or zirconium, will react to form colorless or weakly colored compounds. If oxides of strong coloring elements, such as vanadium, chromium, manganese, iron, cobalt, nickel or copper, are substituted for part of the zinc oxide in these compounds, colored pigments of excellent stability can be produced.¹ Depth of color, as well as the color itself, can be varied by changing the proportions of the constituents and, in addition, the shade of a pigment of a given composition will change with proceeding reaction.

Since the structures of the pure titanates, zinc orthotitanate and nickel titanate, including their cell dimensions, have been worked out by N. W. Taylor,² we selected compositions of zinc oxide, nickel oxide and titanium dioxide for an X-ray diffraction study. Taylor's statement that zinc

(1) U. S. Patent 2,068,294 (1937).

(2) Taylor, *Z. physik. Chem.*, **96**, 242, 243, 259 (1930).