Lifetime of Tetrahedral Intermediates. In a recently advanced theory of stereoelectronic control of the hydrolysis of carboxylic acid esters and amides, it is proposed that the nature of the products obtained is governed by the conformation of the tetrahedral intermediate formed during the course of these reactions.²⁵ This theory requires tetrahedral intermediates to be relatively short lived; specifically in its application to cyclic, e.g., six-membered ring, systems it implies that these intermediates react before chair-chair interconversions can take place.²⁶ Since these tetrahedral intermediates in the case of ester hydrolysis are in fact the same as the hydrogen ortho esters detected in the present study, the measurements of their lifetimes made here have a direct bearing upon this assumption of the stereoelectronic theory. In particular, the data of Table II show that, under certain conditions, tetrahedral intermediates are relatively stable, with lifetimes approaching one second. Since chair-chair interconversions generally occur in times much shorter than this, this assumption of the stereoelectronic theory cannot be universally correct.

Acknowledgment. We are grateful to the Natural Sciences and Engineering Research Council Canada (formerly National Research Council of Canada), the National Science Foundation of the United States, and the donors of the Petroleum Research Fund, administered by the American Chemical Society, for their financial support of this work.

Supplementary Material Available: Tables S1-S4 of rate constants (7 pages). Ordering information is given on any current masthead page.

References and Notes

- (1) For recent reviews, see E. H. Cordes and H. G. Bull., Chem. Rev., 74, 581 (1974) T. H. File, Acc. Chem. Res. 8, 264 (1972).
 M. Ahmad, R. G. Bergstrom, M. J. Cashen, A. J. Kresge, R. A. McClelland,
- and M. F. Powell, J. Am. Chem. Soc., 99, 4827 (1977).
- A. L. Mori, M. A. Parzio, and L. L. Schaleger, *J. Am. Chem. Soc.*, **94**, 5034 (1972); A. L. Mori and L. L. Schaleger, *ibid.*, **94**, 5039 (1972).
 R. F. Atkinson and T. C. Bruice, *J. Am. Chem. Soc.*, **96**, 819 (1974).

- (5) B. Capon, *Pure Appl. Chem.*, **94**, 1001 (1977).
 (6) J. L. Jensen and P. A. Lenz, *J. Am. Chem. Soc.*, **100**, 1291 (1978).
 (7) B. Capon, K. Nimmo, and G. L. Reid, *J. Chem. Soc., Chem. Commun.*, 871
- (1976). (8) B. Capon, J. H. Hall, and D. M. A. Grieve, J. Chem. Soc., Chem. Commun., 1034 (1976).
- (9) R. A. McClelland and M. Ahmad, J. Am. Chem. Soc., 100, 7027 (1978).
- (10) R. A. McClelland, M. Ahmad, and G. Mandrapilias, J. Am. Chem. Soc., 101, 970 (1979).
- (11) R. A. McClelland and M. Ahmad, J. Org. Chem., in press.
 (12) (a) R. G. Bergstrom, M. J. Cashen, Y. Chiang, and A. J. Kresge, J. Org. Chem., in press; (b) Y. Chiang, A. J. Kresge, and C. I. Young, Finn. Chem. Chenr, in press, (b) P. Chang, A.J. Kresge, and C. Froung, *Finit. Chenr. Lett.*, 13 (1978); (c) Y. Chiang, A.J. Kresge, S. Salomaa, and C. I. Young, *J. Am. Chem. Soc.*, 96, 4494 (1974).
 C. F. Stoog, C. M. Chih, F. A. Short, and H. M. Woodburn, *J. Am. Chem. Soc.*, 71, 1710 (1949).
- (14) Tables S1, S2, S3, and S4 appear in the microfilm edition of this journal; see paragraph at end of this paper, regarding their availability. (15) Y. Chiang, A. J. Kresge and C. I. Young, *J. Org. Chem.*, **44**, 619 (1979).
- (16) A. A. Frost and R. G. Pearson, "Kinetics and Mechanisms", 2nd ed., Wiley, New York, 1961, p 167.
- (17) H. Meerwein, V. Hederich, and K. Wunderlich, Arch. Pharm., 291, 541 (1958).
- (18) (a) B. G. Ramsey and R. W. Taft, J. Am. Chem. Soc., 88, 3058 (1966). (b) R. A. McClelland and M. Ahmad, ibid., 99, 5356 (1977); 100, 7031 (1978).
- (19) L. P. Hammett, "Physical Organic Chemistry", 2nd ed., McGraw Hill, New York, 1970, p 290; J. T. Edward and S. C. Wong, J. Am. Chem. Soc., 99, 4229 (1977)
- (20) The observed rate constant obtained from the initial rapid phase of the 2-(*p*-methoxyphenyl)-1,3-dioxolenium ion decay, $k_{obsd} = (1.0 \pm 0.2) \times 10^3 s^{-1}$, measures the rate of approach to contribute 10^8 s⁻¹, measures the rate of approach to equilibrium and is actually equal to $k_0{}^2 + k_{H}{}^{-2}[H^+]$. However, at the acid concentrations where this rapid phase can be observed, i.e., where it is not too fast to be seen by the stopped-flow method, $k_{H^+}^{-2}[H^+]$ is small relative to k_0^2 ; this, coupled with the rather poor accuracy obtainable in measuring the rate for so fast a reaction makes $k_{obsd} \simeq k_0^2$ a good approximation. (21) C. A. Bunton and S. K. Huang, *J. Am. Chem. Soc.*, **94**, 3536 (1972). (22) Constant salt effects on pK values for structurally similar bases are com-
- monly found in acidity function studies; this, in fact, is the basis of the acidity function method.
- (23) Very large hydroxide ion catalytic coefficients of the same magnitude (6 $\times 10^{10} \, \text{M}^{-1} \, \text{s}^{-1}$) were also measured for the decomposition of hydrogen ortho esters derived from 2-(m-chlorophenyl)-, 2-(p-chlorophenyl)-, 2-(p-methylphenyl)-, and 2-(p-methoxyphenyl)-2-(N,N-dimethylamino)-1.3-dioxolanes.
- (24) This factor was originally reported as 42.2 ± 1.1 ,^{12c} but a revised value of $k_{\rm H^+}^1$ for 2-phenyl-2-methoxy-1,3-dioxolane^{12a} drops it down to $30.9 \pm$ 0.7.
- (25) P. Desiongchamps, Tetrahedron, 31, 2463 (1975).
- (26) P. Deslongchamps, R. Chenevert, R. J. Taillefer, C. Moreau, and J. K. Saunders, Can. J. Chem., 53, 1601 (1975).

The Substituent Effect on the Thermal Decomposition of Acetal Hydrotrioxides. Polar and Radical **Decomposition Paths**

Franci Kovač and Božo Plesničar*

Contribution from the Department of Chemistry, University of Ljubljana, 61001 Ljubljana, P.O. Box 537, Yugoslavia. Received October 31, 1978

Abstract: Low-temperature ozonation of aliphatic and aromatic acetals leads to oxygen-rich intermediates, acetal hydrotrioxides 1-3, the NMR spectra of which show two absorptions at δ 13 ppm downfield from Me₄Si. Decomposition of these compounds, which affords among other products singlet oxygen and peroxides, involves nonradical and radical processes. The kinetics of decomposition has been studied by NMR spectroscopy and activation parameters have been determined. Electronwithdrawing substituents in 3 accelerate decomposition in diethyl ether while electron-donating groups retard it (Hammett ρ value 1.2 ± 0.2). Considerable charge separation in the transition state for the decomposition has been proposed.

Introduction

The chemistry of organic polyoxides of the type RO_nR (R = alkyl; n = 3, 4) has been a matter of considerable interest during the past 2 decades.¹ Spectroscopic^{1k,n} and theoretical² evidence has been given in recent years for their existence.

Much less is known about alkyl hydropolyoxides, RO_nH . Alkyl hydrotrioxides have been proposed as unstable intermediates in ozonation of various organic compounds, i.e., ethers,³ silanes,⁴ amines,⁵ alcohols,⁶ ketones,^{6a,8} and aldehydes.^{6b,9} It was only recently that Murray et al. reported the first spectroscopic evidence for their existence.9b

hydrotrioxide	solvent	temp, °C ^a	k_1, s^{-1} b	E_{a} , kcal/mol	log A	δ ΟΟΟΗ (Me ₄ Si) –55 °C
18			2.7×10^{-2}	20.7	15.6	12.8 13.3
		-20	5.6×10^{-3}			(1:2.5)
	CH2Cl2c	-10	4.0×10^{-2}			12.7 13.1
	$(C_2H_3)_2O^2$	-10	2.9×10^{-2}	16.1	11.8	12.8 13.2
	(-25)2-	-20	8.5×10^{-3}			(1:2)
1b		-10	1.1×10^{-1}	14.5	11.1	12.8 13.2
		-20	3.7×10^{-2}			(1:1.5)
	$(C_{2}H_{3})_{2}O^{c}$	-10	7.6×10^{-2}	13.2	9.8	12.8 13.2
	(- 2 5) 2 -	-20	2.8×10^{-2}			(1:1)
2a	CH ₂ Cl ₂ c	-20	6.4×10^{-2}	20.5	16.6	13.20 13.25
	- -	-30	1.2×10^{-2}			(1:1)
	$(C_{2}H_{3})_{2}O^{c}$	-30	6.6×10^{-2}	15.9	13.1	13.10 13.15
	(-2- 0)2	-40	1.6×10^{-2}			
2b	CH2Cl2c	-20	3.8×10^{-2}	18.7	14.7	13.2
		-30	8.1×10^{-3}			
	$(C_{2}H_{5})_{2}O^{c}$	-40	1.1×10^{-2}			13.4
2c	$(C_2H_5)_2O^c$	-40	2.7×10^{-2}			13.9
3a	$(C_2H_5)_2O^c$	-30	7.6×10^{-2}	19.8	16.7	13.00 13.30
		-35	2.8×10^{-2}			(1:2.5)
		-40	1.3×10^{-2}			
3b	$(C_2H_5)_2O^{c}$	-30	6.6×10^{-2}	19.0	15.9	13.00 13.30
	/-	-35	2.9×10^{-2}			
		-40	1.2×10^{-2}			
3c	$(C_2H_5)_2O^{c}$	-30	6.3×10^{-2}	19.5	16.4	12.85 13.15
		-35	3.1×10^{-2}			(1:3.5)
		-40	1.1×10^{-2}			
3d	$(C_2H_5)_2O^{c}$	-35	3.2×10^{-2}	18.4	15.4	12.85 13.15
		-40	1.4×10^{-2}			
3e	$(C_2H_5)_2O^{c}$	-25	2.8×10^{-1}	25.1	21.6	12.65 12.95
		-30	9.5×10^{-2}			(1:2)
		-35	3.2×10^{-2}			
3f	$(C_2H_5)_2O^{c}$	-25	4.4×10^{-2}	17.2	13.7	12.85 13.10
		-30	2.0×10^{-2}			(1:4)
		-35	1.0×10^{-2}			
3g	$(C_2H_5)_2O^c$	-20	2.1×10^{-1}	21.4	17.8	13.05 13.25
		-25	8.1×10^{-2}			
		-30	3.1×10^{-2}			
		-35	1.4×10^{-2}			

Table I. Kinetic and Activation Parameters of the Decomposition of Acetal Hydrotrioxides

^{*a*} Temperature range was dictated by both solubility problems and fast decomposition with gas evolution. ^{*b*} Standard deviations $\pm 8\%$. ^{*c*} Acetal (20 mmol), 30%, solvent, 70% (by weight). The yield of hydrotrioxides estimated by NMR was 70–85%.

As a part of our continuous interest in polyoxides we report additional experimental evidence for the existence of hydrotrioxides in the low-temperature ozonation of acetals.¹⁰ Deslongchamps et al. have already proposed hydrotrioxides as possible intermediates in analogous reactions.¹¹

Results and Discussion

Ozonation of aliphatic and aromatic acetals (ozone-oxygen or ozone-nitrogen mixture) either neat (aliphatic acetals) or in diethyl ether, methylene chloride, and ethyl acetate below -55 °C produces unstable intermediates which upon warming to room temperature decompose vigorously. The low-temperature NMR spectra of these species show two absorptions at δ ca. 13 ppm downfield from Me₄Si. Since their position does not change much with dilution they are tentatively assigned to the OOOH absorptions of the intramolecularly hydrogen bonded six-membered rings of the acetal hydrotrioxides $1-3^{12}$ (Figure 1).

The decomposition of these intermediates was investigated by following the decay of OOOH absorptions and was found to obey first-order kinetics. The NMR data, along with kinetic and activation parameters for the decomposition, are collected in Table I.

The origin of the two absorptions for the OOOH protons remains uncertain. It is interesting to mention that Murray et al. made a similar observation in the case of 2-methyltetrahydrofuran hydrotrioxide.^{9b,13} Both peaks show a tendency to merging and broadening at higher temperatures. A characteristic feature is rather small splitting for **2a** (Figure 2). We assume that this phenomenon might be due either to the presence of two forms of the acyclic acetal hydrotrioxides with OOOH hydrogen bonded to one or the other of available oxygen atoms (nonequivalency of the two acetal oxygens with respect to OOOH group)¹⁴ or the presence of two conformational forms (chair-boat) of the six-membered ring with R groups on the acetal oxygen axial or equatorial. The fivemembered ring in **2** is nearly planar thus not allowing appreciable conformational changes.^{14b,15}

Activation parameters reported in Table I are in good qualitative agreement with those reported for other trioxides.^{1e,9b,16-18} A limited number of solvents suitable for the study of decomposition of compounds under investigation show that the effect of solvent is rather small. Comparable rate constants were obtained by studying decomposition of **1a** in the parent acetal, diethyl ether as well as methylene chloride.

Electron-withdrawing groups in 3 accelerate decomposition while electron-releasing groups retard it. A Hammett plot of the rates of decomposition of substituted derivatives of 3 in diethyl ether vs. σ values gives a fair correlation with $\rho = 1.2 \pm 0.2$ (Figure 3). A similar trend of the substituent effect on decomposition of substituted derivatives of 2 has been ob-



Table II. Singlet Oxygen Determination^a

hydrotrioxide	tetraphenylcyclopenta- dienone, %	1,3-diphenylisobenzo- furan, %
1b	20 ± 5^{b}	54 ± 5^{d}
2a	$55 \pm 3^{\circ}$	85 ± 8^{d}
3a		46 ± 5^{d}
3d	$30 \pm 5^{\circ}$	55 ± 8^d

^{*a*} Percent of absorbed ozone available to react with the reagent. ^{*b*} Neat or in methylene chloride. ^{*c*} Methylene chloride. ^{*d*} Ethyl acetate. Ozonation was performed with ozone-oxygen stream.

served.¹⁹ Considerable charge separation in the transition state for the decomposition is thus indicated.

All of the hydrotrioxides investigated decompose in the temperature range -45 to -10 °C to produce singlet oxygen as determined by typical singlet oxygen acceptors (Table II). The highest yields were obtained by using 1,3-diphenylisobenzofuran, which is known as one of the best singlet oxygen acceptors.²⁰ Since there is some evidence that both scavengers used may not be unambiguous as detectors of singlet oxygen, we studied decomposition of 2a in the presence of 1,2-dimethylcyclohexene. This compound has been known to distinguish between oxidation by singlet oxygen and a free-radical autoxidation on the basis of a different distribution of products for the two reactions.²¹ Although only 20-30% of the absorbed ozone was available to react with this reagent, it was shown that the reacting species is singlet oxygen. Namely, 1methyl-2-methylenecyclohexanol was found to be the major reaction product (80-90%).

A detailed study of the decomposition products of **1b** (ozonation with ozone-nitrogen mixture) by gas-liquid chromatography (GLC) showed ethyl acetate (0.80-0.85 mol/mol of absorbed ozone), ethanol (0.75-0.80 mol), water, acetaldehyde, acetic acid, ethyl formate, diethyl carbonate (0.06-0.12 mol), and a mixture of gases. Preparative thin layer chromatography (TLC) of the decomposition mixture at 5 °C on silica plates afforded α -hydroperoxydiethyl ether and α, α -diethoxydiethyl peroxide as well as hydrogen peroxide. Both organic peroxides account for roughly 5-7% of the decomposition products. The same types of peroxides were isolated in



Figure 2. Segments of NMR spectra of 2a: (A) in diethyl ether at -50 °C; (B) in methylene chloride.



Figure 3. Correlation of logarithms of first-order rate constants for the decomposition of 3 in diethyl ether vs. Hammett σ constants.

the decomposition of aromatic acyclic acetal hydrotrioxides, **3**.

Hydrogen peroxide has also been found in all these cases.²²

0.11

Table III. Spectral and Analytical Data of Peroxides 4 and 5, Isolated from the Deco	omposition Mixture of Hydrotrioxides 1 and 3
--	--

011

n

4a , $R_1 = b$, $R_1 = c$, $R_1 = d$, R_1	$C_{6}H_{5};$ $C_{6}H_{5};$ $4-FC_{6}H_{4};$	$R_2 = C_2 H_5$ $R_2 = CH_3$ $R_2 = C_2 H_5$ $R_2 = CH_3$	$\begin{array}{c} \mathbf{R}_{1} - \mathbf{C} - \mathbf{OOH} \\ \mathbf{C} \\ \mathbf{OR}_{2} \end{array}$	5a, R ₁ b, R ₁ c, R ₁ d, R ₁ e, R ₁	$= C_{6}H_{3};$ = C ₆ H ₅ ; = C ₆ H ₅ ; = 4-FC ₆ H ₄ ; = 4-ClC ₆ H ₄ ;	$R_2 = C_2 R_3$ $R_2 = C H_3$ $R_2 = C_2 H_5$ $R_2 = C H_3$ $R_2 = C H_3$	$\begin{array}{c} H \\ R_{2} - C - O \\ O R_{2} \end{array}$	$-O - C - R_1$
				ID h (n)=1	elemental anal., %		
compa	mpd'H NMR"				IK [®] (^µ OH), cm ·	calcu	Tound
4a <i>c</i>	9.95 (br s, 1 O (m, 6 CH ₃)	ОН), 5.10 (q, 1 Ј _{СН3СН} , = 6 Н	CH), 3.83 (m, 2 CH	H ₂), 1.30	3530, 34	00 (br)		
4b	9.04 (br s, 1 O 5.73 (s, 1 CI	OH), 7.40 (m, H), 3.60 (s, 3 C	5 C ₆ H ₅), CH ₃)		3510, 33	90 (br)	C 62.33 H 6.54	62.37 6.42
4c	9.20 (br s, 1 O 5.88 (s, 1 Cl	OH), 7.50 (m, H), 3.87 (m, 2	5 C ₆ H ₅), CH ₂), 1.18 (t, 3 OCI	H ₃)	3520, 33	90 (br)	C 64.27 H 7.19	64.57 7.07
4d	8.75 (br, 1 OO 5.80 (s, 1 CI	9H), 7.33 (m, 4 H), 3.64 (s, 3 C	$\begin{array}{l} 4\text{-}\mathrm{FC}_{6}\mathrm{H}_{4}\mathrm{),}\\ \mathrm{OCH}_{3}\mathrm{)}\end{array}$		3520, 33	80 (br)		
5a	5.10 (q, 1 CH) 1.32 (m, 6 C	, 3.83 (m, 2 Cl CH ₃), J _{CH3CH2}	H ₂), = 6 Hz					
5b <i>d</i>	7.33 (m, 5 C ₆ H	ł5), 5.77 (s, 1 C	CH), 3.55 (s, 3 OCH	3)				
5c	7.40 (m, 5 C ₆ H 3.80 (m, 2 C	H ₅), 5.93 (s, 1 C CH ₂), 1.25 (m, 4	CH), 6 CH ₃)					
5d	7.20 (m, 4 4-F) 3.55 (s, 3 OC	C ₆ H ₄), 5.77 (s, CH ₃)	1 CH),				C 61.93 H 5.21	62.25 5.81
5e	7.44 (s, 4 4-Cl 3.62 (s, 3 O	C ₆ H ₄), 5.83 (s, CH ₃)	1 CH),				C 55.99 H 4.72	56.01 4.90

^{*a*} CDCl₃, δ from Me₄Si. ^{*b*} CCl₄ (1-2% solutions). ^{*c*} Prepared also independently by the displacement method with 100% hydrogen peroxide (A. Rieche and C. Bischoff, *Chem. Ber.*, **94**, 2722 (1961)). ^{*d*} Prepared also independently by ozonolysis of *trans*-stilbene in methanol (W. P. Keaveney, M. G. Berger, and J. J. Pappas, *J. Org. Chem.*, **32**, 1537 (1967)). **4a**, **4d**, **5a**, **5b**, and **5c** are too volatile for elemental analysis.

It is interesting to mention that a previous attempt to isolate peroxidic products in the autoxidation of aliphatic and aromatic acyclic acetals failed.²³

~ ...

On the basis of the above-mentioned results we conclude that the formation of a large part of ethyl acetate, ethanol, and singlet oxygen can be tentatively ascribed to a nonradical decomposition while the presence of the other decomposition products suggests an alternative free-radical decomposition process accounting for roughly $20 \pm 5\%$ of the products (Scheme I).²⁴⁻²⁶

Scheme I



Experimental Section

Low-temperature NMR spectra were recorded on a JNM-C60H1 spectrometer equiped with a variable-temperature probe. IR spectra were recorded with a Perkin-Elmer 727B or 180 Model instrument. UV and visible spectra were obtained on a Perkin-Elmer Hitachi 200 spectrophotometer while a CEC 21-110C instrument was used to obtain mass spectra.

Materials. Acyclic²⁷ and cyclic²⁸ acetals were prepared by known procedures. The deuterated (C–D) derivative of the dimethyl acetal of benzaldehyde was prepared by the literature procedure.²⁹ All the acetals were checked by GLC, NMR, IR, and elemental analysis and found to be over 99% pure. Methylene chloride, diethyl ether, and ethyl acetate were purest commercially available products which were dried and distilled before use. Namely, it was found that water in these solvents prevents the observation of the OOOH absorptions.

Ozonations. Ozone was produced in a Gallenkamp ozonator which delivers an oxygen-ozone stream containing ca. 0.2 mmol of ozone per min.

Ozonation of the neat substrates (20 mmol) and solutions in diethyl ether, methylene chloride, and ethyl acetate (30% by wt) were carried out at -60 to -55 °C in the dark. Total ozonation times were 3 h for aliphatic acetals and 1 h for aromatic compounds. After ozonation the reaction mixture was thoroughly flushed with dried N₂ to remove excess ozone.

Kinetic Studies. Parts of the reaction mixture were transferred to a series of NMR tubes and the decay of the OOOH proton signals was followed by integrating the peak area with time at various temperatures (Me₄Si as an internal standard). The plots of the logarithm of peak areas vs. time were linear over several half-lives. Each rate constant is an average of at least three runs. First-order rate constants were obtained from a linear least-squares program.

Product Analysis. Singlet oxygen was determined by adding the acceptor (twofold excess per absorbed ozone) to the ozonized acetal at -50 to -40 °C and the reaction mixture allowed to warm up slowly to room temperature. The amount of absorbed ozone available to react with tetraphenylcyclopentadienone and 1,3-diphenylisobenzofuran was determined by measuring the decrease in absorbance at 510 and 405 nm, respectively. The corresponding reaction products, i.e., *cis*-dibenzoylstilbene (mp 214-215.5 °C, lit.²⁰ 215.9-216.3 °C) and *o*-dibenzoylbenzene (mp 143-145 °C, lit.²⁰ 146-147 °C), were isolated, although in lower yields, on silica gel chromatography columns according to the procedure of Wasserman et al.²⁰ Singlet oxygen determination (decomposition of **2a**) with 1,2-dimethylcyclohexene was

4 5

~ 11

performed by determining the amount of 1-methyl-2-methylenecyclohexanol (after sodium sulfite reduction of the corresponding hydroperoxide).9b,21

Decomposition products of 1b (ozonation with an ozone-nitrogen stream³⁰) were determined by a combination of techniques. All products except peroxides were determined by GLC on Porapak Q columns (80-160 °C) using calibrated internal standards and known reference materials. All the products were collected and identified by IR, NMR, and mass spectrum. The peroxidic materials were separated from the rest of the material on preparative silica plates (Merck Kieselgel 60 F₂₅₄, chloroform-n-hexane-methanol (10:10:1)). Preparative TLC was also used for the isolation of other peroxides reported in Table III (R_f , alkyl hydroperoxides 0.2-0.3; dialkyl peroxides, 0.5-0.6; hydrogen peroxide, 0.0).

Acknowledgment. The authors wish to thank the Boris Kidrič Fund for the financial support of this work, Professor O. Exner for stimulating discussions, and Professor B. Stanovnik for his help with the low-temperature NMR equipment.

References and Notes

- (1) (a) G. A. Russell, J. Am. Chem. Soc., 79, 3871 (1957); (b) G. O. Schenck and K. H. Schulte-Elke, Justus Liebigs Ann. Chem., 618, 185 (1958); (c) P. D. Bartlett and T. G. Traylor, J. Am. Chem. Soc., 85, 2407 (1963); (d) R. Hiatt, J. Clipsham, and T. Visser, *Can. J. Chem.*, **42**, 2754 (1964); (e)
 P. D. Bartlett and P. Günther, *J. Am. Chem. Soc.*, **88**, 3288 (1966); (f)
 P. D. Bartlett and G. Guaraldi, *ibid.*, **89**, 4799 (1967); (g)
 P. G. Thompson, *ibid.*, 89, 4316 (1967); (h) L. R. Anderson and W. B. Fox, ibid., 89, 4313 (1967); (i) N. A. Milas and B. Plesničar, ibid., 90, 4450 (1968); (j) J. A. Howard and K. U. Ingold, ibid., 90, 1056 (1968); (k) T. Mill and R. S. Stringham, ibid., 90, 1062 (1968); (I) J. E. Bennett, D. M. Brown, and B. Mile, Trans. Faraday Soc., 66, 397 (1970); (m) K. Adamic, J. A. Howard, and K. U. Ingold, Can. J. Chem., 47, 3803 (1969); (n) P. D. Bartlett and M. Lahav, Isr. J. Chem., Chem, 41, 3803 (1969), (n) P. D. Bartlett and M. Lanav, Isr. O Chem, 10, 101 (1972); (o) N. A. Clinton, R. A. Kenley, and T. G. Traylor, J. Am. Chem. Soc., 97, 3746, 3752 (1975); (p) R. A. Kenley and T. G. Traylor, *ibid.*, 97, 4700 (1975); J. E. Leffler and D. W. Miller, *ibid.*, 99, 480 (1977).
 (2) (a) B. Plesničar, S. Kaiser, and A. Ažman, J. Am. Chem. Soc., 95, 5476 (1973); (b) R. J. Blint and M. D. Newton, J. Chem. Phys., 50, 6220 (1973);
 (1) D. Varian, D. Karian, C. Amara, and A. Azara, b. Azara, b.
- (c) B. Plesničar, D. Kocjan, S. Murovec, and A. Ažman, J. Am. Chem. Soc., 98, 3143 (1976). (a) C. C. Price and A. L. Tumolo, *J. Am. Chem. Soc.*, 86, 4691 (1964); (b)
- (3)R. E. Erickson, R. T. Hansen, and J. Harkins, ibid., 90, 6777 (1968); (c) R. W. Murray, W. C. Lumma, Jr., and J. N.-P. Lin, ibid., 92, 3205 (1970); P. S. Bailey and D. A. Lerdal, ibid., 100, 5820 (1978).
- J. D. Austin and L. Spialter, *Adv. Chem. Ser.*, **No. 77**, 26 (1968). (a) P. S. Bailey, D. A. Mitchard, and A.-J. Y. Khashab, *J. Org. Chem.*, **33**, 2675 (1968); (b) P. S. Bailey and J. E. Keller, *ibid.*, **33**, 2680 (1968); (c) P. (5)S. Bailey, T. P. Carter, Jr., and L. M. Southwick, ibid., 37, 2997 (1972).
- (a) M. C. Whiting, A. J. N. Bolt, and J. H. Parish, *Adv. Chem. Ser.*, No. 77, 4 (1968); G. A. Olah, N. Yoneda, and R. Ohnishi, *J. Am. Chem. Soc.*, 98, 7341 (1976).
- (7) P. S. Bailey, A. M. Reader, P. Kolsaker, H. M. White, and J. C. Barborak, J. Org. Chem., 30, 3042 (1965).
- (a) G. A. Hamilton, B. S. Ribner, and T. M. Hellman, Adv. Chem. Ser., No. 77, 15 (1968); (b) J. E. Batterbe and P. S. Bailey, *J. Org. Chem.*, 32, 3899 (1967); (c) G. A. Olah, N. Yoneda, and D. G. Parker, *J. Am. Chem. Soc.*, (a) H. M. White and P. S. Bailey, J. Org. Chem., 30, 3037 (1965); (b) F. E.
- Stary, D. E. Emge, and R. W. Murray, J. Am. Chem. Soc., 96, 5671 (1975); (b) F. E. 98, 1880 (1976).
- (10) A part of this work was presented in a preliminary form: F. Kovač and B. Plesničar, J. Chem. Soc., Chem. Commun., 122 (1978)
- P. Deslongchamps, P. Atlani, D. Frehel, A. Malaval, and C. Moreau, Can (11)J. Chem., 52, 3651 (1974).

- (12) Ozonation of the dimethyl acetal of C-D deuterated benzaldehyde produces an oxygen-rich intermediate which does not show any absorption around δ 13 ppm
- (13) These authors suggest that the hydrogen bonding can involve either of the lone pairs of the tetrahedral ring oxygens and that some interconversion of these two forms might be taking place. (14) L. S. Rattet, L. Mandel, and J. H. Goldstein, *J. Am. Chem. Soc.*, **89**, 2253
- (1967); O. Exner, V. Jehlička, and B. Uchytil, Collect. Czech. Chem. Commun., 33, 2862 (1968).
- (15) Attempts were made to detect any splitting of methyl proton absorptions. Unfortunately, the spectrum is too complicated owing to the presence of some unreacted acetal and decomposition products. The ¹⁹F NMR spectrum of the ozonized dimethyl acetal of perfluorobenzaldehyde is indistinguishable from that of the parent acetal. (16) R. W. Murray and M. L. Kaplan, *J. Am. Chem. Soc.*, **91**, 5358 (1969). (17) E. Koch, *Tetrahedron*, **26**, 3503 (1970).

- S. W. Benson and R. Shaw in "Organic Peroxides", Vol. 1, D. Swern, Ed., (18)Wiley-Interscience, New York, 1970, p 126. (19) Attempts to study the kinetics of the decomposition of other para-substituted
- hydrotrioxide derivatives were frustrated by precipitation of these species at temperatures employed. These solids begin to decompose vigorously at about -30 °C.
- (20) H. H. Wasserman, J. R. Scheffer, and J. L. Cooper, J. Am. Chem. Soc., 94,
- 4991 (1972).
 (21) C. S. Foote, Acc. Chem. Res., 1, 104 (1968); J. A. Howard and G. D. Mendenhall, Can. J. Chem., 53, 2199 (1975).
- (22) The corresponding esters and alcohols were the major products in the decomposition of 3, accounting for 80-90% of all products as determined by IR.
- (23) H. E. Seyfarth, A. Rieche, and A. Hesse, Chem. Ber., 100, 624 (1967).
- (24) The exact amount of products attributed to radical decomposition paths cannot be determined since the peroxides decompose during the GLC determination to produce ethyl formate (the major product) as well as ethyl acetate, ethanol, acetaldehyde, water, and gases (methane, ethane, butane).
- (25) Free-radical decomposition of the hydrotrioxide 1b might, analogously to hydroperoxides, involve an induced decomposition as shown below (R. Hiatt and T. McCarrick, *J. Am. Chem. Soc.*, **97**, 5234 (1975)).



- (26) Relatively smaller evolution of gas has been observed in most cases even after the disappearance of the OOOH absorption. This might be due to the decomposition of peroxides and/or polyoxides formed by radical recombinations.
- A. C. Bottomley, W. Cocker, and P. Nanny, *J. Chem. Soc.*, 1891 (1937). T. H. Fife and L. K. Jao, *J. Org. Chem.*, **30**, 1492 (1965).
- (28)
- (29)
- K. B. Wiberg, J. Am. Chem. Soc., 76, 5371 (1954).
 P. S. Bailey and A. M. Reader, Chem. Ind. (London), 1063 (1961); H. K. (30)Reimschuessel and G. A. Mountford, J. Colloid Interface Sci., 25, 558 (1967)
- Presented in part at the 4th IUPAC Symposium on Physical Organic (31) Chemistry, York, Sept 4-8, 1978, Abstract B30.