- (12) R. F. Hudson and J. G. Verkade, Tetrahedron Lett., 3231 (1975).
- (13) A. E. Kalinin, V. G. Andrianov, and Y. T. Strouchkov, Zh. Strukt. Khim., 16, 1041 (1975).
- (14) The synthesis of this compound was accomplished by the general reaction of the phosphonous dichloride with 3,3-dimethylpentane-1,5-diol (D. W. White, *Phosphorus*, 1, 33 (1971)) and the compound is characterized by NMR spectroscopy (¹H, ¹³C, ³¹P) and elemental analysis. Although stable at low temperature, it polymerizes within a few hours at room temperature.
- (15) M. Dräger, Z. Anorg. Alig. Chem., 411, 79 (1975); M. Dräger and R. Engler, *ibid.*, 413, 229 (1975); J. J. Daly and F. Sanz, J. Chem. Soc., Dalton Trans., 2051 (1974); P. Maroni, H. Holeman, J. G. Wolf, L. Ricard, and J. Fisher, Tetrahedron Lett., 15, 1193 (1976).
- Tetrahedron Lett., 15, 1193 (1976). (16) Faculty member of the Université Scientifique et Médicale de Grenoble.

J. P. Dutasta, J. B. Robert*¹⁶

Laboratoire de Chimie Organique Physique Département de Recherche Fondamentale Equipe de Recherche Associée No. 674 au CNRS Centre d'Etudes Nucléaires de Grenoble 85 X F. 38041 Grenoble Cedex, France Received August 15, 1977

A Highly Stereoselective Synthesis of Z-Trisubstituted Olefins via [2,3]-Sigmatropic Rearrangement. Preference for a Pseudoaxially Substituted Transition State

Sir:

Sigmatropic rearrangements of the [2,3] and [3,3] type are some of the most widely used reactions for the stereoselective preparation of di- and trisubstituted olefins.¹ Previous examples² of these reactions have led mainly to *E* olefins via cyclic transition states having the greatest possible number of pseudoequatorial substitutents. We have found, however, that this preference for transition states having pseudoequatorial substitution may not be as general as previously assumed. We report here a convenient and efficient [2,3]-sigmatropic Wittig rearrangement which produces *Z*-trisubstituted homoallylic alcohols in high yields and with \geq 95% stereoselectivity. To the best of our knowledge this reaction is the first example of a [2,3]-sigmatropic rearrangement having a transition state which prefers pseudoaxial substitution.³

As outlined below, when the allylic alcohol 1 was deproto-



nated (KH, THF) and alkylated with iodomethyltributyltin⁴ (30 min, 25 °C), the allyl stannylmethyl ether 2 (NMR (CCl₄) $\delta 4.80 (=CH_2, m), 3.45 (-OCH_2Sn, ABq, \Delta \nu_{AB} = 27 Hz, J$ = 10 Hz), 3.30 (>CHO-, m)) was formed quantitatively.⁵ Although this material could be isolated, direct treatment of the chilled reaction mixture with excess n-butyllithium resulted in tin/lithium exchange and smooth [2,3]-sigmatropic rearrangement (30 min, -78 °C) to the homoallylic alcohol 3 $(>95\% \text{ yield}; \text{NMR} (\text{CCl}_4) \delta 5.22 (=CH-, \text{ br t}, J = 8 \text{ Hz}),$ $3.56 (-CH_2OH, t, J = 8 Hz), 2.24 (=C(Me) CH_2-, t, J = 8$ Hz), 1.69 (=CMe-, br s)). VPC (Carbowax) analysis showed 3 to be 96–97% one isomer and an examination by ^{13}C NMR strongly suggested this isomer to be the Z one shown. This tentative assignment followed from the upfield⁶ value (δ 34.4) of carbon 2 and a normal value (δ 21.6) for the relatively unhindered carbon 9.7 Conformation was obtained by comparison with authentic E isomer 4 prepared by the method of Johnson.⁸ Coinjection VPC and ¹H NMR showed that the minor isomer in 4 was identical with the major isomer in 3 and vice versa. Furthermore the ¹³C NMR of **4** showed a shielding pattern for

C-2 (δ 43.0) and C-9 (δ 16.0) which was opposite to that observed in the [2,3]-sigmatropic rearrangement product 3. Thus the olefinic stereochemistry of our product may be unambiguously assigned as Z.

Aside from the obvious synthetic potential of this reaction, the transition-state geometry is of special interest. It is clear from the stereochemistry of 3 that transition state A having a pseudoaxial butyl substituent is strongly preferred over transition state $B.^9$ Since the stereoelectronics of the two



transition states are virtually identical, the observed preference for A implies that the steric interaction of the butyl substituent with the C-1 methylene is much less severe than the alternative butyl/vinyl methyl interaction in B. This state of affairs is understandable given the two reasonable assumptions that the transition state is very early¹⁰ (C-1 is sp² and C-3 is sp³) and that the planes defined by C-1-C-2-C-3 and C-1-C-3-O-4-C-5 are not quite perpendicular ($\theta > 90^\circ$). A careful examination of the geometry of A reveals that, while the pseudoequatorial substituent (hydrogen in A) almost eclipses the methyl attached to C-2, the pseudoaxial substituent (butyl in A) is in fact rather far removed from the C-1 methylene.

Further support for our postulated transition-state geometry came from a study of the [2,3]-sigmatropic rearrangement of allylic ethers having different substitution patterns. If for example the vinyl methyl in 2 was replaced by hydrogen then the model above would predict a marked decrease in the selectivity for transition state A and thus for the Z product. When the allylic alcohol 5 was rearranged as above, a 40:60 mixture of homoallylic alcohols 6 and 7 (>95% yield) was obtained.^{11,12}

$$C_7H_{15} \longrightarrow C_7H_{15} \longrightarrow OH + C_7H_{15} \longrightarrow OH$$

A similar result was found with the trans allylic alcohol 8: a roughly 35:65 mixture of compounds 9 and 10 (96% yield) was produced. The cis allylic alcohol 11 on the other hand would be expected to strongly destabilize transition state A relative to B, and in fact rearrangement of 11 led exclusively to 9: 91% yield; IR (neat) 965 cm⁻¹; NMR (CCl₄) δ 5.35 (-CH=CH-, J = 15 Hz).



Synthetically the reaction has a number of interesting features. The rearrangement proceeds smoothly in cyclic systems so long as the β carbon is relatively unhindered. 2-Cyclohexenol for example was converted to the hydroxymethylcyclohexene 12 in 93% yield. The more hindered octalols 13 and 14 reacted

only sluggishly to yield mixtures of [1,2] and [2,3] rearranged products in low yield. It is also possible to trap the intermediate alkoxymethyllithium intermediates with reactive electrophiles prior to rearrangement. For that reaction the stannylmethyl ether (e.g., 2) is first isolated and then transmetalated in a second step in the absence of strong chelating agents. Thus

addition of n-butyllithium to 2 at -78 °C (THF, 30 s) followed by immediate quenching with cyclohexanone produced 15: 73% yield; IR (neat) 3400 cm⁻¹; NMR (CCl₄) δ 6.84 (=CH₂, br s), 3.55 (>CHO-, m), 3.03 (-OCH₂-, AB q, $\Delta \nu_{AB} = 20$ Hz, J = 9 Hz, 1.62 (==CMe-, s).



At this time the most useful aspect of the reaction seems to be its high stereoselectivity for Z-trisubstituted olefins. While a variety of natural products incorporate double bonds of this type, most trisubstituted olefin syntheses produce E isomers as the major products (see exceptions 13).

The trienvl acetate 19 is an active component in the sex attractant of a major citrus pest, the California red scale.¹⁴ We have prepared this compound from the ethyl ester of β_{β} . dimethylacrylic acid as summarized in Scheme I. Alkylation Scheme I



(LDA, THF/HMPA, -78 °C) with 4-bromobutene gave the β , γ -unsaturated ester 16 in 80% yield. Hydride reduction and NBS-phosphine bromination¹⁵ then gave 17 (93% yield). Conversion to the Grignard reagent and reaction (Et_2O , -20 °C) with methacrolein proceeded in 52% yield and led to a mixture of diastereomeric allylic alcohols 18 which was suitable for the [2,3]-sigmatropic rearrangement described above. Alkylation with iodomethyltributyltin and butyllithium-induced rearrangement (30 min, -78 °C) proceeded as usual. The product was chromatographed and acetylated (Ac₂O/pyr) to yield the racemic red scale triene 19 in 83% yield. VPC and NMR comparison with authentic material¹⁶ showed our product to consist of 95–96% of the desired Z isomer.¹⁷

References and Notes

- (1) Stereoselective olefin synthesis. Reviews: D. J. Faulkner, Synthesis, 175 (1971); J. Reucroft and P. G. Sammes, Quart. Rev., Chem. Soc., 25, 135 1971)
- Trisubstituted [2,3]: (a) G. Büchi and H. Wüest, Helv. Chim. Acta, 50, 2440 (2) (1967); K. B. Sharpless and R. F. Lauer, J. Am. Chem. Soc., 94, 7154
 (1972); (b) P. A. Grieco, J. Chem. Soc., Chem. Commun., 702 (1972); D. A. Evans, G. C. Andrews, T. T. Fujimoto, and D. Wells, Tetrahedron Lett., 1389 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Org. Chem., 38, 959 (1973); (c) P. A. Grieco, D. Boxler and K. Hiroi, J. Chem., 959 (1975); (c) P. A. Grieco, P. Boxler and K. Hiroi, J. Chem., 950 (1975); (c) P. A. Grieco, P. Boxler and K. Hiroi, J. Chem., 950 (1975); (c) P. A. Grieco, P. Boxler and K. Hiroi, J. Chem., 950 (1975); (c) P. A. Grieco, P. Boxler and K. Hiroi, J. Chem., 950 (1975); (c) P. A. Grieco, P. Boxler and F. Hiroi, P. Boxler and F. Hiroi 2572 (1973). Trisubstituted [3,3]: (d) review, F. E. Ziegler, Acc. Chem. Res., 10, 227 (1977).
- (3) Cf. W. Sucrow and W. Richter, *Chem. Ber.*, **104**, 3679 (1971); W. Sucrow,
 B. Schubert, W. Richter and M. Slopianka, *ibid.*, 3689 (1971).
- W. C. Still, J. Am. Chem. Soc., in press.
- With more hindered starting alcohols, addition of an equivalent of 18-crown-6 frequently gives higher yields of the final homoallylic alcohol. γ effect: J. W. De Hann and L. J. M. van de Ven, *Org. Magn. Reson.*, 5, 147 (1973); C. Nishino and W. S. Bowers, Tetrahedron, 32, 2875 (1976); M.
- J. Pettei, F. G. Pilkiewicz and K. Nakanishi, Tetrahedron Lett., 2083 1977)
- (1977).
 (7) Cf. J. D. Roberts, F. J. Weigert, J. I. Kroschwitz, and H. J. Reich, J. Am. Chem. Soc., 92, 1338 (1970); D. E. Dorman, M. Jautelat and J. D. Roberts, J. Org. Chem., 36, 2757 (1971).
 (8) S. F. Brady, M. A. Ilton, and W. S. Johnson, J. Am. Chem. Soc., 90, 2882
- (1968).



(9) Cf. J. E. Baldwin and J. E. Patrick, J. Am. Chem. Soc., 93, 3556 (1971). (10) In a highly exothermic process like this one, an early transition state would be anticipated by the Hammond postulate: L. Melander, Chem. Soc., Spec. Publ., No. 16, 119 (1962).

- (11) Other [2,3]-sigmatropic rearrangements of more stabilized allylic ether anions gave largely trans disubstituted olefins and include U. Schöllkopf and K. Fellenberger, Justus Liebigs Ann. Chem., 698, 80 (1966); Y. Makisumi and S. Notzumoto, Tetrahedron Lett., 6393 (1966); V. Rautenstrauch, J. Chem. Soc., Chem. Commun., 4 (1970); ref 9; G. Büchi, M. Cushman, and H. Wüest, J. Am. Chem. Soc., 96, 5563 (1974).
- (12) Structures were confirmed by comparison with authentic samples prepared from the corresponding acetylene.
- Trom the corresponding activitient.
 B. M. Trost and T. J. Fullerton, J. Am. Chem. Soc., 95, 292 (1973); C. P. Casey and D. F. Marten, Synth. Commun., 3, 321 (1973); S. Kobayashi, H. Takei and T. Mukayama, Chem. Lett., 1097 (1973); S. B. Bowlus and J. A. Katzenellenbogen, J. Org. Chem., 38, 2733 (1973); S. Tanaka, H. Yamamoto, H. Nozaki, K. B. Sharpless, R. C. Michaelson, and J. D. Cutting, J. Am. Chem. Soc., 96, 5254 (1974); B. S. Pritzele, J. S. Baran, and D. H. Steinman, J. Org. Chem., 40, 269 (1975); A. Pelter, K. J. Gould, and C. R. Lucina. Textbackne, 44, 223 (1275); K. Ukbide, K. Michaelson, and J. H. Steinman, J. Org. Chem., 40, 269 (1975); A. Pelter, K. J. Gould, and C. R. Lucina. Textbackne, 44, 223 (1275); K. Ukbide, K. Michaelson, and M. Steinman, Steinman, Steinman, 2010, 2 Harrison, Tetrahedron Lett., 3327 (1975); K. Uchida, K. Utimoto, and H. Nozaki, J. Org. Chem., 41, 2941 (1976); E. J. Corey, P. Ulrich, and A. Venkateswarlu, Tetrahedron Lett., 3231 (1977), and references therein. (14) W. L. Roelofs, M. J. Gieselmann, A. M. Carde, H. Tashiro, D. S. Moreno,
- C. A. Henrick, and R. J. Anderson, Nature, 267, 698 (1977). (15) A. K. Bose and B. Lal, Tetrahedron Lett., 3937 (1973).
- (16) An authentic sample of the red scale triene and its E isomer was kindly provided by R. J. Anderson of Zoecon Corporation, Palo Alto, Calif.
- (17) This work was generously supported by grants from Research Corporation, the donors of the Petroleum Research Fund, administered by the American Chemical Society, and Grant Number CA 23084, awarded by the National Cancer Institute, DHEW.

W. Clark Still,* Abhijit Mitra

Department of Chemistry, Columbia University New York, New York 10027 Received November 7, 1977

Concerning CH2==C-• and Its Reaction with ¹⁴N¹⁵NO

Sir:

In their paper entitled "Gaseous Anion Chemistry",¹ Smit and Field report than an ion of m/e 26 is always present in the negative ion CI spectrum of the bath gas which they used $(N_2O/CH_4 \ 1:1)$. Their work with CD₄, measurements of natural isotope abundance ratios of the ion, and their observation that the relative intensity of the ion was dependent on the temperature of the source filament all suggest that the ion is CN^- rather than $C_2H_2^-$. We would point out that further evidence to support this view comes from the absence of a peak at m/e 40, most significantly with the sample 1-octyne where the m/e 26 intensity was enhanced. It is known that m/e 40, CH_2CN^- , is formed by the reaction of $C_2H_2^-$ with N_2O^2 The route which was suggested to these authors to explain the formation of CN⁻, namely:

$$C^{-} \cdot + N_2 O \to C N^- + N O \cdot \tag{1}$$

seems extremely attractive until it is recalled that Fehsenfeld and Ferguson³ specifically did not observe the reaction under flowing afterglow conditions. Instead, they record that these collisions led to a rapid associative electron detachment reaction, but it was not known if the products of the reaction were $CN + NO + e \text{ or } N_2 + CO + e \text{ or something else.}$

The alternative composition for m/e 26, viz. $C_2H_2^{-1}$, has been observed in a number of independent laboratories.4-7 MINDO/3 calculations⁸ which have now been performed in our laboratory⁹ predict electron affinities of 0 eV for CH₂==C: and -1.8 eV for HC≡CH (in their ground states), so that acetylenic molecular anions should not be stable with respect to electron detachment. Furthermore, we have recently discovered that at near-zero electron energy cis-1,2-difluoroethylene undergoes a resonant dissociative electron attachment process giving in good yield ions of m/e 44:

$$e + cis-CHF \Longrightarrow C_2HF \longrightarrow + HF$$
 (2)

However, we have so far been unable to find any ionization of the isomeric 1,1-difluoroethylene. This divergence of behavior