J.C.S. Dalton

## Reactions of Stable [PtCl<sub>2</sub>( $\eta^2$ -olefin)( $\alpha$ -di-imine)] Complexes. Retention of the Five-co-ordinate Structure upon Halogen Exchange and Ligand Substitution with Olefins, $\alpha$ -Di-imines, and *N*,*N'*-Disubstituted 1,2-Diaminoethanes

By Henk van der Poel, Gerard van Koten,\* and Gerard C. van Stein, Anorganisch Chemisch Laboratorium, University of Amsterdam, J. H. van't Hoff Instituut, Nieuwe Achtergracht 166, 1018 WV Amsterdam, The Netherlands

The axial halogen atoms as well as the equatorial  $\eta^2 \cdot C_2 H_4$  and  $\sigma, \sigma \cdot N, N'$  chelate-bonded R - N = CH - CH = N - R or  $R - N(H) CH_2 CH_2 N(H) - R$  ligands (L-L) in five-co-ordinate  $[PtCI_2(\eta^2 - C_2 H_4)(\sigma, \sigma - N, N' - L - L)]$  can be replaced with retention of the trigonal-bipyramidal structure. Halogen exchange is initiated by formation of an ionic intermediate  $[PtCI(\eta^2 - C_2 H_4)(L-L)]CI$ . The reversible exchange of the equatorial ligands with olefins,  $\alpha$ -di-imines, or 1,2-diamine ligands is proposed to proceed *via* five-co-ordinate intermediates. The selectivity observed in these exchange reactions is explained on the basis of the difference in steric constraints of the various olefins and bidentate ligands used.

EXCHANGE of the N-donor ligand (L) and olefin in  $[PtX(Y)(\eta^2-\text{olefin})L]$  complexes has been demonstrated to proceed via the five-co-ordinate transition states  $[PtX(Y)(\eta^2-\text{olefin})L(L')](L'=N'-\text{donor ligand or olefin})^{1,2}$  The five-co-ordinate transition state is reached either by association with the N'-donor ligand or solvent molecules. These exchange mechanisms have not only been experimentally studied, but theoretical studies have recently been reported also.<sup>1,2</sup> Owing to the transient character of these five-co-ordinate intermediates, aspects such as ligand exchange and intramolecular rearrangements via concerted mechanisms (Berry pseudo-rotation or Turnstile processes) which influence the products formed are difficult to investigate.

Recently, a large series of stable five-co-ordinate  $[PtX_{2}(\eta^{2}-olefin)(L-L)]$  (L-L = bidentate donor ligand) complexes were reported in the literature <sup>3-7</sup> which makes the gathering of important information possible. It was seen that in these complexes where  $\bar{L}\mathchar`-L$  was an  $\alpha$ -di-imine (R-N=CHCH=N-R), R-N(H)CH<sub>2</sub>CH<sub>2</sub>N(H)-R, or the 2,6-disubstituted pyridine R'NC<sub>5</sub>H<sub>3</sub>CH=NR the various ligands show a strong site preference in the trigonal-bipyramidal array. The electronegative X atoms reside in the axial positions, while the neutral ligands are in the equatorial plane. For these complexes, no experimental evidence for the occurrence of stereoisomerization processes could be obtained.4,5,7 However, we did observe dynamic behaviour in the RN= CHCH=NR and RN(H)CH<sub>2</sub>CH<sub>2</sub>N(H)R complexes, as evidenced by multinuclear n.m.r. spectra, involving pure olefin rotation. The  $\Delta G$  values (13–15 kcal \*) for the barrier to olefin rotation were in the range found for fourco-ordinate platinum-olefin species.<sup>4</sup> It was concluded that bending back of the axial atoms in the direction of the bidentate ligand, when the olefin passed the upright conformation, was an important factor. The kinetic stability of these five-co-ordinate species with respect to platinum-olefin bond dissociation to give four-coordinate  $[PtX_2(L-L)]$  complexes depends mainly on steric factors. For example introduction of a methyl group in the 6-position of R'NC<sub>5</sub>H<sub>3</sub>CH=NR (*i.e.* R' = Me) enhanced the solution stability considerably. This was tentatively explained by the fact that in four-co-ordinate  $[PtCl_2(R'NC_5H_3CH=NR)]$  complexes all the ligands reside in one plane which necessarily brings one of the atoms close to the 6-R' substituent.<sup>5</sup>

In this paper we describe studies concerning intermolecular exchange processes of the five-co-ordinate platinum-olefin complexes with olefins, uni- and bidentate N-donor ligands, and sodium halides which occur with retention of the five-co-ordinate geometry.<sup>8</sup>

### EXPERIMENTAL

α-Di-imines (RNCHCHNR), N,N'-substituted 1,2-diaminoethanes [RN(H)CH<sub>2</sub>CH<sub>2</sub>N(H)R], and 6-R'-pyridine-2carbaldimines (R'NC<sub>5</sub>H<sub>3</sub>CH=NR) were prepared by methods described elsewhere.<sup>9</sup> Five-co-ordinate [PtX<sub>2</sub>(η<sup>2</sup>-olefin)(L-L)] complexes [L-L = RNCHCHNR, RN(H)CH<sub>2</sub>CH<sub>2</sub>N(H)R, R'NC<sub>5</sub>H<sub>3</sub>CH=NR, 2,2'-bipyridyl (bipy), or 2,11-dimethylphenanthroline (dmphen)] were obtained from the reaction of Zeise's salt {K[PtCl<sub>3</sub>(η<sup>2</sup>-olefin)]} with L-L.<sup>4-6</sup> Solvents were purified and distilled before use; all exchange reactions were carried out in a nitrogen atmosphere.

*Exchange Reactions.*—The complexes obtained from these experiments were characterized by comparison of their spectroscopic data ( ${}^{1}H$ ,  ${}^{13}C$  n.m.r., and i.r.) with those of authentic samples.<sup>4,5</sup>

Exchange of  $\sigma$ -N-bonded L and  $\sigma,\sigma$ -N,N'-bonded L-L donor ligands in [PtCl<sub>2</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>)L]. The preparation of [PtCl<sub>2</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>){Bu<sup>t</sup>N(H)CH<sub>2</sub>CH<sub>2</sub>N(H)Bu<sup>t</sup>}] from [PtCl<sub>2</sub>-( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>){Bu<sup>t</sup>NCHCHNBu<sup>t</sup>}] (1) described here is representative. The compound Bu<sup>t</sup>N(H)CH<sub>2</sub>CH<sub>2</sub>N(H)Bu<sup>t</sup> (2 mmol) was added to a stirred solution of [PtCl<sub>2</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>)-(Bu<sup>t</sup>NCHCHNBu<sup>t</sup>)] (1) (1 mmol) in dichloromethane (10 cm<sup>3</sup>). The solution was stirred for 10 min, evaporated to dryness, and the residue washed with pentane (4 × 10 cm<sup>3</sup>) to remove Bu<sup>t</sup>NCHCHNBu<sup>t</sup> as well as the excess of Bu<sup>t</sup>-N(H)CH<sub>2</sub>CH<sub>2</sub>N(H)Bu<sup>t</sup>. Yellow microcrystalline [PtCl<sub>2</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>){Bu<sup>t</sup>N(H)CH<sub>2</sub>CH<sub>2</sub>N(H)Bu<sup>t</sup>}] was isolated in 75% yield. When R'NC<sub>5</sub>H<sub>3</sub>CH=NR, bipy, or dmphen were used, the reaction mixture was cooled to -20 °C directly after mixing

<sup>\*</sup> Throughout this paper: 1 cal = 4.184 J; 1 atm = 101 325 Pa.

the reactants in order to minimize decomposition according to  $[PtCl_2(\eta^2-C_2H_4)(L-L)] \rightarrow [PtCl_2(L-L)] + C_2H_4$ .

Halogen exchange.  $[PtI_2(\eta^2-C_2H_4)(L-L)][L-L=Bu^tNCH-CHNBu^t, EtMe_2CNCHCHNCMe_2Et, or Bu^tN(H)CH_2CH_2-N(H)Bu^t] from [PtCl_2(\eta^2-C_2H_4)(L-L)]. The complex [PtCl_2(\eta^2-C_2H_4)(L-L)] (1 mmol) was added to a stirred solution of NaI (6 mmol) in acetone (10 cm<sup>3</sup>). The reaction mixture was stirred for 24 h, after which NaCl was filtered off. The filtrate was evaporated to dryness and the residue extracted with benzene (3 × 30 cm<sup>3</sup>). Concentration of the benzene extract resulted in orange-red crystals of [PtI_2(\eta^2-C_2H_4)(L-L)]. The yield after recrystallization from a hexane-benzene mixture was 40-60%.$ 

Halogen-halogen exchange between  $[PtX_2(\eta^2-C_2H_4)(RNCH-CHNR)] (X = Cl or Br; R = Bu<sup>t</sup> or CMe_2Et). The complex <math>[PtCl_2(\eta^2-C_2H_4)(RNCHCHNR)]$  (0.2 mmol) and  $[PtBr_2(\eta^2-C_2H_4)(RNCHCHNR)]$  (0.2 mmol) were dissolved in oxygen-free CDCl<sub>3</sub> (0.5 cm<sup>3</sup>) and then sealed in a n.m.r. tube. For the exchange reaction monitored by <sup>1</sup>H and <sup>13</sup>C n.m.r. spectrometry for 3 weeks see Results section.

Exchange of ethylene in  $[PtCl_2(\eta^2-C_2H_4)(Bu^tNCHCHNBu^t)]$ (1). The preparation of  $[PtCl_2(\eta^2-CH_2=CHCO_2Me)(Bu^t-NCHCHNBu^t)]$  from  $[PtCl_2(\eta^2-C_2H_4)(Bu^tNCHCHNBu^t)]$  described here is representative.

Methyl acrylate (4 mmol) was added to a stirred solution of  $[PtCl_2(\eta^2-C_2H_4)(Bu^tNCHCHNBu^t)]$  (1 mmol) in dichloromethane (10 cm<sup>3</sup>). The solution was stirred for 24 h and then concentrated below 25 °C. The residue was washed

with pentane  $(4 \times 10 \text{ cm}^3)$  to remove excess of methyl acrylate and dried *in vacuo*. Y1eld 80% {Found: C, 31.85; H, 5.10; N, 5.35; O, 5.70. Calc. for [PtCl<sub>2</sub>{ $\eta^2$ -CH<sub>2</sub>= CHCO<sub>2</sub>Me)(Bu<sup>t</sup>NCHCHNBu<sup>t</sup>)]: C, 32.3; H, 5.05; N, 5.40; O, 6.15%. Found: C, 31.75; H, 4.95; N, 4.65. Calc. for [PtCl<sub>2</sub>{ $\eta^2$ -C<sub>2</sub>H<sub>2</sub>(CO<sub>2</sub>Me)<sub>2</sub>}(Bu<sup>t</sup>NCHCHNBu<sup>t</sup>)]: C, 33.25; H, 4.90; N, 4.85% }.

Separation of a 1:1 cis-trans mixture of  $(MeO_2C)CH=$ CH(CO<sub>2</sub>Me) by means of  $[PtCl_2(\eta^2-C_2H_4)(Bu^tNCHCHNBu^t)]$ (1). A solution of dimethyl maleate (*cis* 0.5 mmol) and dimethyl fumarate (*trans* 0.5 mmol) was added to a solution of (1) (2.0 mmol) in dichloromethane (10 cm<sup>3</sup>) which was stirred for 48 h and then concentrated to 2 cm<sup>3</sup>. Addition of pentane (10—15 cm<sup>3</sup>, -20 °C) resulted in the precipitation of a mixture of (1) and  $[PtCl_2\{\eta^2-MeO_2C(CH)_2CO_2Me\}-(Bu^tNCHCHNBu^t)]$  which was filtered off and washed with pentane (3 × 5 cm<sup>3</sup>). From the filtrate, dimethyl fumarate could be isolated. For the isolation of free dimethyl maleate from  $[PtCl_2\{\eta^2-MeO_2C(CH)_2CO_2Me\}(Bu^tNCHCHN-Bu^t)]$  see below.

 $[PtCl_2(\eta^2-C_2H_4)(Bu^tNCHCHNBu^t)]$  (1) from  $[PtCl_2(\eta^2-olefin)(Bu^tNCHCHNBu^t)]$  (olefin = methyl acrylate or dimethyl maleate). A solution of  $[PtCl_2(\eta^2-olefin)(Bu^tNCH-CHNBu^t)]$  (0.2 mmol) in CDCl<sub>3</sub> was kept under a  $C_2H_4$ atmosphere (50 atm) in an autoclave for 6 h. The solution was degassed and N<sub>2</sub> bubbled through the reaction mixture. Nuclear magnetic resonance spectra were recorded directly after this work-up procedure and showed exclusively the

### TABLE 1

## $\label{eq:proton n.m.r. data for [PtX_2(\eta^2 - olefin)(RNCHCHNR)] and [PtX_2(\eta^2 - olefin)\{RN(H)CH_2CH_2N(H)R\}] complexes and a second second$

						Method of
				HC=0		prepar-
Complex	CH <sub>3</sub>	CH <sub>2</sub>	HC=N	CH3-C=0	HC=	ation •
$[PtCl_2(\eta^2 - C_2H_4)(Bu^t NCHCHNBu^t)] c (1)$	1.66(s, br)		8.63(s,br) [36]		3.50(s) [70]	A
$[PtClBr(\eta^2-C_2H_4)(Bu^tNCHCHNBu^t)]$	1.66(s, br)		8.63(s, Dr) [36]		3.57(S) [70]	B
$[PtBr_2(\eta^2-C_2H_4)(Bu^tNCHCHNBu^t)]^{\circ}$	1.66(s,br)		8.63(s), br) [36]		3.63(s) [70]	А, В
$[PtI_2(\eta^2 - C_2H_4)(Bu^*NCHCHNBu^*)]$	1.77(S)	1.00(-)	8.93(S) [38]		3.08(S) [72]	В
$[PtCi_2(\eta^2-C_2H_4)(Etme_2CNCHCHNCMe_2Et)]$	1.57(c br)	1.98(q)	8.48(S,Dr) [30]		5.40(S) [09]	A
[PtClBr(m <sup>2</sup> -C_H_)/FtMe_CNCHCHNCMe_Ft)] <sup>d</sup>	0.83(t)	1.98(a)	8 48 (s br) [36]		3 52(s) [70]	в
	1.57(s.br)	1.00(4)	0.10 (3,01) [00]		0.02(3) [10]	D
$[PtBr_{a}(n^{2}-C_{a}H_{a})(EtMe_{a}CNCHCHNCMe_{a}Et)]^{d}$	0.83(t).	1.98(a)	8.48(s.br) [36]		3.60(s) [69]	A. B
	1.57(s.br)	(1/				,
$[PtI_{2}(\eta^{2}-C_{2}H_{4})(EtMe_{2}CNCHCHNCMe_{2}Et)]$	0.93(t),	2.05(q)	8.47(s) [37]		3.88(s) [72]	в
	1.68(s)					
$[PtCl_2(\eta^2-CH_2=CHCO_2Me)(Bu^tNCHCHNBu^t)]$	1.63(s)		8.70(s) [40]	3.68(s)	3.43(d) [70],	в
					3.98(d) [64],	
			a aa() <b>f</b> (aa	0.0=()	4.75(dd) [84]	_
$[PtCl_2(\eta^2-CH_2=CHCO_2Me)(EtMe_2CNCHCHNCMe_2Et)]$	0.87(t),	2.02(q)	8.63(s) [40]	3.67(s)	3.38(d),	в
	1.60(s),				3.93(d),	
(D(C) ( A .: C II (CO MA) )(D INCHCUND I))	1.64(s)		0 59/-> 5407	0 70(-)	4.67(dd)	ъ
$[PtCl_2(\eta^*-cls-C_2H_2(CO_2Me)_2)(Du^*NCHCHNDu^*)]$	1.07(S)		8.73(S) [42] 9.05(c) [15] c	3.70(S)	4.00(S) [77]	а а
$[PtCl_2(\eta^2-cis-C_2\Pi_2(CO_2Me)_2)(Du^2N(\Pi)C\Pi_2C\Pi_2N-(II)P_2+t)]$	1.46(S)		3.05(S) [15] *	3.71(S)	4.21(S) [75]	Б
(П) Du' {]						
$[PtCl_{2} \{\eta^{2} - OCCH = CHC(O)O\}(Bu^{t}NCHCHNBu^{t})]$	1.67(s)		9.08(s) [50]		4.57(s) [74]	в
$[P_{1}]$ $(*^{2}-OCCH=CHC(O)O)(F_{1}M_{e}CNCHCHNCM_{e}F_{1})]$	0.87(+)		9 27(s) [50]		4 65(s) [76]	в
	1.66(s)		J.27(3) [00]		4.00(3) [10]	D
[PtCl <sub>a</sub> (n <sup>2</sup> -CH <sub>a</sub> =CHCHO)(Bu <sup>t</sup> NCHCHNBu <sup>t</sup> )]	1.67(s.br)		8.70(s.br) [4]]	9.73(s)	3.68(d).	в
	,				4.25(d)	_
[PtCl <sub>2</sub> ( $\eta^2$ -CH <sub>2</sub> =CHCO <sub>2</sub> Me){EtMe <sub>2</sub> CN(H)CH <sub>2</sub> CH <sub>2</sub> N-	0.93(t),	1.80(q)	3.00(s,br) [16] *	3.68(s)	3.33(d) [68],	в
$(H)CMe_2Et\}]$	1.37(s),			.,	3.98(d),	
	1.42(s),				4.67(dd)	
$[PtCl_2(\eta^2-CH_2=CHPh)(Bu^tNCHCHNBu^t)]$	1.58(s)		8.57(s) [38]		3.84(d) (70),	А, В
					4.51(d) [62],	
DICL ( ) C II M VD INCLUED IN	1.00()		0.00/-> 5/11		5.78(dd) [82]	
$[PtCl_2(\eta^*-cis-C_2H_2Me_2)(Bu^*NCHCHNBu^*)]$	1.60(s)		8.80(s) [41]		5.22(m) [72]	A, B
$ PtCl_2(\eta^*-trans-C_2\Pi_2Ne_2)(Du^*NCHCHNBu^*) $	1.02(S)		8.00(S) [41]		$\mathbf{0.3(m)} \ [02]$	А, В

<sup>a</sup>  $\delta$  In p.p.m., values of <sup>a</sup>  $J(^{195}\text{Pt}^{-1}\text{H})$  in Hz are given in square brackets; q = quartet. <sup>b</sup> Route A: directly from K[PtX<sub>3</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>)] or K[PtX<sub>3</sub>( $\eta^2$ -Olefin)]; B by ligand exchange, see also the Scheme. <sup>c,d</sup> Spectra obtained of a 1:1 mixture of [PtCl<sub>3</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>)] (RNCHCHNR)] and [PtBr<sub>2</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>)(RNCHCHNR)] with R = Bu<sup>t</sup> (c) or R = CMe<sub>2</sub>Et (d). <sup>e</sup> H<sub>2</sub>C-N-.

## J.C.S. Dalton

characteristic resonance patterns of (1) and the free olefin, see Table 1.

*Physical Measurements.*—Hydrogen-1 n.m.r. spectra were recorded on a Varian T-60 or XL-100 (Fouriertransform) spectrometer with SiMe<sub>4</sub> as internal standard; <sup>13</sup>C n.m.r. spectra were obtained on a Varian CFT-20 with SiMe<sub>4</sub> as internal standard. Infrared spectra were measured on a Beckman 4250 spectrophotometer.

#### RESULTS

Exchange of  $\sigma$ -N- and  $\sigma$ , $\sigma$ -N,N'-bonded Ligands.—The general route for the preparation of trans-[PtX<sub>2</sub>( $\eta^2$ -olefin)L] complexes comprises the reaction of Zeise's salt or Zeise's dimer with the appropriate ligand L. When L is bidentate [e.g. RNCHCHNR, RN(H)CH<sub>2</sub>CH<sub>2</sub>N(H)R, R'NC<sub>5</sub>H<sub>3</sub>CH=NR, bipy, or dmphen] this route affords the five-co-ordinate complex [PtCl<sub>2</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>)(L-L)] containing the olefin and the bidentate ligand in the equatorial plane of a trigonal-bipyramidal array.

In the course of our study we found new alternative routes for the preparation of five-co-ordinate platinum-olefin complexes. First, the reaction of four-co-ordinate *trans*-[PtCl<sub>2</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>)(NC<sub>5</sub>H<sub>2</sub>Me<sub>3</sub>-2,4,6)] with one of the  $\sigma$ , $\sigma$ -N,N' ligands in a 1:1—1:3 molar ratio resulted in formation of [PtCl<sub>2</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>)(L-L)] in almost quantitative yields. Secondly, the reaction starting from five-coordinate (1) with RNCHCHNR, RN(H)CH<sub>2</sub>CH<sub>2</sub>N(H)R, R'NC<sub>5</sub>H<sub>3</sub>CH=NR, bipy, and dmphen likewise resulted in quantitative formation of the five-co-ordinate complexes [PtCl<sub>2</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>)(L-L)].

These substitution reactions were carried out in dichloromethane, acetone, or toluene using a 1:1-1:3 platinumligand molar ratio. The five-co-ordinate complexes formed reactions could readily be followed by n.m.r. spectroscopy. This is illustrated by two examples.

The substitution reaction of  $[PtCl_2(\eta^2-C_2H_4)(tmpy)]$  with Bu<sup>t</sup>NCHCHNBu<sup>t</sup> (3:4 molar ratio) in CDCl<sub>3</sub> at -40 °C followed by <sup>1</sup>H n.m.r. revealed a concomitant disappearance of  $[PtCl_2(\eta^2-C_2H_4)(tmpy)]$  and free Bu<sup>t</sup>NCHCHNBu<sup>t</sup> ligand, and appearance of the five-co-ordinate complex (1) and accompanying exchanged tmpy ligand. The resonance patterns belonging to each of the components of the equilibrium mixture remained sharp while chemical-shift and coupling data [ ${}^{n}J({}^{196}Pt^{-1}H)$ ] were similar to the data obtained for the pure compounds. This indicates that intermolecular exchange between the species is slow on the n.m.r. time scale, but that equilibrium is attained rapidly on the laboratory time scale.

The <sup>13</sup>C n.m.r. spectra of a l : l molar mixture of (l) and EtMe<sub>2</sub>CNCHCHNCMe<sub>2</sub>Et in CDCl<sub>3</sub> (34 °C) showed four sharp resonance patterns corresponding to the respective components of the equilibrium mixture {*i.e.* Bu<sup>t</sup>NCHCHN-Bu<sup>t</sup>, EtMeCNCHNCMe<sub>2</sub>Et, (l), and [PtCl<sub>2</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>)(EtMe<sub>2</sub>-CNCHCHNCMe<sub>2</sub>Et)]}.

While  $MeNC_5H_3CH=NR$  and bipy are neither substituted by RNCHCHNR nor  $RN(H)CH_2CH_2N(H)R$ ,  $MeNC_5H_3$ -CH=NR is irreversibly substituted by bipy. This was concluded from the reaction of  $[PtCl_2(\eta^2-C_2H_4)(MeNC_5H_3-CH=NR)]$ , which is stable in solution for several days, with bipy followed by <sup>1</sup>H n.m.r. spectroscopy. Slow formation of free  $MeNC_5H_3CH=NCMe_2Et$ , as well as unco-ordinated  $C_2H_4$ , pointed to the occurrence of the reaction sequence given in the Scheme.

One of the reaction products  $[PtCl_2(\eta^2-C_2H_4)(bipy)]$ , which is unstable in solution, has been studied in detail by Natile and co-workers.<sup>10</sup> In our reaction a mixture of the red and yellow forms.<sup>11</sup> of the [PtCl<sub>2</sub>(bipy)] complex was obtained.

$$[PtCl_{2}(\eta^{2}-C_{2}H_{4})(tmpy)] \xrightarrow{(1)}_{\{fast\}(ii)} [PtCl_{2}(\eta^{2}-C_{2}H_{4})(RNCHCHNR)] \xrightarrow{(iii)}_{(fast)(1)} [PtCl_{2}(\eta^{2}-C_{2}H_{4})\{RN(H)CH_{2}CH_{2}N(H)R\}]$$

$$(iv) \qquad (iv) \qquad (iv)$$

$$[PtCl_2(bipy)] \xrightarrow{(vi)}_{(fast)} [PtCl_2(\eta^2 - C_2H_4)(bipy)] \xrightarrow{(v)}_{(slow)} [PtCl_2(\eta^2 - C_2H_4)(NC_5H_3CH=NR)]$$

were characterized by comparison of their <sup>1</sup>H and <sup>13</sup>C n.m.r. and i.r. spectra with those of authentic samples obtained via the Zeise's salt route. Clear advantages of the two alternative methods over this latter route are the excellent solubilities of the four- and five-co-ordinate starting materials, absence of KCl or  $H_2O$  in the reaction mixture, and the easy removal of the soluble and volatile 2,4,6-trimethylpyridine or Bu<sup>t</sup>NCHCHNBu<sup>t</sup> from the final mixtures. The reactions, which were virtually instantaneous at 25 °C (slow at -50 °C), were reversible when the L-L ligand in the five-co-ordinate complexes was RNCHCHNR or RN(H)CH<sub>2</sub>CH<sub>2</sub>N(H)R.

Addition of excess of either unidentate 2,4,6-trimethylpyridine (tmpy) or bidentate R'NCHCHNR' or R'N(H)CH<sub>2</sub>-CH<sub>2</sub>N(H)R' to a solution of [PtCl<sub>2</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>) (L-L)] afforded a quantitative substitution of the  $\sigma$ , $\sigma$ -N,N'-bonded ligand. For L-L = R'NC<sub>5</sub>H<sub>3</sub>CH=NR or bipy no substitution was observed even when a large excess of tmpy, R'NCHCHNR', or R'N(H)CH<sub>2</sub>CH<sub>2</sub>N(H)R' was used. These substitution Exchange of the Halogen Atoms.—The 1:6 molar ratio reactions of  $[PtCl_2(\eta^2-C_2H_4)(L-L)]$  [L-L = RNCHCHNR or  $RN(H)CH_2CH_2N(H)R]$  with NaI in acetone afforded ultimately (24 h) orange-red  $[PtI_2(\eta^2-C_2H_4)(L-L)]$ . The displacement of the Cl atoms by I was established by the observation of the characteristic Pt-X absorptions in the i.r. spectra of the compounds and the chemical-shift position of the ethylene resonance in the <sup>1</sup>H n.m.r. spectrum [v(Pt-I)172 cm<sup>-1</sup>,  $\delta(CH_2=CH_2)$  3.68 p.p.m., <sup>2</sup>J(<sup>195</sup>Pt-<sup>1</sup>H) 72 Hz], see also Table 1.

The reaction with NaBr is considerably slower and requires repeated removal of the NaCl-NaBr precipitate followed by addition of fresh NaBr. A better route for obtaining  $[PtBr_2(\eta^2-C_2H_4)(L-L)]$  appeared to be the reaction of K $[PtBr_3(\eta^2-C_2H_4)]$  with L-L { $[PtBr_2(\eta^2-C_2H_4)(Bu^tNCH-CHNBu^t)]$ , v(Pt-Br) 256 cm<sup>-1</sup>}.

In order to exclude the possibility that these halogenexchange reactions occur via intermediate formation of Na[PtX<sub>3</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>)] the 1:1 reaction of [PtCl<sub>2</sub>( $\eta^2$ -C<sub>2</sub>H<sub>4</sub>)-

TABLE 2

$Carbon-10$ m.m.r. data for $[1 CX_0/\eta]$ -otenin/(Du NOITOIINDU)) complexes	Carbon-13 n.m.r.	data for [PtX	L <sub>2</sub> (η <sup>2</sup> -olefin)(Bu <sup>t</sup> N	[CHCHNBu <sup>t</sup> )]	complexes a
--	------------------	---------------	---	--------------------------	-------------

	×2\11 01011	multiple refite		mpienes		
Complex	$CH_3$	C-N	C=N	C=C	O=CCH <sub>3</sub>	C=O
$[PtCl_{2}(\eta^{2}-C_{2}H_{4})(Bu^{t}NCHCHNBu^{t})]^{b}$ (1)	30.45	64.31(24)	156.69	37.98 (297)		
$[PtClBr(\eta^2 - C_2H_4)(Bu^tNCHCHNBu^t)]^{b}$	30.87	64.31(24)	157.14	36.78 (286)		
$[PtBr_2(\eta^2 - C_2H_4)(Bu^t NCHCHNBu^t)]^b$	31.27	64.31 (24)	157.59	35.46 (295)		
$[PtI_2(\eta^2 - C_2H_4)(Bu^t NCHCHNBu^t)]$	32.17	65.05 (24)	162.14	33.92 (294)		
$[PtCl_2(\eta^2-CH_2=CHCO_2Me)(Bu^tNCHCHNBu^t)] - 16 \ ^{\circ}C$	29.43,	64.71,	156.89,	33.68 (289),	50.97	171.38
	30.35	65.17(24)	158.28	38.54 (317)		
40 °C	39.29	64.95 (24)	157.68	34.64 (290),	50.50	17.1.21
				39.40 (321)		
$[PtCl_2{\eta^2-cis-C_2H_2(CO_2Me)_2}(Bu^tNCHCHNBu^t)]$	29.94	65.87	158.43	37.55 (340)	51.32	168.71
$f \in I_{n}$ as a manual of $n I/(195Dt - 13C)$ are given in perenth		dad as CDCL	antina h	Constant abtained	1	inture of

<sup>a</sup>  $\delta$  In p.p.m. values of <sup>a</sup> $J(^{196}Pt^{-13}C)$  are given in parentheses, recorded as CDCl<sub>3</sub> solutions. <sup>b</sup> Spectra obtained of a 1 : 1 mixture of  $[PtCl_2(\eta^2-C_2H_4)(Bu^tNCHCHNBu^t)]$  and  $[PtBr_2(\eta^2-C_2H_4)(Bu^tNCHCHNBu^t)]$  in CDCl<sub>3</sub> after 15 d.

(RNCHCHNR)] with  $[\text{PtBr}_2(\eta^2 - C_2H_4)(\text{RNCHCHNR})]$  (R = Bu<sup>t</sup> or CMe<sub>2</sub>Et) in CDCl<sub>3</sub> was followed by <sup>1</sup>H and <sup>13</sup>C n.m.r. spectroscopy. (The various halogen complexes have distinctly different resonance patterns, see Tables 1 and 2.)

In addition to the pattern of  $[PtCl_2(\eta^2-C_2H_4)-(RNCHCHNR)]$  and  $[PtBr_2(\eta^2-C_2H_4)(RNCHCHNR)]$ , a new set of resonances was observed which was assigned to the mixed complex  $[PtClBr(\eta^2-C_2H_4)(RNCHCHNR)]$ , see Figure 1.



FIGURE 1 Proton n.m.r. spectrum of a mixture of  $[PtX_2(\eta^2-C_2H_4)(Bu^tNCHCHNBu^t)]$  (X = Cl or Br) in CDCl<sub>3</sub> after standing for 15 d. Besides  $[PtCl_2(\eta^2-C_2H_4)(Bu^tNCHCHNBu^t)]$ and  $[PtBr_2(\eta^2-C_2H_4)(Bu^tNCHCHNBu^t)]$ , the mixed-halide product  $[PtClBr(\eta^2-C_2H_4)(Bu^tNCHCHNBu^t)]$  is also present in solution.  ${}^{n} f({}^{195}Pt^{-1}H)$  is indicated by asterisks

Olefin Exchange.—The number of  $[PtX_2(\eta^2-\text{olefin})L]$  and  $[PtX_2(\eta^2-\text{olefin})(L-L)]$  complexes which are directly available from the reaction of olefin analogues of Zeise's salt with L-L is restricted because only a few stable K[PtX\_3(\eta^2-\text{olefin})] complexes are available. The last species have to be prepared (i) via the exchange of  $C_2H_4$  in Zeise's salt by the respective olefin or (ii) via a reaction of olefin with K<sub>2</sub>[PtCl<sub>4</sub>] (catalyzed by SnCl\_2·nH\_2O): K[PtCl\_3(\eta^2-C\_2H\_4)] + olefin  $\longrightarrow$  K[PtCl\_3(\eta^2-\text{olefin})] + C\_2H\_4.

During our study we found that direct substitution of the ethylene by olefin in five-co-ordinate  $[PtX_2(\eta^2-ethylene)(L-L)]$  was also possible. In Table 1 it has been indicated by which routes the compounds are accessible. These results show that this direct olefin exchange depends on the nature of the olefin and the chelating coligand. In general, exchange was observed only with RNCHCHNR and RN(H)-

 $CH_2CH_2N(H)R$  as a coligand and with certain mono-(styrene, methyl acrylate, or acrylaldehyde) or di-substituted (*cis-* and *trans-*but-2-ene, dimethyl maleate, or maleic anhydride) olefins.

Reaction of (1) with stilbene, dimethyl fumarate, crotonaldehyde, or  $CH_2=C(Me)CHO$  was not observed. In these cases the starting material could be recovered almost quantitatively after reaction periods of up to 170 h. Likewise, the  $[PtCl_2(\eta^2-C_2H_4)(MeNC_5H_3CH=NCMe_2Et)]$  complex appeared to be unreactive with respect to exchange with, for example, methyl acrylate or maleic anhydride.

The study of the olefin-ethylene exchange in bipy complexes was hampered by solution instability.

The  $[PtCl_2(\eta^2-olefin)(L-L)]$  complexes were characterized by <sup>1</sup>H and <sup>18</sup>C n.m.r. and i.r. spectroscopy (see Tables 1 and 2) using the assignment criteria emerging from our detailed structural investigation of five-co-ordinated ethylene, butene, and styrene complexes  $[PtX_2(\eta^2-olefin)(L-L)]$  [L-L= RNCHCHNR, RN(H)CH<sub>2</sub>CH<sub>2</sub>N(H)R, or R'NC<sub>5</sub>H<sub>3</sub>CH= NR].<sup>4,5</sup> In several cases the complexes were compared with samples of the complexes obtained *via* other routes. The styrene complexes, for example, were also prepared starting from K[PtCl<sub>3</sub>( $\eta^2$ -styrene)]. It appeared that the new complexes (olefin = methyl acrylate, maleic anhydride, dimethyl maleate, and acrylaldehyde) also have a trigonal-bipyramidal structure indicating that substitution of olefins takes place with retention of configuration.

#### DISCUSSION

The present results show that the  $[PtX_2(\eta^2 \text{-olefin})(L-L)]$ [RNCHCHNR or RN(H)CH<sub>2</sub>CH<sub>2</sub>N(H)R] complexes can undergo ligand-displacement reactions of all three types of ligands with retention of the special arrangement of the ligands in the trigonal-bipyramidal array. Exchange of  $C_2H_4$  in  $[PtCl_2(\eta^2-C_2H_4)(RR'NCH_2CH_2NRR')]$  [R =  $CH_3$ ,  $R' = (R) - CH(CH_3)C_6H_5$ ] was already established, using labelled <sup>14</sup>C<sub>2</sub>H<sub>4</sub>, by Panunzi<sup>12</sup> and co-workers.<sup>7</sup> The mechanism of these displacement reactions has to account for the following general observations. The rate of ligand displacement, which is in all cases slow on the n.m.r. time scale, follows the order halide < olefin < Ndonor ligands, while the reactions strongly depend on the nature of the bidentate N-donor ligand. Whereas with RNCHCHNR and RN(H)CH<sub>2</sub>CH<sub>2</sub>N(H)R the displacement is a reversible process, exchange of these N-donor ligands with bipy or pyridinecarbaldimines appeared to be irreversible. Finally, the occurrence of ligand displacement via an associative process as the first step seems unlikely because this would require rehybridization of the platinum(II) orbitals in order to allow increase of the co-ordination number around the platinum centre from five (18-electron)  $[PtX_2(\eta^2-olefin)(L-L)]$  to six (20 electron) in  $[PtX_2(\eta^2-olefin)(L-L)L']$ .

Halogen-Halogen Exchange.—An important aspect of the halogen-halogen exchange reaction of  $[PtX_2(\eta^2-C_2H_4)-(L-L)]$  is the observation that no *trans*-to-*cis* isomerization occurs. This can be explained on the basis of the strong site preference of the N-donor ligand for equatorial sites in the trigonal-bipyramidal array. As a consequence, the olefin will reside in the equatorial plane because of the mutual stabilizing effect of these ligands with the platinum centre (*cf.* the M-olefin bonding scheme of Dewar, and Chatt and Duncanson).<sup>4,13,14</sup>

It is known that, in the presence of excess of NaX,  $[PtX_3(\eta^2-C_2H_4)]^-$  undergoes fast halide exchange.<sup>1</sup> In principle, formation of a  $[PtX_3(\eta^2-C_2H_4)]^-$  species from  $[PtX_{2}(\eta^{2}-C_{2}H_{4})(L-L)]$  by displacement of the ligand L-L is possible. Subsequent association of L-L with concomitant Pt-X bond cleavage could explain the halogen exchange observed for  $[PtCl_2(\eta^2-C_2H_4)(RNCHCHNR)]$ and  $[PtCl_2(\eta^2-C_2H_4)\{RN(H)CH_2CH_2N(H)R\}]$  with NaBr or NaI. This process requires complete loss of L-L. Moreover, the  $[PtX_3(\eta^2-C_2H_4)]^-$  intermediate could not explain the scrambling of halogens in the reaction of  $[PtCl_2(\eta^2-C_2H_4)(RNCHCHNR)]$  with its bromo-analogue. Therefore we prefer to describe this exchange reaction by initial Pt-X bond dissociation leading to cationic [PtX- $(\eta^2 - C_2 H_4)(L-L)$ ]X intermediates.

Recently Maresca *et al.*<sup>10</sup> found in their study of the solvent-dependent decomposition of  $[PtCl_2(\eta^2-C_2H_4)-(L-L)]$  [L-L = N, N, N', N'-tetramethylethylenediamine (tmen) or bipy] complexes that stable intermediates  $[PtCl(\eta^2-C_2H_4)(L-L)]X$  were formed. Complexes of this type were stabilized with suitable counter ions  $(ClO_4^{-})$  and the structure of  $[PtCl(\eta^2-C_2H_4)(tmen)][ClO_4]$  was determined by X-ray methods <sup>15</sup> (see Figure 2).



FIGURE 2 Crystal structure of  $[PtCl(\eta^2-C_2H_4)(tmen)][ClO_4]$  (ref. 15)

However, our attempts to prepare stable  $[PtX(\eta^2 - olefin)(RNCHCHNR)]X$  complexes by treating (1) with Na[BF<sub>4</sub>], Tl[PF<sub>6</sub>], or  $[NH_4][PF_6]$  failed, although we did observe decomposition  $(Pt-\eta^2-C_2H_4 \text{ bond dissociation})$ . With silver salts (Ag[NO<sub>3</sub>], Ag[BF<sub>4</sub>], or Ag[O<sub>3</sub>SCF<sub>3</sub>]) by contrast, rapid Bu<sup>t</sup>NCHCHNBu<sup>t</sup> transfer from (1) to Ag<sup>+</sup> did occur.

On the basis of the results of Maresca *et al.*,  $^{10,15}$  a plausible route for the halogen-halogen exchange would

# J.C.S. Dalton

be formation of a  $[PtX(\eta^2-C_2H_4)(L-L)]^+$  cation which can then interact with neutral  $[PtX_2(\eta^2-C_2H_4)(L-L)]$  via one of the axial halogen ligands. This would result in formation of a binuclear cationic intermediate  $[(L-L)(\eta^2-C_2H_4)ClPt-Cl-PtBr(\eta^2-C_2H_4)(L-L)]^+$  which can dissociate to produce either the starting species or the exchanged products. In this proposed mechanism the N-donor ligand remains  $\sigma, \sigma - N, N'$ -chelate bonded during the whole displacement reaction. Likewise, olefin dissociation is not possible since in separate experiments we found that this leads irreversibly to cis-[PtCl\_2(L-L)] complexes.

Olefin and N-Donor Ligand Displacement.—Since both the olefin- and the ligand-displacement reactions are much faster than the halogen-exchange reactions it seems unlikely that these displacements proceed via a cationic intermediate  $[PtCl(\eta^2-olefin)(L-L)]^+$ . Furthermore, we have shown that the platinum-olefin bonds in the five-coordinate  $[PtCl_2(\eta^2-olefin)(L-L)]$  [L-L = RNCHCHNR, $R'NC_5H_3CH=NR$ , or  $RN(H)CH_2CH_2N(H)R]$  complexes are inert on the n.m.r. time scale with the coupling constants and chemical-shift positions in both <sup>1</sup>H and <sup>13</sup>C n.m.r. spectra being independent of the concentration of excess of olefin or L-L.

On the basis of these considerations, the olefin and L-L displacement reactions can be better described by a process which involves Pt-N bond dissociation as the initial step. This would imply a change of the chelate bonding mode  $(\sigma, \sigma \cdot N, N')$  of the N-donor ligand L-L to a  $\sigma \cdot N$  unidentate mode. By this process a free coordination site is created *via* which either an olefin or an N-donor ligand (L or L-L) can enter the co-ordination sphere of the platinum atom. The so formed five-coordinate intermediate (Figure 3) produces either the starting complex or the exchanged product. This picture is consistent not only with the N-donor ligands (L, L-L) but also with the olefin-displacement reactions (see Figure 3).

In previous papers extensive evidence for the occurrence of reversible  $\sigma, \sigma-N, N'$  chelate to  $\sigma-N$  unidentate processes in square-planar  $[PtX_{2-n}(PR_3)_{1+n}(L-L)]X_n$  (n = 0 or 1) complexes has been put forward.<sup>16-19</sup> In these intramolecular rearrangements the bidentate N-donor ligand changes its point of attachment to the Pt atom (head-to-tail rearrangement) via five-co-ordinate intermediates. The kinetics of this fluxional behaviour depend strongly on the stereochemistry of the N-donor ligands. A common feature of these ligands shown in Figure 4 is that in the syn conformation the electron pairs on the nitrogen atoms are suitably positioned for  $\sigma, \sigma$ -N,N' chelation. A difference, however, is the flexibility of the skeleton for rotation around the central C-C bond in the unidentate co-ordination mode and the steric consequences of such a rotation with respect to possible interaction of the metal atom with the substituents connected to the  $\beta$ -imine C atom. Considering the series shown in Figure 4, it is obvious that the tendency towards unidentate Pt-N interaction will increase on going from bipy to RN(H)CH<sub>2</sub>CH<sub>2</sub>N(H)R. This is indeed the case. For example, Dixon 17 studied [PtCl-



FIGURE 3 Mechanism proposed for olefin exchange in  $[PtCl_2(\eta^2-olefin)(L-L)]$  complexes. Initial Pt-N bond dissociation provides a four-co-ordinate species (not shown) which has a free co-ordination site for the incoming olefin. Similar intermediates are proposed for the bidentate N-donor ligand exchange when olefin is replaced by L-L' for the incoming ligand and olefin is replaced by RNCHCHNR as leaving group

 $(PR_3)_2(L-L)]X$  (L = bipy or phen) complexes and found that the slow exchange limit was not reached at -80 °C. However, <sup>1</sup>H and <sup>13</sup>C n.m.r. studies of  $[PtCl_2(PR_3)-(RNCHCHNR)]$  having the less sterically hindered ligand (see above) revealed that in the slow-exchange limit (ca. -40 °C) the RNCHCHNR ligand is unidentate with a trans ligand conformation as is shown in Figure 4. In this case the  $\beta$ -imine H atom (CH=N) can be favourably positioned with respect to the co-ordination plane.<sup>16</sup> These conclusions were confirmed by a study of the <sup>15</sup>Nlabelled derivatives.<sup>18</sup> Above -40 °C, rapid intramolecular N,N' site exchange took place. It should be mentioned that in these complexes a phosphine ligand acts as coligand instead of an olefinic ligand.

Maresca *et al.*<sup>19</sup> showed that  $[PtCl_2(\eta^2-C_2H_4)\{R^1R^2N-(CH_2)_nNR^1R^2\}]$  complexes for n = 3 exist only at low temperature in the five-co-ordinate form; at higher temperature they have to be formulated as four-co-ordinate platinum compounds with unidentate  $R^1R^2N-(CH_2)_nNR^1R^2$ .

These results provide an explanation for the observation that displacement of L<sup>-L</sup> in  $[PtCl_2(\eta^2-C_2H_4)(L^-L)]$ 



**FIGURE 4** Comparison of the  $\sigma, \sigma-N, N'$ -chelating [(a)-(d)] and  $\sigma-N$  unidentate [(a')-(d'), (c'')] bonding mode of several bidentate *N*-donor ligands. Structure (c') is unfavourable since the pyridine *N*-donor atom is a much poorer electron-releasing atom than the alkyl imine *N* atom

# J.C.S. Dalton

by pyridine (py), RNCHCHNR, and RN(H)CH<sub>2</sub>CH<sub>2</sub>-N(H)R was reversible whereas the reactions with pyridine-2-carbaldimine or bipy appeared to be irreversible. For the latter two N-donor ligands, the formation of a  $\sigma$ -N unidentate Pt-N intermediate has a high activation barrier because both for bipy and pyridine-2-carbaldimine (which would be bonded via the imine nitrogen rather than via the pyridine N atom)  $H^3$  would be unfavourably positioned (see Figure 4) with respect to the platinum centre.

In the series RNCHCHNR, RN(H)CH<sub>2</sub>CH<sub>2</sub>N(H)R, and py the tendency towards four-co-ordination increases and for the py ligand it has been shown that only under special conditions (below -30 °C with excess of py) could a five-co-ordinate species, in low concentration, be observed by n.m.r.20

In the olefin-displacement reactions of  $[PtCl_2(\eta^2-C_2H_d)-$ (L-L)] it was only for L-L = RNCHCHNR or RN(H)- $CH_{2}CH_{2}N(H)R$  that we observed (reversible) olefin displacement. The proposed displacement of one N coordination site of the L-L ligand by the second olefin ligand is analogous to the mechanisms proposed for olefin exchange in square-planar platinum-olefin complexes. In the latter case it is always the non-olefinic ligand which is the leaving 'group.<sup>1,2</sup>

An interesting aspect of these olefin-displacement reactions is the influence of the nature of the olefin. There is no evidence for a steric influence on the course of the reaction with respect to the five-co-ordinate intermediate which most probably contains the two olefins in the equatorial co-ordination sites. Studies of Stone and co-workers<sup>21</sup> of the bonding and barriers to olefin rotation in trigonal  $[Pt^0(\eta^2-olefin)_2L]$  complexes are of interest in this respect. These complexes might be isostructural in the trigonal plane with our proposed intermediate.

It is well established that *cis*-disubstituted olefins bond more strongly to a metal centre than the corresponding trans olefin.<sup>1b,22</sup> Upon  $\eta^2$ -co-ordination of cisand trans-disubstituted olefins to the platinum centre there is a net positive energy gain, which is of greater importance for the *cis* isomer due to its particular bond oppositions. However, in complexes of the [PtCl<sub>2</sub>{ $\eta^2$ - $H_{3}C(H)C=C(H)CH_{3}(RNCHCHNR)$ ] type the *cis*- and trans-but-2-ene species are exceptional because both cis and *trans* isomers can be exchanged.

A more detailed investigation directed to the study of olefin-isomer complex formation should clarify this point.23

Conclusions.—The present results clearly show that in five-co-ordinate platinum(II) olefin complexes of the type  $[PtCl_2(\eta^2-C_2H_4)(L-L)]$  it is possible to exchange all the ligands (halide, olefin, or L-L) by intermolecular processes.

A prerequisite for reversible neutral ligand exchange in the proposed mechanism is the possibility of a (intermediate) unidentate L-L ligand. It has been shown on both stereochemical and electronic grounds that the  $\alpha$ di-imine ligand and its hydrogenated analogue, 1,2diaminoethane, possess the capability of such versatile bonding behaviour.

Moreover it is reasonable to assume, especially for the  $\alpha$ -di-imine ligand that the  $\sigma$ -N unidentate or  $\sigma$ -N,  $\sigma$ -N' bridge bonding mode [with a s-trans N=C-C=N skeleton] is important for the formation of intermediates in either stoicheiometric or catalytic processes in which such ligands are involved. Examples are C-C bond formation in the reversible coupling of two  $\alpha$ -di-imines in ruthenium(0) and molybdenum(0) carbonyl chemistry 24 and complex formation of  $\alpha$ -di-imines with alkyl aluminium (AlR<sub>3</sub>) which occurs with concomitant -C=N- bond reduction and alkyl transfer.<sup>25</sup>

Thanks are due to Professor K. Vrieze for his contributions and stimulating interest in this work. The experimental assistance of Mr. J. M. Ernsting (n.m.r.) and Mr. G. C. Schoemaker (i.r.) is gratefully acknowledged. Dr. D. M. Grove is thanked for critical reading of the manuscript.

[1/404 Received, 11th March, 1981]

#### REFERENCES

<sup>1</sup> (a) F. Basolo and R. G. Pearson, 'Mechanisms of Inorganic Reactions,' Wiley, New York, 1967; (b) F. R. Hartley, 'The Chemistry of Platinum and Palladium,' Applied Science, London,

1973.
<sup>2</sup> S. S. Hupp and G. Dahlgren, *Inorg. Chem.*, 1976, 15, 2349;
P. D. Kaplan, P. Schmidt, A. Brause, and M. Orchin, *J. Am. Chem.*

 Soc., 1969, 91, 85.
 <sup>3</sup> H. van der Poel, G. van Koten, K. Vrieze, M. Kokkes, and C. H. Stam, J. Organomet. Chem., 1979, 175, C21.

<sup>4</sup> H. van der Poel, G. van Koten, M. Kokkes, and C. H. Stam,

Inorg. Chem., 1981, in the press. <sup>5</sup> H. van der Poel and G. van Koten, Inorg. Chem., 1981, in the

Press. <sup>6</sup> L. Maresca, G. Natile, M. Calligaris, P. Delise, and L.

Randaccio, J. Chem. Soc., Dalton Trans., 1976, 2386. <sup>7</sup> A. De Renzi, B. Di Blasio, A. Saporito, M. Scalone, and A. Vitagliano, Inorg. Chem., 1980, **19**, 960.

H. van der Poel and G. van Koten, J. Organomet. Chem., 1980, 187, C17.

 O. Exner and J. M. Kliegman, J. Org. Chem., 1971, 36, 2014;
 H. Bock and H. tom Dieck, Chem. Ber., 1967, 100, 288, and refs. therein; H. van der Poel, Ph.D. Thesis, University of Amsterdam, 1981.

<sup>19</sup> L. Maresca, G. Natile, and L. Cattalani, J. Chem. Soc., Dalton Trans., 1979, 1140; L. Maresca, G. Natile, and G. Riz-zardi, Inorg. Chim. Acta, 1980, 38, 137.
 <sup>10</sup> E. Bielli, P. M. Gidney, R. D. Gillard, and B. T. Heaton, J.

Chem. Soc., Dalton Trans., 1974, 2133.

<sup>12</sup> A. De Renzi, A. Panunzi, A. Saporito, and A. Vitagliano, Gazz. Chim. Ital., 1977, 107, 549.

<sup>13</sup> M. J. S. Dewar, Bull. Soc. Chim. Fr., 1953, 18, C79; M.

<sup>14</sup> T. A. Albright, R. Hoffmann, J. C. Thibeault, and D. L. Thorn, J. Am. Chem. Soc., 1973, 103, 2939.
 <sup>14</sup> T. A. Albright, R. Hoffmann, J. C. Thibeault, and D. L. Thorn, J. Am. Chem. Soc., 1979, 101, 3801; T. Ziegler and A. Rauk, Inorg. Chem., 1979, 18, 1558.
 <sup>15</sup> A. Tiripicchio, M. Tiripicchio-Camelini, L. Maresca, G. Natile, and G. Rizzardi, Cryst. Struct. Commun., 1979, 8, 689.
 <sup>16</sup> H. van der Poel, G. van Koten, and K. Vrieze, Inorg. Chem., 1960, 10, 1145.

1980, 19, 1145.

<sup>17</sup> K. R. Dixon, Inorg. Chem., 1977, 16, 2618.

<sup>18</sup> H. van der Poel, G. van Koten, D. M. Grove, P. S. Pregosin, and K. Ostoja Starzewski, *Helv. Chim. Acta*, 1981, **64**, 1174.

<sup>19</sup> L. Maresca, G. Natile, and L. Cattalani, J. Chem. Soc., Dalton Trans., 1979, 1140.

<sup>20</sup> J. C. Chottard, D. Mansuy, and J. F. Bartoli, J. Organomet. Chem., 1974, 65, C19; F. Pesa and M. Orchin, *ibid.*, 1976, 108, 135; I. Al-Najjar and M. Green, J. Chem. Soc., Chem. Commun., 1977, 212.

<sup>21</sup> N. C. Harrison, M. Murray, J. L. Spencer, and F. G. A. Stone, J. Chem. Soc., Dalton Trans., 1978, 1337; T. Chicote, M. Green, J. L. Spencer, F. G. A. Stone, and J. Vicente, *ibid.*, 1979, 536.
<sup>22</sup> M. A. Muhs and F. T. Weiss, J. Am. Chem. Soc., 1962, 84, 4697; S. Sakaki, Theor. Chim. Acta, 1973, 30, 159.

- <sup>23</sup> H. van der Poel, G. van Koten, L. Maresca, and G. Natile, unpublished work.
  <sup>24</sup> L. H. Staal, A. Oskam, K. Vrieze, E. Roosendaal, and H. Schenk, *Inorg. Chem.*, 1979, 18, 1634.
  <sup>25</sup> J. M. Klerks, D. J. Stufkens, G. van Koten, and K. Vrieze, *J. Organomet. Chem.*, 1979, 181, 271.