

A Novel Exchange Reaction between Diselenides and Bis(*N,N*-dialkylselenocarbamoyl)selenides. Application in the Synthesis of Tetraselenafulvalene (TSF)

Ib Johannsen,^a Knud Lerstrup,^b Lars Henriksen,^c and Klaus Bechgaard^c

^a Physics Laboratory III, Technical University of Denmark, DK-2800 Lyngby, Denmark.

^b Department of Chemistry, the Johns Hopkins University, Baltimore, Maryland 21218, U.S.A.

^c Department of General and Organic Chemistry, The H. C. Ørsted Institute, DK-2100 Copenhagen, Denmark

A high yield synthesis of tetraselenafulvalene (TSF) involving a novel acid catalysed exchange reaction between diselenides and bis(*N,N*-dialkylselenocarbamoyl)selenides is reported.

The recent discovery of superconductivity in cation radical salts of tetramethyltetraselenafulvalene¹ has increased the interest in the synthesis of tetraselenafulvalenes (TSF's). Key intermediates in the synthesis of substituted TSF's are 2-oxoalkyl *N,N*-dialkyldiselenocarbamoyl esters (in the 'carbamate route').^{2,3}

Unsubstituted TSF (**1**) has so far been prepared by two routes^{4,5} (yield 5–20%) from CSe₂, which is difficult to obtain. We have been interested in preparing (**1**) by the 'carbamate' route especially since *N,N*-dimethyldiselenocarbamate salts are now more readily available from *N,N*-dimethylphosgeneiminium chloride and NaHSe or H₂Se.³ However, α-halogenoacetaldehydes (or their acetal derivatives) do not react with diselenocarbamate salts. Instead we obtained the esters (**4**) in high yields by a novel acid catalysed exchange reaction between the diselenide (**2**) and bis(*N,N*-dialkyldiselenocarbamoyl)selenides (**3a**) or (**3b**) (Scheme 1).


The carbamoylselenides (**3**) are readily obtained from the corresponding diselenocarbamate^{3,6} salts by oxidation and treatment with trimethyl phosphite.⁶ Bis(2-oxoethyl)diselenide (**7**) was prepared as shown in Scheme 2. Na₂Se₂⁷ reacted with 2-bromoacetaldehyde diethyl acetal in an ethanol-water (1:1) solution. The reaction mixture was extracted with ether, and distilled to give (**6**) as a yellow oil, b.p. 120 °C

(0.05 mmHg) (80%). The acetal was then hydrolysed with dilute formic acid and the aldehyde (**7**) was obtained by extraction with ether and by removal of the solvent.


Stoichiometric amounts of (**7**) (prepared as in Scheme 2) and (**3a**) were stirred in a CHCl₃ solution containing a catalytic amount of toluene-*p*-sulphonic acid. The precipitate of the triselenide (**5a**) was removed by filtration and the solvent was evaporated to give the carbamate ester (**4a**) as a partially crystalline orange oil.

The conversion of (**4a**) into TSF is shown in Scheme 3. (**4a**) was cyclized by slow dissolution into ice cold conc. H₂SO₄. The iminium compound (**8a**) was isolated as the PF₆[−] salt after treatment with HPF₆ in ice-water. After dissolution in CH₂Cl₂ and precipitation with ether (**8a**) was obtained as white needles, m.p. 160 °C (decomp.), 65% yield based on (**3a**). In a similar manner (**8b**) was obtained in 60% yield.


Treatment of compounds (**8**) with an excess of H₂Se in MeOH-H₂O solution followed by recrystallization of the precipitate from hexane gave the selenone (**9**).³ (**9**) was converted into TSF by treatment with P(OMe)₃.



Scheme 2. i, EtOH-H₂O, reflux, 16 h; ii, 1 M HCO₂H, 50 °C, 1 h.



Scheme 1



Scheme 3. i, 3 mm toluene-*p*-sulphonic acid in CHCl₃, room temp., 2 h; ii, conc. H₂SO₄, 0 °C, 2 h; 2 M HPF₆, 0 °C, ½ h; iii, H₂Se, MeOH-H₂O, 0 °C, 5 h; iv, P(OMe)₃, benzene, reflux, 2 h.

Thus, TSF can be prepared in 42—48% overall yield or from phosgeneiminium chloride and H_2Se or NaHSe^3 with comparable yields.

Received, 8th August 1983; Com. 1075

References

- 1 D. Jérôme, A. Mazaud, M. Ribault, and K. Bechgaard, *J. Phys. (Paris), Lett.*, 1980, **41**, L15.
- 2 K. Bechgaard, D. O. Cowan, A. N. Bloch, and L. Henriksen, *J. Org. Chem.*, 1975, **40**, 746; J. R. Andersen and K. Bechgaard, *ibid.*, p. 2016.
- 3 F. Wudl and D. Nalewajek, *J. Chem. Soc., Chem. Commun.*, 1980, 866; L. Chiang, T. O. Poehler, A. N. Bloch, and D. O. Cowan, *ibid.*, p. 866; A. Moradpour, V. Peyrussan, I. Johannsen, and K. Bechgaard, *J. Org. Chem.*, 1983, **48**, 388.
- 4 E. M. Engler and V. Patel, *J. Am. Chem. Soc.*, 1974, **96**, 7376.
- 5 M. V. Lakshmikantham and M. P. Cava, *J. Org. Chem.*, 1976, **41**, 882.
- 6 L. Henriksen, *Synthesis*, 1982, 771.
- 7 D. L. Klayman and T. S. Griffin, *J. Am. Chem. Soc.*, 1973, **95**, 197.