


This article was downloaded by: [University of Louisville]

On: 25 January 2015, At: 08:57

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Synthesis of Benzofuryl Substituted Unsymmetrical Ureas Under Microwave Irradiation

Lan-Qin Chai^a, Yony-Xi Chen^a, Wei-Peng Chen^a & Qi Ding^a

^a School of Chemical and Biological Engineering, Lanzhou Jiaotong University, Lanzhou, Gansu, P. R. China

Published online: 03 Feb 2009.

To cite this article: Lan-Qin Chai, Yony-Xi Chen, Wei-Peng Chen & Qi Ding (2009) Synthesis of Benzofuryl Substituted Unsymmetrical Ureas Under Microwave Irradiation, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 184:2, 453-459, DOI: [10.1080/10426500802176689](https://doi.org/10.1080/10426500802176689)

To link to this article: <http://dx.doi.org/10.1080/10426500802176689>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages,

and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Synthesis of Benzofuryl Substituted Unsymmetrical Ureas Under Microwave Irradiation

Lan-Qin Chai, Yony-Xi Chen, Wei-Peng Chen, and Qi Ding

School of Chemical and Biological Engineering, Lanzhou Jiaotong University, Lanzhou, Gansu, P. R. China

A series of 2-benzofuryl substituted unsymmetrical ureas were synthesized by reactions of benzofuroyl isocyanate, which was prepared from benzofuroyl azide by Curtius rearrangement, with various aromatic amines, 2-amino-5-(benzo-2-furyl)-1,3,4-thiadiazole, and 2-amino-5-aryloxymethylene-1,3, 4-thiadiazoles under microwave irradiation. Compared to conventional methods, this synthesis has the advantages of mild reaction conditions, easy handling, and high yields. The products have been characterized by analytical and spectral (IR and ^1H NMR) data.

Keywords 1,3,4-thiadiazoles; curtius rearrangement; isocyanate; microwave irradiation (MWI); unsymmetrical urea

INTRODUCTION

In recent years, the use of microwave irradiation to promote reactions has received considerable attention, and dramatic rate enhancements have been reported.¹ Organic cyclization reactions yielding heterocycles under microwave irradiation have attracted the attention of chemists.^{2,3} Examples of such applications are the syntheses of unsymmetrically substituted ureas. Unsymmetrical ureas are widely used as herbicides, pesticides, plant growth regulators, and medicinal intermediates.^{4–6} Ureas bearing heterocyclic substituents, for example 1,3,4-thiadiazole, have been shown to exert anti-inflammatory,^{7–9} anti-bacterial,¹⁰ and anti-convulsant activities.¹¹ The synthetic protocols of ureas generally utilize phosgene or phosgene-based isocyanates

Received 18 March 2008; accepted 2 May 2008.

We are thankful for the financial support from the Foundation of Lanzhou Jiaotong University (No. 06-40611) and the Innovation and Technology Fund of Lanzhou Jiaotong University (No. DXS-2007-007).


Address correspondence to Lan-Qin Chai, School of Chemistry and Biological Engineering, Lanzhou Jiaotong University, Lanzhou, 730070, Gansu, P.R. China. E-mail: chailq@mail.lzjtu.cn

as starting materials,^{12,13} both of which are toxic or unstable. These methods also involve longer reaction times. Therefore, it is necessary to develop phosgene-free and straight-toward routes for unsymmetrical ureas.

These reasons prompted us to develop an environmentally benign methodology to synthesize some new series of compounds bearing both urea and 1,3,4-thiadiazole moieties, with the objective to investigate the properties and structure–activity relationships of these new compounds and to obtain new biologically active compounds.

RESULTS AND DISCUSSION

We herein report a fast and efficient method for the preparation of a series of unsymmetrical ureas. As described in Scheme 1, N-aryl-N'-(benzofuran-2-yl) ureas (**3a–k**), N-benzofuran-2-yl-N'-[5-(benzofuran-2-yl)-1,3,4-thiadiazol-2-yl] urea (**4a**), and N-benzofuran-2-yl-N'-(1,3,4-thiadiazol-2-yl) ureas (**5a–i**) were synthesized by reactions of benzofuran isocyanate with various aromatic amines, 2-amino-5-(benzo-2-furyl)-1,3,4-thiadiazole, and 2-amino-5-aryloxymethylene-1,3,4-thiadiazoles, respectively, under microwave irradiation. Benzofuran-2-yl isocyanate was prepared by treating benzofuran-2-carboxylic acid with sodium azide and ethyl chloro-carbamate in the presence of triethylamine followed by Curtius rearrangement.¹⁴



SCHEME 1

To investigate the effects of microwave irradiation, all the reactions were performed in an oil bath at 120°C. When compared to classical heating, the reactions performed under microwave irradiation are at least 30 times faster and proceed with high yields. The results obtained are reported in Table I.

TABLE I Yields, Reaction Time, Melting Points, and Elemental Analyses of Compounds 3a–k, 4a, and 5a–i

	Ar	Yield (%)		Reaction Time		M.p. °C lit.	Elemental analysis (%) found (calcd.)			
		MWt ^a	Reflux ^b	MWt	Reflux (hour) ^b		C	H		N
3a	C ₆ H ₅	84	74	10	5	232–233 (180–181) ^[15c]	71.54 (71.42)	4.91 (4.79)		11.23 (11.10)
3b	2-CH ₃ C ₆ H ₄	82	77	8	4	244–246	71.89 (72.17)	5.66 (5.30)		10.48 (10.52)
3c	4-CH ₃ C ₆ H ₄	85	78	8	4	250–252 (214–215) ^[15c]	71.83 (72.17)	5.59 (5.30)		10.67 (10.52)
3d	2-NO ₂ C ₆ H ₄	75	64	14	8	224–226	60.40 (60.61)	4.06 (3.73)		14.09 (14.14)
3e	3-NO ₂ C ₆ H ₄	71	61	14	8	226–228	60.84 (60.61)	4.01 (3.73)		14.28 (14.14)
3f	4-NO ₂ C ₆ H ₄	72	62	14	8	>300	60.78 (60.61)	3.94 (3.73)		14.39 (14.14)
3g	2-ClC ₆ H ₄	79	69	10	6	240–241	62.67 (62.84)	4.01 (3.87)		9.91 (9.77)
3h	4-ClC ₆ H ₄	80	70	10	6	256–258 (216–218) ^[15c]	62.61 (62.84)	4.06 (3.87)		9.95 (9.77)
3i	4-BrC ₆ H ₄	81	71	12	7	222–224 (230–232) ^[15c]	54.24 (54.40)	3.57 (3.35)		8.63 (8.46)
3j	1-Naphthyl	80	70	12	7	252–254	75.72 (75.48)	4.80 (4.67)		9.46 (9.27)
3k	2-Naphthyl	78	74	12	7	276–278	75.69 (75.48)	4.86 (4.67)		9.39 (9.27)
4a	benzofuran-2	78	70	25	14	280–282	60.85 (60.63)	3.38 (3.21)		15.11 (14.89)
5a	C ₆ H ₅	80	73	25	14	288–290	59.25 (59.01)	3.93 (3.85)		15.45 (15.29)
5b	4-CH ₃ OC ₆ H ₄	82	78	20	12	>300	57.64 (57.57)	4.23 (4.07)		14.28 (14.13)
5c	2-CH ₃ C ₆ H ₄	80	75	20	13	268–270	60.14 (59.99)	4.46 (4.24)		14.91 (14.73)
5d	3-CH ₃ C ₆ H ₄	73	68	22	13	246–248	60.21 (59.99)	4.40 (4.24)		14.95 (14.73)
5e	4-CH ₃ C ₆ H ₄	76	70	22	13	272–274	60.18 (59.99)	4.38 (4.24)		14.89 (14.73)
5f	4-ClC ₆ H ₄	72	66	25	15	290–292	53.78 (53.94)	3.38 (3.27)		13.79 (13.98)
5g	2,4-Cl ₂ C ₆ H ₃	68	64	25	15	294–296	49.88 (49.67)	2.93 (2.78)		12.76 (12.87)
5h	4-NO ₂ C ₆ H ₄	65	61	28	16	276–278	52.73 (52.55)	3.34 (3.19)		17.25 (17.02)
5i	1-Naphthyl	70	65	25	14	>300	63.59 (63.45)	3.98 (3.87)		13.57 (13.45)

^aIrradiated by microwave at less than 490 W; ^bHeated at 120°; ^cRecrystallized from either acetone or alcohol, recorded in an open capillary tube and are uncorrected.

We selected the synthesis of compound **3a** as a model reaction to study the effects of irradiation power and time on the yields. The best yields obtained are 84% after 10 min of irradiation with 490 W using toluene as solvent. A greater power or longer irradiation time induces a decrease in yield (only 78% with 700 W or 68% after 12 min) due to the decomposition of benzofuran-2-yl isocyanate.

In summary, the synthesis of unsymmetrical ureas has been accomplished employing the Curtius rearrangement of 2-benzofuroyl azide followed by nucleophile addition of amines to the NCO moiety under microwave irradiation. Compared to conventional thermal heating, microwave irradiation decreased the reaction time from 4–16 h to 8–28 min. The main advantages of this method are short reaction times, high yields, less byproducts, and simple handling of slashing materials and products.

EXPERIMENTAL

IR spectra were recorded using KBr pellets on a Nicolet AVATAR 360 FT-IR spectrophotometer. ^1H NMR spectra were obtained with a Bruker Avanci-D2X-200 instrument using DMSO-d_6 as solvent and TMS as internal standard. Elemental analyses were performed on a Vario E-l Elemental Analysis instrument. Melting points were determined with a XT-4 thermal apparatus and are uncorrected. Microwave irradiation was carried out in a Galanz domestic microwave oven.

Benzofuran-2-carboxylic acid,¹⁶ 2-amino-5-(benzo-2-furyl)-1,3,4-thiadiazole,¹⁷ and 2-amino-5-aryloxymethylene-1,3,4-thiadiazoles¹⁸ were prepared according to literature procedures. Aryloxy acetic acids were commercially available and used as received.

Preparation of Benzofuroyl Azide 1

A mixture of benzofuran-2-carboxylic acid (10 mmol, 1.87 g), triethylamine (11 mmol, 1.111 g), and ethyl chlorocarbamate (11 mmol, 1.194 g) in dry acetone (30 mL) was stirred at 0°C for 1 h. Then sodium azide (11 mmol, 0.715 g) dissolved in 15 mL water was added and the mixture was kept at 0°C for 7 h. After the reaction was completed (monitored by TLC), the mixture was poured onto ice. The precipitated product was separated by filtration. Yield: 96.3 %. White crystal. M.p. $110\text{--}111^\circ\text{C}$. IR (KBr, ν/cm^{-1}): 2162 ($\text{N}\equiv\text{N}$), 1703 ($\text{C}=\text{O}$), 1333 ($\text{N}=\text{N}$); ^1H NMR (DMSO-d_6) δ : 6.90–7.36 (m, 5H, benzofuran H). MS: m/z = 187. Anal. Calcd. for $\text{C}_9\text{H}_5\text{O}_2\text{N}_3$: C, 57.76; H, 2.69; N, 22.45. Found: C, 57.83; H, 2.71; N, 22.52.

TABLE II IR and ^1H NMR Spectroscopic Data for Compounds 3a–k, 4a, and 5a–i

Product	IR (KBr) cm^{-1}	^1H NMR δ /(ppm)
3a	3314, 3276 (N–H); 1649 (C=O)	9.82 (s, 1H, NH); 9.01 (s, 1H, NH); 7.95–6.49 (m, 10H, ArH & benzofuran H).
3b	3284, 3264 (N–H); 1653 (C=O)	9.79 (s, 1H, NH); 8.99 (s, 1H, NH); 7.93–6.47 (m, 9H, ArH & benzofuran H); 2.44 (s, 3H, CH_3).
3c	3282, 3274 (N–H); 1659 (C=O)	9.78 (s, 1H, NH); 9.00 (s, 1H, NH); 7.94–6.48 (m, 9H, ArH & benzofuran H); 2.43 (s, 3H, CH_3).
3d	3288, 3269 (N–H); 1646 (C=O)	9.90 (s, 1H, NH); 9.04 (s, 1H, NH); 8.11–6.54 (m, 9H, ArH & benzofuran H).
3e	3280, 3261 (N–H); 1655 (C=O)	9.87 (s, 1H, NH); 9.03 (s, 1H, NH); 8.13–6.51 (m, 9H, ArH & benzofuran H).
3f	3302, 3273 (N–H); 1654 (C=O)	9.88 (s, 1H, NH); 9.04 (s, 1H, NH); 8.21–6.60 (m, 9H, ArH & benzofuran H).
3g	3290, 3271 (N–H); 1660 (C=O)	9.80 (s, 1H, NH); 9.01 (s, 1H, NH); 7.96–6.52 (m, 9H, ArH & benzofuran H).
3h	3281, 3264 (N–H); 1648 (C=O)	9.79 (s, 1H, NH); 9.03 (s, 1H, NH); 8.09–6.51 (m, 9H, ArH & benzofuran H).
3i	3288, 3269 (N–H); 1657 (C=O)	9.82 (s, 1H, NH); 9.02 (s, 1H, NH); 7.95–6.49 (m, 9H, ArH & benzofuran H).
3j	3278, 3261 (N–H); 1665 (C=O)	9.81 (s, 1H, NH); 9.01 (s, 1H, NH); 8.01–6.51 (m, 12H, ArH & benzofuran H).
3k	3275, 3265 (N–H); 1661 (C=O)	9.80 (s, 1H, NH); 9.00 (s, 1H, NH); 8.03–6.47 (m, 12H, ArH & benzofuran H).
4a	3283, 3175 (N–H); 1658 (C=O); and 1574, 1495, 1324, 1044 (C=N–N=C–S)	9.80 (s, 1H, NH); 9.22 (s, 1H, NH); 7.95–6.86 (m, 10H, benzofuran H); 5.41 (s, 2H, CH_2O).
5a	3277, 3179 (N–H); 1667 (C=O); and 1577, 1496, 1321, 1045 (C=N–N=C–S)	9.82 (s, 1H, NH); 9.20 (s, 1H, NH); 7.96–6.89 (m, 10H, ArH & benzofuran H); 5.42 (s, 2H, CH_2O).
5b	3283, 3176 (N–H); 1661 (C=O); and 1584, 1494, 1322, 1048 (C=N–N=C–S)	9.80 (s, 1H, NH); 9.18 (s, 1H, NH); 7.95–6.87 (m, 9H, ArH & benzofuran H); 5.38 (s, 2H, CH_2O); 2.28 (s, 3H, CH_3O).

(Continued on next page)

TABLE II IR and ^1H NMR Spectroscopic Data for Compounds 3a–k, 4a, and 5a–i (Continued)

Product	IR (KBr) cm^{-1}	^1H NMR $\delta/(\text{ppm})$
5c	3280, 3174 (N–H); 1653 (C=O); and 1575, 1493, 1323, 1047 (C=N–N=C–S)	9.79 (s, 1H, NH); 9.17 (s, 1H, NH); 7.92–6.84 (m, 9H, ArH & benzofuran H); 5.37 (s, 2H, CH_2O); 2.26 (s, 3H, CH_3).
5d	3285, 3172 (N–H); 1658 (C=O); and 1583, 1495, 1325, 1049 (C=N–N=C–S)	9.81 (s, 1H, NH); 9.16 (s, 1H, NH); 7.96–6.85 (m, 9H, ArH & benzofuran H); 5.40 (s, 2H, CH_2O); 2.27 (s, 3H, CH_3).
5e	3278, 3170 (N–H); 1653 (C=O); and 1576, 1490, 1326, 1046 (C=N–N=C–S)	9.78 (s, 1H, NH); 9.14 (s, 1H, NH); 7.93–6.86 (m, 9H, ArH & benzofuran H); 5.38 (s, 2H, CH_2O); 2.24 (s, 3H, CH_3).
5f	3281, 3168 (N–H); 1662 (C=O); and 1574, 1492, 1319, 1045 (C=N–N=C–S)	9.81 (s, 1H, NH); 9.21 (s, 1H, NH); 7.95–6.88 (m, 9H, ArH & benzofuran H); 5.45 (s, 2H, CH_2O).
5g	3288, 3174 (N–H); 1652 (C=O); and 1573, 1493, 1320, 1044 (C=N–N=C–S)	9.84 (s, 1H, NH); 9.18 (s, 1H, NH); 7.98–6.92 (m, 8H, ArH & benzofuran H); 5.47 (s, 2H, CH_2O).
5h	3293, 3167 (N–H); 1654 (C=O); and 1570, 1491, 1317, 1042 (C=N–N=C–S)	9.83 (s, 1H, NH); 9.23 (s, 1H, NH); 7.96–6.90 (m, 9H, ArH & benzofuran H); 5.48 (s, 2H, CH_2O).
5i	3287, 3173 (N–H); 1657 (C=O); and 1573, 1488, 1327, 1044 (C=N–N=C–S)	9.82 (s, 1H, NH); 9.21 (s, 1H, NH); 7.98–6.89 (m, 12H, ArH & benzofuran H); 5.44 (s, 2H, CH_2O).

General Procedure for the Synthesis of Compounds 3a–k, 4a, and 5a–i

A solution of 2-benzofuroyl azide (**1**) (0.5 mmol) in toluene (20 mL) was heated at 120°C for 4 h to give benzofuran-2-yl isocyanate **2**, which is not isolated and treated in situ with the respective aromatic amine, 2-amino-5-(benzo-2-furyl)-1,3,4-thiadiazole, or 2-amino-5-aryloxymethylene-1,3,4-thiadiazole under microwave irradiation at 490 W for the time given in Table I. After the completion of the reaction (monitored by TLC using ethyl acetate and petroleum ether (2:3) as eluent), the solvent was removed under reduced pressure and from the residue the products **3a–k**, **4a**, and **5a–i** were isolated by recrystallization from DMF-EtOH. The spectral data of compounds are listed in Table II.

REFERENCES

- [1] A. Bose, M. S. Manhas, S. N. Ganguly, A. H. Sharma, and B. K. Banik, *Synthesis*, **32**, 1579 (2002).
- [2] X. C. Wang, M. G. Wang, Z. J. Quan, and Z. Li, *Synth. Commun.*, **35**, 2881 (2005).
- [3] L. Perrux and A. Loupy, *Tetrahedron*, **57**, 9199 (2001).
- [4] M. Takumi, K. Takanobu, I. Takatoshi, and M. Toshiyuki, *Synth. Commun.*, **30**, 1675 (2000).
- [5] M. J. Gil, M. A. Manu, C. Arteaga, M. Migliaccio, and I. Encio, *Bioorg. Med. Chem. Lett.*, **9**, 2321 (1999).
- [6] E. G. Chalina, L. Chakarova, and D. T. Staneva, *Eur. J. Med. Chem. Chim. Ther.*, **33**, 985 (1998).
- [7] M. Amir, A. Oberoi, and S. Alam, *Indian J. Chem.*, **38B**, 237 (1999).
- [8] H. S. Chen, Z. M. Li, and Y. F. Han, *J. Agric. Food Chem.*, **50**, 3757 (2002).
- [9] X. C. Wang, Z. Li, and Y. X. Da, *Indian J. Chem.*, **40B**, 422 (2001).
- [10] X. P. Hui, L. M. Zhang, Z. Y. Zhang, Q. Wang, and F. Wang, *Indian J. Chem.*, **38B**, 1066 (1999).
- [11] E. E. Chufan, J. C. Pedregosa, O. N. Badini, and L. Bruno-Blanch, *Farmaco*, **54**, 838 (1999).
- [12] A. Arrieta and C. Palomo, *Tetrahedron Lett.*, **22**, 1729 (1981).
- [13] R. A. Batey, V. Santhakumar, C. Yoshina-Ishii, and S. D. Taylor, *Tetrahedron Lett.*, **39**, 6267 (1998).
- [14] X. C. Wang, L. Q. Chai, M. G. Wang, Z. J. Quan, and Z. Li, *Synth. Commun.*, **36**, 645 (2006).
- [15] D. R. Shridhar, C. V. Reddy Sastry K. B. Lal, O. P. Bansal, and S. M. Sondhi, *J. Indian Chem. Soc.*, **55**, 910 (1978).
- [16] S. Tanaka, *J. Am. Chem. Soc.*, **73**, 872 (1951).
- [17] A. A. F. Wasfy, S. A. Nassar, and A. M. F. Eissa, *Indian J. Chem.*, **35B**, 1218 (1996).
- [18] X. C. Wang, Z. Li, Z. J. Quan, X. S. Lu, and R. H. Gou, *Synth. Commun.*, **33**, 2891 (2003).